(19) World Intellectual Property Organization International Bureau





(43) International Publication Date 4 January 2007 (04.01.2007) (10) International Publication Number WO 2007/002543 A2

(51) International Patent Classification: Not classified

1 TOT CIUSE

(21) International Application Number:

PCT/US2006/024717

(22) International Filing Date: 23 June 2006 (23.06.2006)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:

60/693,603 23 June 2005 (23.06.2005) US 60/699,614 15 July 2005 (15.07.2005) US

- (71) Applicant (for all designated States except US): MEDIM-MUME, INC. [US/US]; One Medimmune Way, Gaithersburg, MD 20878 (US).
- (72) Inventors; and
- (75) Inventors/Applicants (for US only): WEI, Ziping [US/US]; 11102 Welland Street, North Potomac, MD 21136 (US). TOUS, Guillermo, I. [US/US]; 146 Barnsfield Court, Gaithersburg, MD 20878 (US). SCHENER-MAN, Mark [US/US]; 1 Cornfield Court, Reisterstown, MD 21136 (US). ALLAN, Christian, B. [US/US]; 4401 Gregg Road, Brookeville, MD 20833 (US).

- (74) Agents: INSOGNA, Anthony, M. et al.; JONES DAY, 222 East 41st Street, New York, NY 10017-6702 (US).
- (81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LV, LY, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RS, RU, SC, SD, SE, SG, SK, SL, SM, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.
- (84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, IIU, IE, IS, IT, LT, LU, LV, MC, NL, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

 without international search report and to be republished upon receipt of that report

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: ANTIBODY FORMULATIONS HAVING OPTIMIZED AGGREGATION AND FRAGMENTATION PROFILES

Conditioned Medium (From Cell Culture) HS50 Chromatography Column Cycle Benzonase Treatment Protein A Chromatography Column Cycle(s) Nanofitration Low pH.Treatment Q Chromatography Column Cycle

(Liquid Formulations of Anti-RSV Antibodies)

(57) Abstract: The present invention provides methods of optimizing the production and purification of antibody formulations that immunospecifically bind to antigens of interest and are suitable for parenteral administration to a subject, which formulations exhibit increased stability due to reduced degradation and aggregation of the antibody component on long term storage. Such methods provide formulations that offer multiple advantages over formulations produced by non-optimized methods including less stringent or more readily available transportation/storage conditions, and less frequent dosing or smaller dosage amounts in the therapeutic, prophylactic and diagnostic use of such formulations. The invention further provides methods of utilizing the formulations of the present invention.

ANTIBODY FORMULATIONS HAVING OPTIMIZED AGGREGATION AND FRAGMENTATION PROFILES

This application claims the benefit under 35 U.S.C. § 119(e) of U.S. Provisional Patent Application No. 60/693,603, filed on June 23, 2005, and U.S. Provisional Patent Application No. 60/699,614, filed on July 15, 2005, each of which is incorporated by reference herein in its entirety.

1. **INTRODUCTION**

[0001] The present invention provides methods of optimizing the production and purification of antibody formulations that immunospecifically bind to antigens of interest and are suitable for parenteral administration to a subject, which formulations exhibit increased stability due to reduced degradation and aggregation of the antibody component on long term storage. Such methods provide formulations that offer multiple advantages over formulations produced by non-optimized methods including less stringent or more readily available transportation/storage conditions, and less frequent dosing or smaller dosage amounts in the therapeutic, prophylactic and diagnostic use of such formulations. The invention further provides methods of utilizing the formulations of the present invention. In a specific embodiment, the invention provides methods of optimizing the production and purification of antibody formulations that immunospecifically bind to RSV antigens, which formulations exhibit increased stability due to reduced degradation and aggregation of the antibody component on long term storage. Such formulations may be used in the diagnostic, therapeutic or prophylactic treatment of RSV infections.

2. **BACKGROUND OF THE INVENTION**

Respiratory Syncytial Virus

[0002] Respiratory infections are common infections of the upper respiratory tract (e.g., nose, ears, sinuses, and throat) and lower respiratory tract (e.g., trachea, bronchial tubes, and lungs). Symptoms of upper respiratory infection include runny or stuffy nose, irritability, restlessness, poor appetite, decreased activity level, coughing, and fever. Viral upper respiratory infections cause and/or are associated with sore throats, colds, croup, and the flu. Clinical manifestations of a lower respiratory infection include shallow coughing that produces sputum in the lungs, fever, and difficulty breathing.

[0003] Respiratory syncytial virus (RSV) is one of the leading causes of respiratory disease worldwide. In the United States, it is responsible for tens of thousands of hospitalizations and thousands of deaths per year (see Black, C.P., Resp. Care 2003 48(3):209-31 for a recent review of the biology and management of RSV). Infants and children are most at risk for serious RSV infections which migrate to the lower respiratory system, resulting in pneumonia or bronchiolitis. In fact, 80% of childhood bronchiolitis cases and 50% of infant pneumonias are attributable to RSV. The virus is so ubiquitous and highly contagious that almost all children have been infected by two years of age. Although infection does not produce lasting immunity, reinfections tend to be less severe so that in older children and healthy adults RSV manifests itself as a cold or flu-like illness affecting the upper and/or lower respiratory system, without progressing to serious lower respiratory tract involvement. However, RSV infections can become serious in elderly or immunocompromised adults. (Evans, A.S., eds., 1989, Viral Infections of Humans. Epidemiology and Control, 3rd ed., Plenum Medical Book, New York at pages 525-544; Falsey, A.R., 1991, Infect. Control Hosp, Epidemiol. 12:602-608: and Garvie et al., 1980, Br. Med. J. 281:1253-1254; Hertz et al., 1989, Medicine 68:269-281).

At present, there is no vaccine against RSV, nor is there any effective treatment. Recent clinical data has failed to support the early promise of the antiviral agent ribavirin, which is the only drug approved for treatment of RSV infection (Black, C.P., Resp. Care 2003 48(3):209-31). Consequently, the American Academy of Pediatrics issued new guidelines suggesting that use of ribavirin be restricted to only the most severe cases (Committee on Infectious Disease, American Academy of Pediatrics. 1996. Pediatrics 97:137-140; Randolph, A.G., and E.E. Wang., 1996, Arch. Pediatr. Adolesc. Med. 150:942-947).

[0005] While a vaccine or effective treatment have proven elusive, some success has been achieved in the area of prevention for infants at high risk of serious upper and/or lower respiratory tract RSV infection. In particular, there are two immunoglobulin-based therapies approved to protect high-risk infants from serious lower respiratory tract RSV infection, RSV-IGIV (RSV-immunoglobulin intravenous, also known as RespiGamTM) and palivizumab (SYNAGIS®). However, neither RSV-IGIV nor palivizumab has been approved for use other than as a prophylactic agent for lower respiratory tract RSV infections.

[0006] RSV is easily spread by physical contact with contaminated secretions. The virus can survive for at least half an hour on hands and for hours on countertops and used tissues. The highly contagious nature of RSV is evident from the risk factors associated with contracting serious infections. One of the greatest risk factors is hospitalization, where in some cases in

excess of 50% of the staff on pediatric wards were found to be infected (Black, C.P., Resp. Care 2003 48(3):209-31). Up to 20% of these adult infections are asymptomatic but still produce substantial shedding of the virus. Other risk factors include attendance at day care centers, crowded living conditions, and the presence of school-age siblings in the home. Importantly, an agent that is effective at clearing the virus from the upper and/or lower respiratory tract is likely to be effective in preventing its transmission. Thus, one promising approach to preventing serious RSV infections is the development of therapies to clear or block the virus from the upper and/or lower respiratory tract.

[0007] Although RSV-IVIG and palivizumab represent significant advances in the prevention of lower respiratory tract RSV infections, neither has demonstrated efficacy at permissible doses against the virus in the upper respiratory tract. In fact, RSV-IVIG failed to clear nasal RSV when administered as a nasal spray in amounts that were effective to clear pulmonary RSV in every animal of the treatment group (Prince et al., U.S. Patent No. 4,800,078, issued January 24, 1989). The interperitoneal route of administration also failed to clear RSV from the upper respiratory tract with the same efficacy as the lower respiratory tract. It has recently been noted that the immune response elicited by upper respiratory tract infections differs from that induced by lower respiratory infections (van Benten I.J. et al., J. Med. Virol. 2003 Oct.;71(2):290-7). Thus, a need exists for the prevention and treatment of upper and/or lower respiratory tract RSV infections.

Otitis Media

Otitis media is an infection or inflammation of the middle ear. This inflammation often begins when infections that cause sore throats, colds, or other respiratory or breathing problems spread to the middle ear. These can be viral or bacterial infections. RSV is the principal virus that has been correlated with otitis media. Seventy-five percent of children experience at least one episode of otitis media by their third birthday. Almost half of these children will have three or more ear infections during their first 3 years. It is estimated that medical costs and lost wages because of otitis media amount to \$5 billion a year in the United States (Gates GA, 1996, Cost-effectiveness considerations in otitis media treatment. Otolaryngol Head Neck Sur. 114 (4): 525-530). Although otitis media is primarily a disease of infants and young children, it can also affect adults.

[0009] Otitis media not only causes severe pain but may result in serious complications if it is not treated. An untreated infection can travel from the middle ear to the nearby parts of the head, including the brain. Although the hearing loss caused by otitis media is usually

temporary, untreated otitis media may lead to permanent hearing impairment. Persistent fluid in the middle ear and chronic otitis media can reduce a child's hearing at a time that is critical for speech and language development. Children who have early hearing impairment from frequent ear infections are likely to have speech and language disabilities.

[0010] Although many physicians recommend the use of antibiotics for the treatment of ear infections, antibiotic resistance has become an important problem in effective treatment of the disease. Further, new therapies are needed to prevent or treat viral infections that are associated with otitis media, particularly RSV.

Asthma and Reactive Airway Disease (RAD)

[0011] About 12 million people in the U.S. have asthma and it is the leading cause of hospitalization for children. *The Merck Manual of Diagnosis and Therapy* (17th ed., 1999).

Asthma is an inflammatory disease of the lung that is characterized by airway hyperresponsiveness ("AHR"), bronchoconstriction (*i.e.*, wheezing), eosinophilic inflammation, mucus hypersecretion, subepithelial fibrosis, and elevated IgE levels. Asthmatic attacks can be triggered by environmental triggers (*e.g.* acarids, insects, animals (*e.g.*, cats, dogs, rabbits, mice, rats, hamsters, guinea pigs, mice, rats, and birds), fungi, air pollutants (*e.g.*, tobacco smoke), irritant gases, fumes, vapors, aerosols, chemicals, or pollen), exercise, or cold air. The cause(s) of asthma is unknown. However, it has been speculated that family history of asthma (London *et al.*, 2001, Epidemiology 12(5):577-83), early exposure to allergens, such as dust mites, tobacco smoke, and cockroaches (Melen *et al.*, 2001, 56(7):646-52), and respiratory infections (Wenzel *et al.*, 2002, Am J Med, 112(8):672-33 and Lin *et al.*, 2001, J Microbiol Immuno Infect, 34(4):259-64), such as RSV, may increase the risk of developing asthma. A review of asthma, including risk factors, animal models, and inflammatory markers can be found in O'Byrne and Postma (1999), Am. J. Crit. Care. Med. 159:S41-S66, which is incorporated herein by reference in its entirety.

[0013] Current therapies are mainly aimed at managing asthma and include the administration of β-adrenergic drugs (e.g. epinephrine and isoproterenol), theophylline, anticholinergic drugs (e.g., atropine and ipratorpium bromide), corticosteroids, and leukotriene inhibitors. These therapies are associated with side effects such as drug interactions, dry mouth, blurred vision, growth suppression in children, and osteoporosis in menopausal women. Cromolyn and nedocromil are administered prophylatically to inhibit mediator release from inflammatory cells, reduce airway hyperresponsiveness, and block responses to allergens.

However, there are no current therapies available that prevent the development of asthma in subjects at increased risk of developing asthma. Thus, new therapies with fewer side effects and better prophylactic and/or therapeutic efficacy are needed for asthma.

[0014] Reactive airway disease is a broader (and often times synonymous) characterization for asthma-like symptoms, and is generally characterized by chronic cough, sputum production, wheezing or dyspenea.

Wheezing

[0015] Wheezing (also known as sibilant rhonchi) is generally characterized by a noise made by air flowing through narrowed breathing tubes, especially the smaller, tight airways located deep within the lung. It is a common symptom of RSV infection, and secondary RSV conditions such as asthma and brochiolitis. The clinical importance of wheezing is that it is an indicator of airway narrowing, and it may indicate difficulty breathing.

[0016] Wheezing is most obvious when exhaling (breathing out), but may be present during either inspiration (breathing in) or exhalation. Wheezing most often comes from the small bronchial tubes (breathing tubes deep in the chest), but it may originate if larger airways are obstructed.

[0017] Citation or discussion of a reference herein shall not be construed as an admission that such is prior art to the present invention.

3. <u>SUMMARY OF THE INVENTION</u>

[0018] The present invention is based upon the inventors' use of sensitive analytical techniques, such as analytical ultracentrifugation (AUC), size exclusion chromatography (SEC), Liquid Chromatography Mass Spectrometry (LC-MS) or particle counter analysis to analyze the fragmentation and aggregation profiles of formulations of full-length IgG1 monoclonal antibodies, particularly those that have been recombinantly expressed in myeloma cells, such as, but not limited to, NS0 cells. Thus, the present invention provides antibody formulations having fragmentation and aggregation profiles that are improved (*i.e.*, have decreased total fragmentation and/or aggregation or have decreased amounts of certain types of fragments or aggregates or have reduced rates of aggregation or fragmentation) as compared to prior antibody formulations.

[0019] In a particular embodiment, the invention provides an antibody formulation comprising a full length IgG1 antibody, preferably specific for a therapeutic or prophylactic target, wherein no more than 0.5 % of the total protein fraction (in certain embodiments, however, at least 0.1% of the total protein fraction or is below detectable levels) of said formulation comprises (or, in other embodiments, consists of as impurities or as fragments to detectable levels) antibody type I fragments. In other embodiments, no more than 0.5% of the total protein fraction (and, in certain embodiments, at least 0.1% of the total protein fraction or is below detectable levels) of said formulation comprises (or, in other embodiments, consists of as impurities or as fragments to detectable levels) antibody type I fragments and antibody type II fragments. Preferably, the antibody type I fragments comprise one or more C-terminal portions of the heavy chain of the antibody, which heavy chain C-terminal portion has a molecular weight of about 25.6 kD, about 25.7 kD, about 25.8 kD, about 26.0 kD, or about 26.1 kD as determined by Liquid Chromatography Mass Spectrometry (LC-MS) analysis of samples of the antibody that have been deglycosylated, reduced and alkylated. Morevoer, the antibody type II fragments comprise one or more N-terminal portions of the heavy chain of the antibody, which heavy chain N-terminal portion has a molecular weight of about 24.4 kD, about 24.6 kD, about 24.7 kD, about 24.9 kD, or about 25.1 kD as determined by LC-MS analysis of samples of the antibody that have been deglycosylated, reduced and alkylated. In addition, the antibody type I fragments may comprise one or more C-terminal portions of the heavy chain, which heavy chain C-terminal portion comprises amino acid residues 223-449 of the IgG1 heavy chain (according to Kabat numbering), amino acid residues 224-449 of the IgG1 heavy chain, amino acid residues 225-449 of the IgG1 heavy chain, amino acid residues 226-449 of the IgG1 heavy chain, amino acid residues 227-449 of the IgG1 heavy chain, amino acid residues 228-449 of the IgG1 heavy chain and amino acid residues 229-449 of the IgG1 heavy chain and the antibody type II fragments comprise one or more heavy chain N-terminal portions which comprises amino acid residues 1-222 of the IgG1 heavy chain, amino acid residues 1-223 of the IgG1 heavy chain, amino acid residues 1-224 of the IgG1 heavy chain, amino acid residues 1-225 of the IgG1 heavy chain, amino acid residues 1-226 of the IgG1 heavy chain, amino acid residues 1-227 of the IgG1 heavy chain or amino acid residues 1-228 of the IgG1 heavy chain. In certain embodiments, the antibody formulation does not contain detectable levels of any other types of fragments. In certain embodiments, the antibody formulation contains one, two, three, four, five, six or seven of the type I fragments and/or contains one, two, three, four, five, six or seven of the type II fragments.

[0020] In particular embodiments, the formulations of the invention comprise (or consists of as the aggregate fraction) a particle profile of less than about 3.4 E +5 particles/ml of diameter 2-4 μm , less than about 4.0 E +4 particles/ml of diameter 4-10 μm , less than about 4.2 E +3 particles/ml of diameter 10-20 μm , less than about 5.0 E +2 particles/ml of diameter 20-30 μm , less than about 7.5 E +1 particles/ml of diameter 30-40 μm , and less than about 9.4 particles/ml of diameter 40-60 μm as determined by a particle multisizer. In certain embodiments, the formulation contains no detectable particles greater than 40 μm , or greater than 30 μm . In other embodiments, the formulations of the invention have a turbidity value of a degassed sample of said formulation of about 6.4 NTU (in certain embodiments from 4-8 NTU, in other embodiments less than 10 NTU, less than 8 NTU, less than 7 NTU, or less than 6.5 NTU).

[0021] The antibody formulations of the invention may likewise have a combination of one or more of the above parameters of fragmentation and aggregation.

[0022] The antibody formulations of the invention are preferably at least 10 mg/ml antibody, more preferably, 15 mg/ml, 25 mg/ml, 50 mg/ml, 75 mg/ml, 100 mg/ml, 150 mg/ml or 200 mg/ml. The antibody in the antibody formulations of the invention may be any antibody that has a therapeutic, prophylactic or diagnostic utility. In preferred embodiments, the antibody in the formulations of the invention is specific for RSV and, in a specific embodiment, is not palivizumab. In more specific and preferred embodiments, the anti-RSV antigen binds to the F protein of RSV, and, in particular embodiments, the RSV antigen comprises or even consists of the F protein epitope NSELLSLINDMPITNDQKKLMSNN (SEQ ID NO:337). In other embodiments, the antibody is one of the antibodies listed in Table 2, preferably is the antibody A4B4L1FR-S28R or competes for binding with one of the antibodies listed in Table 2, preferably A4B4L1FR-S28R.

[0023] The antibody formulations of the invention preferably maintain improved aggregation and fragmentation profiles upon storage, for example, for extended periods (for example, but not limited to 6 months, 1 year, 2 years, 3 years or 5 years) at room temperature or 4°C or for periods (such as, but not limited to 1 week, 2 weeks, 3 weeks, 1 month, 2 months, 3 months, 6 months or 1 year) at elevated temperatures such as 38°C-42°C. Such formulations may be at pH 5-7, preferably at pH 6.0. Thus, in a particular embodiment, an antibody formulation of the invention comprising a full length IgG₁ antibody, upon storage at 38-42°C, pH 6.0 for 1 month, 6 months, 9 months or 14 months, comprises or, alternatively consists (other than the full length IgG1 antibody or as the fragment fraction) one or more antibody type I

fragments. In another particular embodiment, an antibody formulation of the invention comprising a full length IgG_1 antibody, upon storage at $38-42^{\circ}C$, pH 6.0 for 1 month, 6 months, 9 months or 14 months, comprises or, alternatively consists (other than the full length IgG_1 antibody or as the fragment fraction) one or more antibody type I fragments and one or more antibody type II fragments. Upon storage, the level of fragments as a percentage of the total amount of protein is preferably less than 0.5% and, in certain embodiments is at least 0.1% or is below detectable levels of fragments.

Additionally, during storage, such formulations preferably exhibit constant aggregation and fragmentation rates at temperatures, such as, but not limited to, 0-4°C, 10-15°, 20-24°C room temperature, or elevated temperatures 38-42°C, and extended periods, such as, but not limited to, two weeks, one month, six months, one year, three years or five years. In certain embodiments, the antibody formulation does not contain detectable levels of any other types of fragments. Thus, in a particular embodiment, an antibody formulation fo the invention comprising a full length IgG1 will increase in aggregate percentage relative to total protein, by 0.2%/month - 0.35%/month at 20-24°C and preferably by not more than 0.02%/month at 4°C. In a further embodiment, an antibody formulation of the invention comprising a full length IgG1 will not increase in fragment percentage, relative to total protein, by more than 0.015%/month - 0.03%/month at 20-24°C and preferably by not more than 0.00%/month at 4°C. In certain embodiments, the antibody formulation contains one, two, three, four, five or six or the type I fragments and/or contains one, two, three, four, five, six or seven of the type II fragments.

In particular embodiments, after storage, the formulations of the invention comprise (or consists of as the aggregate fraction) a particle profile of less than about 3.4 E + 5 particles/ml of diameter $2-4 \mu m$, less than about 4.0 E + 4 particles/ml of diameter $4-10 \mu m$, less than about 4.2 E + 3 particles/ml of diameter $10-20 \mu m$, less than about 5.0 E + 2 particles/ml of diameter $20-30 \mu m$, less than about 7.5 E + 1 particles/ml of diameter $30-40 \mu m$, and less than about 9.4 particles/ml of diameter $40-60 \mu m$ as determined by a particle multisizer. In certain embodiments, the formulation contains no detectable particles greater than $40 \mu m$, or greater than $30 \mu m$. In other embodiments, the formulations of the invention, after storage, have a turbidity value of a degassed sample of said formulation of about 6.4 NTU (in certain embodiments from 4-8 NTU, in other embodiments less than 10 NTU, less than 8 NTU, less than 7 NTU, or less than 6.5 NTU).

[0026] The antibody formulations of the invention, after storage, may likewise have a combination of one or more of the above parameters of fragmentation and aggregation.

[0027] Other aspects of the invention provide for methods of optimizing a particular antibody formulation for the fragmentation and aggregation parameters set forth above. Such methods comprise production, purification and formulation of the antibody and monitoring at one or more steps, or of the final formulation, for the levels of fragmentation and/or aggregation using methods such as, but not limited to AUC, SEC, LC-MS or particle multisizing, and then varying one or more parameters of one or more steps of the production, purification and/or formulation process or the formulation itself and evaluating whether varying the parameter reduces the level of fragmentation and/or aggregation. By such screening and monitoring steps, the method of the invention may be used to optimize an antibody formulation. Such parameters include, the termperature at which one or more steps is carried out, the reduction in or elimination of freeze/thaw cycles of the antibody, introduction of filtration steps, such as ultrafiltration, addition or change in one or more column chromatography steps, change in pH, etc.

[0028] The invention provides an antibody comprising a Fab fragment, which immunospecifically binds to an RSV antigen (e.g., the F protein epitope NSELLSLINDMPITNDQKKLMSNN (SEQ ID NO:337)), wherein the Tm of the Fab fragment is at least about 87 °C, and wherein said antibody is not any of palivizumab, AFFF, P12f2, P12f4, P11d4, Ale9, A12a6, A13c4, A17d4, A4B4, A8c7, 1X-493L1FR, H3-3F4, M3H9, Y10H6, DG, AFFF(1), 6H8, L1-7E5, L2-15B10, A13a11, A1h5, A4B4(1), A4B4L1FR-S28R (motavizumab), A4B4-F52S, A17d4(1), A3e2, A14a4, A16b4, A17b5, A17f5, and A17h4. In a specific embodiment, the Fab in such an antibody is different from the Fab of palivizumab. In another embodiment, such an antibody comprises a VH or VL domain that is different from the VH or VL domain of palivizumab. In preferred embodiment, the Tm of the Fab fragment is at least about 90 °C or at least about 93 °C. In another preferred embodiment, the pI of the antibody is between about 8.5 to 9.5 or between about 9.0 to 9.5.

[0029] In another specific embodiment, the antibody comprises a VH domain of the antibody A4B4L1FR-S28R (SEQ ID NO:48). In still another embodiment, the antibody comprises a VL domain of the antibody A4B4L1FR-S28R (SEQ ID NO:11). In still another embodiment, said Fab is the Fab of antibody A4B4L1FR-S28R.

[0030] The invention also provides an antibody formulation comprising the above described antibody, said formulation having a viscosity of less than 10.00 cP at any temperature in the range of 1 to 26 °C.

[0031] The invention also provides an antibody formulation comprising the above described antibody, said formulation having an aggregation rate of less than 15% per day at any temperature in the range of 38 to 42 °C.

[0032] The invention also provides a method of preventing, treating, or ameliorating one or more symptoms associated with a RSV infection in a subject, e.g., otitis media, asthma, and wheezing, said method comprising administering a prophylactically or therapeutically effective amount of an antibody formulation comprising such antibody. In one embodiment, the formulation is administered parenterally, intramuscularly, intravenously, subcutaneously or intranasally.

[0033] The invention also provides an antibody formulation comprising a full length IgG₁ antibody, which immunospecifically binds to an RSV antigen, said formulation having a viscosity of less than 10.00 cP at any temperature in the range of 1 to 26 °C. The invention also provides an antibody formulation comprising any such antibody, said formulation having an aggregration rate of less than 15% per day at any temperature in the range of 38 to 42 °C. In one embodiment, the antibody is not palivizumab. In another embodiment, the antibody is not any of AFFF, P12f2, P12f4, P11d4, Ale9, A12a6, A13c4, A17d4, A4B4, A8c7, 1X-493L1FR, H3-3F4, M3H9, Y10H6, DG, AFFF(1), 6H8, L1-7E5, L2-15B10, A13a11, A1h5, A4B4(1), A4B4L1FR-S28R (motavizumab), A4B4-F52S, A17d4(1), A3e2, A14a4, A16b4, A17b5, A17f5, and A17h4.

3.1 TERMINOLOGY

[0034] In the context of a polypeptide, the term "analog" as used herein refers to a polypeptide that possesses a similar or identical function as a RSV polypeptide, a fragment of a RSV polypeptide, or an antibody but does not necessarily comprise a similar or identical amino acid sequence of a RSV polypeptide, a fragment of a RSV polypeptide, or an antibody, or possess a similar or identical structure of a RSV polypeptide, a fragment of a RSV polypeptide, or an antibody. A polypeptide that has a similar amino acid sequence refers to a polypeptide that satisfies at least one of the following: (a) a polypeptide having an amino acid sequence that is at least 30%, at least 35%, at least 40%, at least 45%, at least 50%, at least 55%, at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 99% identical to the amino acid sequence of a RSV polypeptide, a fragment of a RSV polypeptide, or an antibody described herein; (b) a polypeptide encoded by a nucleotide sequence that hybridizes under stringent conditions to a nucleotide sequence encoding a RSV polypeptide, a fragment of a RSV polypeptide, or an antibody described herein of at least 5

amino acid residues, at least 10 amino acid residues, at least 15 amino acid residues, at least 20 amino acid residues, at least 25 amino acid residues, at least 40 amino acid residues, at least 50 amino acid residues, at least 60 amino residues, at least 70 amino acid residues, at least 80 amino acid residues, at least 90 amino acid residues, at least 100 amino acid residues, at least 125 amino acid residues, or at least 150 amino acid residues; and (c) a polypeptide encoded by a nucleotide sequence that is at least 30%, at least 35%, at least 40%, at least 45%, at least 55%, at least 55%, at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95% or at least 99% identical to the nucleotide sequence encoding a RSV polypeptide, a fragment of a RSV polypeptide, or an antibody described herein. A polypeptide with similar structure to a RSV polypeptide, a fragment of a RSV polypeptide, or an antibody described herein refers to a polypeptide that has a similar secondary, tertiary or quaternary structure of a RSV polypeptide, a fragment of a RSV, or an antibody described herein. The structure of a polypeptide can determined by methods known to those skilled in the art, including but not limited to, X-ray crystallography, nuclear magnetic resonance, and crystallographic electron microscopy.

[0035] To determine the percent identity of two amino acid sequences or of two nucleic acid sequences, the sequences are aligned for optimal comparison purposes (e.g., gaps can be introduced in the sequence of a first amino acid or nucleic acid sequence for optimal alignment with a second amino acid or nucleic acid sequence). The amino acid residues or nucleotides at corresponding amino acid positions or nucleotide positions are then compared. When a position in the first sequence is occupied by the same amino acid residue or nucleotide as the corresponding position in the second sequence, then the molecules are identical at that position. The percent identity between the two sequences is a function of the number of identical positions shared by the sequences (i.e., % identity = number of identical overlapping positions/total number of positions X 100%). In one embodiment, the two sequences are the same length.

[0036] The determination of percent identity between two sequences can also be accomplished using a mathematical algorithm. A preferred, non limiting example of a mathematical algorithm utilized for the comparison of two sequences is the algorithm of Karlin and Altschul, 1990, Proc. Natl. Acad. Sci. U.S.A. 87:2264 2268, modified as in Karlin and Altschul, 1993, Proc. Natl. Acad. Sci. U.S.A. 90:5873 5877. Such an algorithm is incorporated into the NBLAST and XBLAST programs of Altschul et al., 1990, J. Mol. Biol. 215:403. BLAST nucleotide searches can be performed with the NBLAST nucleotide program parameters

set, e.g., for score=100, wordlength=12 to obtain nucleotide sequences homologous to a nucleic acid molecules of the present invention. BLAST protein searches can be performed with the XBLAST program parameters set, e.g., to score 50, wordlength=3 to obtain amino acid sequences homologous to a protein molecule of the present invention. To obtain gapped alignments for comparison purposes, Gapped BLAST can be utilized as described in Altschul et al., 1997, Nucleic Acids Res. 25:3389 3402. Alternatively, PSI BLAST can be used to perform an iterated search which detects distant relationships between molecules (Id.). When utilizing BLAST, Gapped BLAST, and PSI Blast programs, the default parameters of the respective programs (e.g., of XBLAST and NBLAST) can be used (see, e.g., http://www.ncbi.nlm.nih.gov). Another preferred, non limiting example of a mathematical algorithm utilized for the comparison of sequences is the algorithm of Myers and Miller, 1988, CABIOS 4:11 17. Such an algorithm is incorporated in the ALIGN program (version 2.0) which is part of the GCG sequence alignment software package. When utilizing the ALIGN program

[0037] The percent identity between two sequences can be determined using techniques similar to those described above, with or without allowing gaps. In calculating percent identity, typically only exact matches are counted.

for comparing amino acid sequences, a PAM120 weight residue table, a gap length penalty of

12, and a gap penalty of 4 can be used.

[0038] The terms "antibodies that immunospecifically bind to a RSV antigen" and analogous terms as used herein refers to antibodies that specifically bind to a RSV polypeptide or a fragment of a RSV polypeptide and do not non-specifically bind to other polypeptides. Antibodies that immunospecifically bind to a RSV polypeptide or fragment thereof may have cross-reactivity with other antigens. Preferably, antibodies that immunospecifically bind to a RSV polypeptide or fragment thereof do not cross-react with other antigens. Antibodies that immunospecifically bind to a RSV polypeptide can be identified, for example, by immunoassays or other techniques known to those of skill in the art.

[0039] Antibodies of the invention include, but are not limited to, synthetic antibodies, monoclonal antibodies, recombinantly produced antibodies, multispecific antibodies (including bi-specific antibodies), human antibodies, humanized antibodies, chimeric antibodies, intrabodies, single-chain Fvs (scFv) (e.g., including monospecific and bi-specific, etc.), Fab fragments, F(ab') fragments, disulfide-linked Fvs (sdFv), anti-idiotypic (anti-Id) antibodies, and epitope-binding fragments of any of the above. In particular, antibodies of the present invention include immunoglobulin molecules and immunologically active portions of immunoglobulin

molecules, *i.e.*, molecules that contain an antigen-binding site that immunospecifically binds to a RSV antigen (preferably, a RSV F antigen) (*e.g.*, one or more complementarity determining regions (CDRs) of an anti-RSV antibody). The antibodies of the invention can be of any type (*e.g.*, IgG, IgE, IgM, IgD, IgA and IgY), class (*e.g.*, IgG₁, IgG₂, IgG₃, IgG₄, IgA₁ and IgA₂) or a subclass of immunoglobulin molecule.

[0040] As used herein, the term "analogue" in the context of a non-proteinaceous analog refers to a second organic or inorganic molecule which possess a similar or identical function as a first organic or inorganic molecule and is structurally similar to the first organic or inorganic molecule.

[0041] The term "antibody fragment" as used herein refers to a fragment of an antibody that immunospecifically binds to a RSV antigen. Antibody fragments may be generated by any technique known to one of skill in the art and by proteolytic or non-proteolytic cleavage. For example, Fab and F(ab')₂ fragments may be produced by proteolytic cleavage of immunoglobulin molecules, using enzymes such as papain (to produce Fab fragments) or pepsin (to produce F(ab')₂ fragments). F(ab')₂ fragments contain the complete light chain, and the variable region, the CH1 region and the hinge region of the heavy chain. Antibody fragments can be also produced by recombinant DNA technologies. Antibody fragments may be one or more complementarity determining regions (CDRs) of antibodies.

[0042] The term "antibody type I fragment" as used herein refers to a multimeric protein comprising a full length antibody light chain, a full length antibody heavy chain and a C-terminal portion of an antibody heavy chain that, in human IgG₁ immunoglobulins, has an N-terminus at cysteine 223, aspartic acid 224, lysine 225, threonine 226, histidine 227, threonine 228 or cysteine 229 and a C-terminus at lysine 449. Amino acid numbering for the constant domain is given according to the Kabat EU numbering scheme (Kabat, E. A., T. T. Wu, H. M. Perry, K. S. Gottesman, and Foeller. 1991. Sequences of Proteins of Immunological Interest, U.S. Public Health Service, National Institutes of Health, Washington, D.C., which is incorporated herein by reference), unless otherwise indicated. In a specific embodiment, the full length antibody light chain, full length antibody heavy chain and C-terminal portion of an antibody heavy chain are linked by disulfide bonds as depicted in FIG. 14. In another specific embodiment, the type I fragment is capable of immunospecifically binding to an antigen of interest.

[0043] The term "antibody type II fragment" as used herein refers to a peptide, polypeptide, or protein comprising an antibody light chain and an N-terminal portion of an antibody heavy chain that, in human IgG₁ immunoglobulins, has a C-terminus at serine 222, cysteine 223, aspartic acid 224, lysine 225, threonine 226, histidine 227 or threonine 228 and an N-terminus at glycine 1. In a specific embodiment, the full length antibody light chain and N-terminal portion of an antibody heavy chain are linked by disulfide bonds as depicted in FIG. 14. In another specific embodiment, the type II fragment is capable of immunospecifically binding to an antigen of interest.

In the context of a polypeptide, the term "derivative" as used herein refers to a [0044] polypeptide that comprises an amino acid sequence of a RSV polypeptide, a fragment of a RSV polypeptide, or an antibody that immunospecifically binds to a RSV polypeptide which has been altered by the introduction of amino acid residue substitutions, deletions or additions. The term "derivative" as used herein also refers to a RSV polypeptide, a fragment of a RSV polypeptide, or an antibody that immunospecifically binds to a RSV polypeptide which has been modified, i.e., by the covalent attachment of any type of molecule to the polypeptide. For example, but not by way of limitation, a RSV polypeptide, a fragment of a RSV polypeptide, or an antibody may be modified, e.g., by glycosylation, acetylation, pegylation, phosphorylation, amidation, derivatization by known protecting/blocking groups, proteolytic cleavage, linkage to a cellular ligand or other protein, etc. A derivative of a RSV polypeptide, a fragment of a RSV polypeptide, or an antibody may be modified by chemical modifications using techniques known to those of skill in the art, including, but not limited to specific chemical cleavage, acetylation, formylation, metabolic synthesis of tunicamycin, etc. Further, a derivative of a RSV polypeptide, a fragment of a RSV polypeptide, or an antibody may contain one or more nonclassical amino acids. A polypeptide derivative possesses a similar or identical function as a RSV polypeptide, a fragment of a RSV polypeptide, or an antibody described herein.

[0045] As used herein, the term "derivative" in the context of a non-proteinaceous derivative refers to a second organic or inorganic molecule that is formed based upon the structure of a first organic or inorganic molecule. A derivative of an organic molecule includes, but is not limited to, a molecule modified, e.g., by the addition or deletion of a hydroxyl, methyl, ethyl, carboxyl or amine group. An organic molecule may also be esterified, alkylated and/or phosphorylated.

[0046] The term "effective amount" as used herein refers to the amount of a therapy (e.g., an antibody of the invention) which is sufficient to reduce and/or ameliorate the severity

and/or duration of a disease or disorder. For example, the "effective amount" of an anti RSV antibody is that which is sufficient to reduce and/or ameliorate the severity and/or duration of an upper and/or lower respiratory tract RSV infection, otitis media, and/or a symptom or respiratory condition relating thereto (including, but not limited to, asthma, wheezing, RAD, or a combination thereof), prevent the advancement or progression of the upper and/or lower respiratory tract RSV infection, otitis media and/or a symptom or respiratory condition relating thereto (e.g., prevent the progression of an upper respiratory tract RSV infection to a lower respiratory tract RSV infection), prevent the recurrence, development, or onset of an upper and/or lower respiratory tract RSV infection, otitis media, and/or a symptom or respiratory condition relating thereto (including, but not limited to, asthma, wheezing, RAD, or a combination thereof), and/or enhance/improve the prophylactic or therapeutic effect(s) of another therapy (e.g., a therapy other than an antibody of the invention). Non-limiting examples of effective amounts of an antibody of the invention are provided in Section 5.3, infra. With respect to the treatment of a RSV infection, a therapeutically effective amount refers to the amount of a therapeutic agent sufficient to reduce or inhibit the replication of a virus, inhibit or reduce the infection of cell with the virus, inhibit or reduce the production of the viral particles, inhibit or reduce the release of viral particles, inhibit or reduce the spread of the virus to other tissues or subjects, or ameliorate one or more symptoms associated with the infection. In a specific embodiment, a therapeutically effective amount of a therapeutic agent reduces one or more of the following steps of a RSV life cycle: the docking of the virus particle to a cell, the introduction of viral genetic information into a cell, the expression of viral proteins, the production of new virus particles and the release of virus particles from a cell by at least 5%, preferably at least 10%, at least 15%, at least 20%, at least 25%, at least 30%, at least 35%, at least 40%, at least 45%, at least 50%, at least 55%, at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, or at least 100%. In another specific embodiment, a therapeutically effective amount of a therapeutic agent reduces the replication, multiplication or spread of a virus by at least 5%, preferably at least 10%, at least 15%, at least 20%, at least 25%, at least 30%, at least 35%, at least 40%, at least 45%, at least 50%, at least 55%, at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, or at least 100%.

[0047] The term "effective neutralizing titer" of an anti-RSV antibody as used herein refers to the amount of antibody which corresponds to the amount present in the serum of animals (human or cotton rat) that has been shown to be either clinically efficacious (in humans)

or to reduce virus by 99% in, for example, cotton rats. The 99% reduction is defined by a specific challenge of, e.g., 10^3 pfu, 10^4 pfu, 10^5 pfu, 10^6 pfu, 10^7 pfu, 10^8 pfu, or 10^9 pfu of RSV.

[0048] The term "elderly" as used herein refers to a human subject who is age 65 or older.

[0049] The term "epitopes" as used herein refers to fragments of a polypeptide having antigenic or immunogenic activity in an animal, preferably a mammal, and most preferably in a human. An epitope having immunogenic activity is a fragment of a polypeptide that elicits an antibody response in an animal. An epitope having antigenic activity is a fragment of a polypeptide to which an antibody immunospecifically binds as determined by any method well known in the art, for example, by the immunoassays described herein. Antigenic epitopes need not necessarily be immunogenic.

[0050] The term "excipients" as used herein refers to inert substances which are commonly used as a diluent, vehicle, preservatives, binders, or stabilizing agent for drugs and includes, but not limited to, proteins (e.g., serum albumin, etc.), amino acids (e.g., aspartic acid, glutamic acid, lysine, arginine, glycine, histidine, etc.), fatty acids and phospholipids (e.g., alkyl sulfonates, caprylate, etc.), surfactants (e.g., SDS, polysorbate, nonionic surfactant, etc.), saccharides (e.g., sucrose, maltose, trehalose, etc.) and polyols (e.g., mannitol, sorbitol, etc.). Also see Remington's Pharmaceutical Sciences (by Joseph P. Remington, 18th ed., Mack Publishing Co., Easton, PA), which is hereby incorporated in its entirety.

In the term "fragment" as used herein refers to a peptide or polypeptide comprising an amino acid sequence of at least 5 contiguous amino acid residues, at least 10 contiguous amino acid residues, at least 20 contiguous amino acid residues, at least 20 contiguous amino acid residues, at least 25 contiguous amino acid residues, at least 40 contiguous amino acid residues, at least 50 contiguous amino acid residues, at least 60 contiguous amino residues, at least 70 contiguous amino acid residues, at least 80 contiguous amino acid residues, at least 90 contiguous amino acid residues, at least 125 contiguous amino acid residues, at least 150 contiguous amino acid residues, at least 175 contiguous amino acid residues, at least 200 contiguous amino acid residues, or at least 250 contiguous amino acid residues of the amino acid sequence of a polypeptide or an antibody that immunospecifically binds to a polypeptide. In a specific embodiment, a fragment of a polypeptide or an antibody of that immunospecifically binds to an antigen retains at least 1, at least 2, or at least 3 functions of the polypeptide or antibody.

[0052] The term "fusion protein" as used herein refers to a polypeptide that comprises an amino acid sequence of an antibody and an amino acid sequence of a heterologous polypeptide or protein (i.e., a polypeptide or protein not normally a part of the antibody (e.g., a non-anti-RSV antigen antibody)).

[0053] The terms "high concentration" and "concentrated antibody" as used herein refer to a concentration of 50 mg/ml or higher, preferably 95 mg/ml or higher of an antibody or antigen-binding fragment thereof in an antibody formulation.

[0054] The term "high potency" as used herein refers to antibodies that exhibit high potency as determined in various assays for biological activity (e.g., neutralization of RSV) such as those described herein. For example, high potency antibodies of the invention have an IC₅₀ value less than 5 nM, less than 4 nM, less than 3 nM, less than 2 nM, less than 1.75 nM, less than 1.5 nM, less than 1.25 nM, less than 1 nM, less than 0.75 nM, less than 0.5 nM, less than 0.25 nM, less than 0.1 nM, less than 0.05 nM, less than 0.025 nM, or less than 0.01 nM, as measured by a microneutralization assay described herein. Further, high potency anti-RSV antibodies of the invention result in at least a 75%, preferably at least a 95% and more preferably a 99% lower RSV titer in a cotton rat 5 days after challenge with 10⁵ pfu relative to a cotton rat not administered said antibodies. In certain embodiments of the invention, high potency anti-RSV antibodies of the present invention exhibit a high affinity and/or high avidity for one or more RSV antigens (e.g., antibodies having an affinity of at least 2 X 108 M⁻¹, preferably at least $2.5 \times 10^8 \text{ M}^{-1}$, at least $5 \times 10^8 \text{ M}^{-1}$, at least 10^9 M^{-1} , at least 10^9 M^{-1} , at least 10^{10} M^{-1} , at least $5 \times 10^{10} \, \text{M}^{-1}$, at least $10^{11} \, \text{M}^{-1}$, at least $5 \times 10^{11} \, \text{M}^{-1}$, at least $10^{12} \, \text{M}^{-1}$, or at least $5 \times 10^{12} \, \text{M}^{-1}$ for one or more RSV antigens).

[0055] The term "host" as used herein refers to an animal, preferably a mammal, and most preferably a human.

[0056] The term "host cell" as used herein refers to the particular subject cell transfected with a nucleic acid molecule and the progeny or potential progeny of such a cell. Progeny of such a cell may not be identical to the parent cell transfected with the nucleic acid molecule due to mutations or environmental influences that may occur in succeeding generations or integration of the nucleic acid molecule into the host cell genome.

[0057] The term "human infant" as used herein refers to a human less than 24 months, preferably less than 16 months, less than 12 months, less than 6 months, less than 3 months, less than 2 months, or less than 1 month of age.

[0058] The term "human infant born prematurely" as used herein refers to a human born at less than 40 weeks gestational age, preferably less than 35 weeks gestational age, who is less than 6 months old, preferably less than 3 months old, more preferably less than 2 months old, and most preferably less than 1 month old.

As used herein, the term "in combination" refers to the use of more than one therapy. The use of the term "in combination" does not restrict the order in which therapies are administered to a subject with an infection. A first therapy can be administered before (*e.g.*, 1 minute, 45 minutes, 30 minutes, 45 minutes, 1 hour, 2 hours, 4 hours, 6 hours, 12 hours, 24 hours, 48 hours, 72 hours, 96 hours, 1 week, 2 weeks, 3 weeks, 4 weeks, 5 weeks, 6 weeks, 8 weeks, or 12 weeks), concurrently, or after (*e.g.*, 1 minute, 45 minutes, 30 minutes, 45 minutes, 1 hour, 2 hours, 4 hours, 6 hours, 12 hours, 24 hours, 48 hours, 72 hours, 96 hours, 1 week, 2 weeks, 3 weeks, 4 weeks, 5 weeks, 6 weeks, 8 weeks, or 12 weeks) the administration of a second therapy to a subject which had, has, or is susceptible to a disease or disorder. Any additional therapy can be administered in any order with the other additional therapies. In certain embodiments, the antibodies of the invention can be administered in combination with one or more non-antibody therapies. Non-limiting examples of therapies that can be administered in combination with an antibody of the invention include analgesic agents, anesthetic agents, antibiotics, or immunomodulatory agents.

[0060] As used herein, the term "infection" refers to all stages of RSV's life cycle in a host (including, but not limited to the invasion by and replication of RSV in a cell or body tissue), and the pathological state resulting from the invasion by and replication of a RSV. The invasion by and multiplication of a RSV includes, but is not limited to, the following steps: the docking of the RSV particle to a cell, the introduction of viral genetic information into a cell, the expression of RSV proteins, the production of new RSV particles and the release of RSV particles from a cell.

[0061] The term "inorganic salt" as used herein refers to any compounds containing no carbon that result from replacement of part or all of the acid hydrogen or an acid by a metal or a group acting like a metal and are often used as a tonicity adjusting compound in pharmaceutical compositions and preparations of biological materials. The most common inorganic salts are NaCl, KCl, NaH₂PO₄, etc.

[0062] An "isolated" or "purified" antibody is substantially free of cellular material or other contaminating proteins from the cell or tissue source from which the protein is derived, or

substantially free of chemical precursors or other chemicals when chemically synthesized. The language "substantially free of cellular material" includes preparations of an antibody in which the antibody is separated from cellular components of the cells from which it is isolated or recombinantly produced. Thus, an antibody that is substantially free of cellular material includes preparations of antibody having less than about 30%, 20%, 10%, or 5% (by dry weight) of heterologous protein (also referred to herein as a "contaminating protein"). When the antibody is recombinantly produced, it is also preferably substantially free of culture medium, *i.e.*, culture medium represents less than about 20%, 10%, or 5% of the volume of the protein preparation. When the antibody is produced by chemical synthesis, it is preferably substantially free of chemical precursors or other chemicals, *i.e.*, it is separated from chemical precursors or other chemicals which are involved in the synthesis of the protein. Accordingly such preparations of the antibody have less than about 30%, 20%, 10%, 5% (by dry weight) of chemical precursors or compounds other than the antibody of interest. In a preferred embodiment, antibodies of the invention are isolated or purified.

[0063] An "isolated" nucleic acid molecule is one which is separated from other nucleic acid molecules which are present in the natural source of the nucleic acid molecule. Moreover, an "isolated" nucleic acid molecule, such as a cDNA molecule, can be substantially free of other cellular material, or culture medium when produced by recombinant techniques, or substantially free of chemical precursors or other chemicals when chemically synthesized. In a specific embodiment, a nucleic acid molecule(s) encoding an antibody of the invention is isolated or purified.

[0064] The phrase "low to undetectable levels of aggregation" as used herein refers to samples containing no more than 5%, no more than 4%, no more than 3%, no more than 2%, no more than 1% and most preferably no more than 0.5% aggregation by weight of protein as measured by high performance size exclusion chromatography (HPSEC) or by a multi-sizer.

[0065] The term "low to undetectable levels of fragmentation" as used herein refers to samples containing equal to or more than 95%, 98%, 99%, 99.5% or 99.9% of the total protein, for example, as detrmined by AUC or LC-MS.

[0066] The term "lower respiratory" tract refers to the major passages and structures of the lower respiratory tract including the windpipe (trachea) and the lungs, including the bronchi, bronchioles, and alveoli of the lungs.

[0067] As used herein, the term "low tolerance" refers to a state in which the patient suffers from side effects from a therapy so that the patient does not benefit from and/or will not continue therapy because of the adverse effects and/or the harm from side effects outweighs the benefit of the therapy.

[0068] As used herein, the terms "manage", "managing" and "management" refer to the beneficial effects that a subject derives from a therapy (e.g., a prophylactic or therapeutic agent), which does not result in a cure of the infection. In certain embodiments, a subject is administered one or more therapies (e.g., prophylactic or therapeutic agents) to "manage" a infection, one or more symptoms thereof, or a respiratory condition associated with, potentiated by, or potentiating a RSV infection, so as to prevent the progression or worsening of the infection.

[0069] The terms "non-responsive" and "refractory" as used herein describe patients treated with a currently available therapy (such as but not limited to, a prophylactic or therapeutic agent) for a RSV infection, one or more symptoms thereof, or a respiratory condition associated with, potentiated by, or potentiating a RSV infection, which is not clinically adequate to relieve one or more symptoms associated with the infection. Typically, such patients suffer from severe, persistently active infection and require additional therapy to ameliorate the symptoms associated with their infection or respiratory condition.

[0070] As used herein, the terms "nucleic acids" and "nucleotide sequences" include DNA molecules (*e.g.*, cDNA or genomic DNA), RNA molecules (*e.g.*, mRNA), combinations of DNA and RNA molecules or hybrid DNA/RNA molecules, and analogues of DNA or RNA molecules. Such analogues can be generated using, for example, nucleotide analogues, which include, but are not limited to, inosine or tritylated bases. Such analogues can also comprise DNA or RNA molecules comprising modified backbones that lend beneficial attributes to the molecules such as, for example, nuclease resistance or an increased ability to cross cellular membranes. The nucleic acids or nucleotide sequences can be single-stranded, double-stranded, may contain both single-stranded and double-stranded portions, and may contain triple-stranded portions, but preferably is double-stranded DNA.

[0071] The term "pharmaceutically acceptable" as used herein means being approved by a regulatory agency of the Federal or a state government, or listed in the U.S. Pharmacopia, European Pharmacopia or other generally recognized pharmacopia for use in animals, and more particularly in humans.

[0072] The term "polyol" as used herein refers to a sugar that contains many -OH groups compared to a normal saccharide.

As used herein, the terms "prevent," "preventing," and "prevention" refer to the prevention or inhibition of the development or onset of a disease or disorder, such as an upper and/or lower respiratory tract RSV infection, otitis media or a respiratory condition related thereto in a subject, the prevention or inhibition of the progression of an upper respiratory tract RSV infection to a lower respiratory tract RSV infection, otitis media or a respiratory condition related thereto resulting from the administration of a therapy (e.g., a prophylactic or therapeutic agent), the prevention of a symptom of an upper and/or lower tract RSV infection, otitis media or a respiratory condition related thereto, or the administration of a combination of therapies (e.g., a combination of prophylactic or therapeutic agents).

As used herein, the term "prophylactic agent" refers to any agent that can prevent the recurrence, spread or onset of a disease or disorder, such as an upper and/or lower respiratory tract RSV infection, otitis media, or a symptom or respiratory condition relating thereto (including, but not limited to, asthma, wheezing, RAD, or a combination thereof), and/or prevent the progression of an upper respiratory tract RSV infection to a lower respiratory tract RSV infection or otitis media. In certain embodiments, the term "prophylactic agent" refers to an antibody of the invention. In certain other embodiments, the term "prophylactic agent" refers to an agent other than an antibody of the invention. Preferably, a prophylactic agent is an agent which is known to be useful to or has been or is currently being used to prevent or impede the onset, development, progression and/or severity of a RSV infection (preferably an upper and/or lower respiratory tract RSV infection) otitis media, and/or a symptom or respiratory condition related thereto.

[0075] In certain embodiments of the invention, a "prophylactically effective serum titer" is the serum titer in a subject, preferably a human, that reduces the incidence of an upper and/or lower respiratory tract RSV infection, otitis media and/or a symptom or respiratory condition related thereto in said subject. In some embodiments, the prophylactically effective serum titer prevents the progression of an upper respiratory tract RSV infection to a lower respiratory tract RSV infection or otitis media. Preferably, the prophylactically effective serum titer reduces the incidence of RSV infections in humans with the greatest probability of complications resulting from RSV infection (e.g., a human with cystic fibrosis, bronchopulmonary dysplasia, congenital heart disease, congenital immunodeficiency or acquired immunodeficiency, a human who has had a bone marrow transplant, a human infant, or an

elderly human). In certain other embodiments of the invention, a "prophylactically effective serum titer" is the serum titer in a cotton rat that results in a RSV titer 5 days after challenge with 10⁵ pfu that is 99% lower than the RSV titer 5 days after challenge with 10⁵ pfu of RSV in a cotton rat not administered an antibody that immunospecifically binds to a RSV antigen.

[0076] As used herein, the term "refractory" refers to an upper and/or lower respiratory tract RSV infection, otitis media or a respiratory condition related thereto that is not responsive to one or more therapies (e.g., currently available therapies). In a certain embodiment, an upper and/or lower respiratory tract RSV infection, otitis media or a respiratory condition related thereto is refractory to a therapy means that at least some significant portion of the symptoms associated with said upper and/or lower respiratory tract RSV infection, otitis media or a respiratory condition related thereto are not eliminated or lessened by that therapy. The determination of whether an upper and/or lower respiratory tract RSV infection, otitis media or a respiratory condition related thereto is refractory can be made either in vivo or in vitro by any method known in the art for assaying the effectiveness of therapy for the infection, otitis media or the respiratory condition related thereto.

[0077] The term "RSV antigen" refers to a RSV polypeptide to which an antibody immunospecifically binds. A RSV antigen also refers to an analog or derivative of a RSV polypeptide or fragment thereof to which an antibody immunospecifically binds.

[0078] The term "serum titer" as used herein refers to an average serum titer in a population of least 10, preferably at least 20, and most preferably at least 40 subjects.

[0079] The term "saccharide" as used herein refers to a class of molecules that are derivatives of polyhydric alcohols. Saccharides are commonly referred to as carbohydrates and may contain different amounts of sugar (saccharide) units, *e.g.*, monosaccharides, disaccharides and polysaccharides.

[0080] As used herein, the term "side effects" encompasses unwanted and adverse effects of a therapy (e.g., a prophylactic or therapeutic agent). Adverse effects are always unwanted, but unwanted effects are not necessarily adverse. An adverse effect from a therapy (e.g., a prophylactic or therapeutic agent) might be harmful or uncomfortable or risky. Examples of side effects include, but are not limited to, nausea, vomiting, anorexia, abdominal cramping, fever, pain, loss of body weight, dehydration, alopecia, dyspnea, insomnia, dizziness, mucositis, nerve and muscle effects, fatigue, dry mouth, and loss of appetite, rashes or swellings at the site of administration, flu-like symptoms such as fever, chills and fatigue, digestive tract

problems and allergic reactions. Additional undesired effects experienced by patients are numerous and known in the art. Many are described in the Physician's Desk Reference (58th ed., 2004).

[0081]The terms "stability" and "stable" as used herein in the context of a formulation comprising an antibody or antigen-binding fragment refer to the resistance of the antibody or antibody fragment in the formulation to thermal and chemical unfolding, aggregation, degradation or fragmentation under given manufacture, preparation, transportation and storage conditions. The "stable" formulations of the invention retain biological activity equal to or more than 80%, 85%, 90%, 95%, 98%, 99%, 99.5%, or 99.9% under given manufacture, preparation, transportation and storage conditions. The stability of the antibody or antibody fragment can be assessed by degrees of aggregation, degradation or fragmentation or levels of particular fragments (e.g., Fragment Type I or Fragment Type II) or types or sizes of aggregates by methods known to those skilled in the art, including but not limited to reduced AUC, SEC, LC-MS, particle multisizer Capillary Gel Electrophoresis (rCGE), Sodium Dodecyl Sulfate Polyacrylamide Gel Electrophoresis (SDS-PAGE) and HPSEC, compared to a reference, for example, a commercially available lyophilized palivizumab reconstituted to 100 mg/ml in 50 mM histidine/3.2 mM glycine buffer with 6% mannitol at pH 6.0. The reference regularly gives a single peak (>97% area) by HPSEC. The overall stability of a formulation comprising an antibody or fragment thereof that immunospecifically binds to a RSV antigen can be assessed by various immunological assays including, for example, ELISA and radioimmunoassay, using the specific epitope of RSV.

As used herein, the terms "subject" and "patient" are used interchangeably. As used herein, a subject is preferably a mammal such as a non-primate (e.g., cows, pigs, horses, cats, dogs, rats, etc.) and a primate (e.g., monkey and human), most preferably a human. In one embodiment, the subject is a mammal, preferably a human, with an upper and/or lower respiratory tract RSV infection or otitis media. In another embodiment, the subject is a mammal, preferably a human, at risk of developing an upper and/or lower respiratory tract RSV infection or otitis media (e.g., an immunocompromised or immunosuppressed mammal, or a genetically predisposed mammal). In one embodiment, the subject is a human with a respiratory condition (including, but not limited to asthma, wheezing or RAD) that stems from, is caused by or associated with a RSV infection.

[0083] As used herein, the term "palivizumab standard reference" or analogous terms refer to commercially available lyophilized palivizumab, as described in the Physicians' Desk

Reference, 56th edition, 2002. Reconstituted palivizumab may contain, *e.g.*, the following excipients: 47 mM histidine, 3.0 mM glycine and 5.6% manitol and the active ingredient, the antibody, at a concentration of 100 milligrams per ml solution.

[0084] As used herein, the terms "subject" and "patient" are used interchangeably. As used herein, the terms "subject" and "subjects" refer to an animal, preferably a mammal including a non-primate (e.g., a cow, pig, horse, cat, dog, rat, and mouse) and a non-primate (e.g., a monkey such as a cynomolgous monkey and a human), and more preferably a human.

[0085] The term "substantially free of surfactant" as used herein refers to a formulation of an antibody or fragment thereof that immunospecifically binds to a RSV antigen, said formulation containing less than 0.0005%, less than 0.0003%, or less than 0.0001% of surfactants and/or less than 0.0005%, less than 0.0003%, or less than 0.0001% of surfactants.

[0086] The term "substantially free of salt" as used herein refers to a formulation of an antibody or fragment thereof that immunospecifically binds to a RSV antigen, said formulation containing less than 0.0005%, less than 0.0003%, or less than 0.0001% of inorganic salts.

[0087] The term "surfactant" as used herein refers to organic substances having amphipathic structures; namely, they are composed of groups of opposing solubility tendencies, typically an oil-soluble hydrocarbon chain and a water-soluble ionic group. Surfactants can be classified, depending on the charge of the surface-active moiety, into anionic, cationic, and nonionic surfactants. Surfactants are often used as wetting, emulsifying, solubilizing, and dispersing agents for various pharmaceutical compositions and preparations of biological materials.

[0088] The term "synergistic" as used herein refers to a combination of therapies (e.g., use of prophylactic or therapeutic agents) which is more effective than the additive effects of any two or more single therapy. For example, a synergistic effect of a combination of prophylactic or therapeutic agents permits the use of lower dosages of one or more of the agents and/or less frequent administration of said agents to a subject with a RSV infection. The ability to utilize lower dosages of prophylactic or therapeutic therapies and/or to administer said therapies less frequently reduces the toxicity associated with the administration of said therapies to a subject without reducing the efficacy of said therapies in the prevention, management or treatment of a RSV infection. In addition, a synergistic effect can result in improved efficacy of therapies in the prevention or treatment of a RSV infection. Finally, synergistic effect of a

combination of therapies (e.g., prophylactic or therapeutic agents) may avoid or reduce adverse or unwanted side effects associated with the use of any single therapy.

[0089] As used herein, the term "therapeutic agent" refers to any agent that can be used in the treatment, management, prevention or amelioration of a disease or disorder, for example, an upper and/or lower respiratory tract RSV infection, otitis media or a respiratory condition related thereto. In certain embodiments, the term "therapeutic agent" refers to an antibody of the invention. In certain other embodiments, the term "therapeutic agent" refers to an agent other than an antibody of the invention. Preferably, a therapeutic agent is an agent which is known to be useful for, or has been or is currently being used for the prevention, treatment, management or amelioration of a RSV infection (*i.e.*, an upper and/or lower respiratory tract RSV infection), otitis media, or one or more symptoms or respiratory conditions related thereto.

In certain embodiments of the invention, a "therapeutically effective serum titer" is the serum titer in a subject, preferably a human, that reduces the severity, the duration and/or the symptoms associated with a RSV infection in said subject. Preferably, the therapeutically effective serum titer reduces the severity, the duration and/or the number symptoms associated with upper and/or lower respiratory tract RSV infections in humans with the greatest probability of complications resulting from the infection (*e.g.*, a human with cystic fibrosis, bronchopulmonary dysplasia, congenital heart disease, congenital immunodeficiency or acquired immunodeficiency, a human who has had a bone marrow transplant, a human infant, or an elderly human). In certain other embodiments of the invention, a "therapeutically effective serum titer" is the serum titer in a cotton rat that results in a RSV titer 5 days after challenge with 10⁵ pfu of RSV in a cotton rat not administered an antibody that immunospecifically binds to a RSV antigen.

[0091] As used herein, the term "therapy" refers to any protocol, method and/or agent that can be used in the prevention, treatment or management of a disease or disorder, such as an RSV infection (*i.e.*, an upper and/or lower respiratory tract RSV infection), otitis media, or a symptom or respiratory condition relating thereto (including, but not limited to, asthma, wheezing, RAD, or a combination thereof). In certain embodiments, the terms "therapies" and "therapy" refer to a biological therapy, supportive therapy, and/or other therapies useful in the treatment, management, prevention and/or amelioration of a RSV infection (*i.e.*, an upper and/or lower respiratory tract RSV infection), otitis media, or a symptom or respiratory condition relating thereto (including, but not limited to, asthma, wheezing, RAD, or a combination thereof) known to one of skill in the art such as medical personnel.

[0092] As used herein, the terms "treat," "treatment" and "treating" refer to the reduction or amelioration of the progression, severity, and/or duration of a disease or disorder, such as an upper and/or lower respiratory tract RSV infection, otitis media, or a symptom or respiratory condition related thereto (such as asthma, wheezing, RAD, or a combination thereof) resulting from the administration of one or more therapies (including, but not limited to, the administration of one or more prophylactic or therapeutic agents). In specific embodiments, such terms refer to the reduction or inhibition of the replication of RSV, the inhibition or reduction in the spread of RSV to other tissues or subjects (e.g., the spread to the lower respiratory tract), the inhibition or reduction of infection of a cell with a RSV, or the amelioration of one or more symptoms associated with an upper and/or lower respiratory tract RSV infection or otitis media.

[0093] The term "upper and/or lower respiratory" tract refers to the major passages and structures of the upper and/or lower respiratory tract including the nose or nostrils, nasal cavity, mouth, throat (pharynx), and voice box (larynx).

4. <u>DESCRIPTION OF THE FIGURES</u>

[0094] FIG. 1 is a schematic diagram showing an outline for preparing purified antibodies that immunospecifically bind to RSV antigen.

[0095] FIG. 2 is a schematic diagram showing an outline for preparing purified antibodies that immunospecifically bind to RSV antigen.

[0096] FIG. 3A-3B show the amino acid sequences of the (A) light chain variable region and (B) heavy chain variable region of a monoclonal antibody that binds to a RSV antigen, the potency of which can be increased by methods described herein or in Applicants' copending applications Serial Nos. 60/168,426 and 60/186,252 and U.S. Patent No. 6,656,467. For reference purposes, this is the amino acid sequence of the palivizumab antibody disclosed in Johnson et al., 1997, J. Infect. Dis. 176:1215-1224 and U.S. Patent No. 5,824,307. Here, the CDR regions are underlined while non-underlined residues form the framework (FR) regions of the variable regions of the antibody. In this antibody, the CDRs are derived from a mouse antibody while the framework regions are derived from a human antibody. The constant regions (not shown) are also derived from a human antibody.

[0097] FIG. 4A-4B show the (A) light chain variable region and (B) heavy light chain variable region for an antibody sequence. CDR regions are underlined, and the non-underlined

residues form the framework of the variable regions of the antibody. This sequence differs from the sequence disclosed in Figures 1A-1B in the first 4 residues of VH CDR1 of the light chain, residue 103 of the light chain FR4 and residue 112 of the heavy chain FR4. For reference purposes, these VL and VH sequences are identical to the VL and VH domains of IX-493L1FR (see Table 2).

- [0098] FIG. 5A-5B show the nucleotide and translated amino acid sequence of the A4B4L1FR-S28R (A) VH domain and (B) VL domain. CDR sequences are underlined. Where palivizumab differs from A4B4L1FR-S28R, the palivizumab amino acid is shown below the motavizumab sequence. Residues that were introduced on the IX-493L1FR template (see also Figure 2) are indicated in bold.
- [0099] FIG. 6A-6C. Quantitation of aggregates, fragments and monomers of A4B4L1FR-S28R during storage at (♦) 2-8 °C, (□) 20-24 °C and (▲) 38-42 °C; as determined by SEC with UV detection. (A) Percent Aggregates; (B) Percent Fragments and (C) Percent Purity (monomers).
- [00100] FIG. 7. Plot of Aggregation and fragmentation rates of A4B4L1FR-S28R based on the SEC data of FIGs. 19A-19C; (♦) rate of aggregation, (■) rate of fragmentation.
- [00101] FIG. 8. SEC profile of A4B4L1FR-S28R formulated in 25 mM histidine-HCl, pH 6.0 after storage at 38-42 °C for one month.
- [00102] FIG. 9. Comparison of AUC and SEC analysis of A4B4L1FR-S28R at initial, 9-month and 14-month time points. All samples were formulated in 25 mM histidine-HCl, pH 6.0 and, for the 9 and 14 moth points, stored at 38-42 °C. (A) AUC; (B) SEC.
- [00103] FIG. 10. Comparison of antibody sample concentration dependence of signal/noise ratio for AUC analysis.
- [00104] FIG. 11. AUC analysis of A4B4L1FR-S28R formulated in 25 mM histidine-HCl, pH 6.0 and stored at 38-42 °C over the course of 5 days.
- [00105] FIG. 12. LC-MS analysis of deglycosylated, reduced and alkylated antibody type I fragment. Sample collected from SEC of A4B4L1FR-S28R formulated in 25 mM histidine-HCl, pH 6.0 and stored at 38-42 °C for 1 month.

[00106] FIG. 13. LC-MS analysis of deglycosylated, reduced and alkylated antibody type II fragment. Sample collected from SEC of A4B4L1FR-S28R formulated in 25 mM histidine-HCl, pH 6.0 and stored at 38-42 °C for 1 month.

- [00107] FIG. 14A-14B is a diagram showing the characteristic fragmentation pattern of A4B4L1FR-S28R, forming antibody type I and antibody type II fragments. (A) Cleavage sites within the hinge region of the antibody heavy chain. Bold arrows indicate preferred or predominant cleavage sites. (B) Schematic showing characteristics of antibody type I and antibody type II fragments. an outline for preparing purified antibodies that immunospecifically bind to RSV antigen. The antibody type I fragment comprises a full length antibody light chain, a full length antibody heavy chain and a C-terminal portion of an antibody heavy chain that, in human IgG₁ immunoglobulins, has an N-terminus at cysteine 223, aspartic acid 224, lysine 225, threonine 226, histidine 227, threonine 228 or cysteine 229. The antibody type II fragment comprises an antibody light chain and an N-terminal portion of an antibody heavy chain that, in human IgG₁ immunoglobulins, has a C-terminus at serine 222, cysteine 223, aspartic acid 224, lysine 225, threonine 226, histidine 227 or threonine 228.
- [00108] FIG. 15. Chromatograms of Lys-C digested aggregates, monomers and fragments collected from SEC of A4B4L1FR-S28R formulated in 25 mM histidine-HCl, pH 6.0 and stored at 38-42 °C for 1 month. The arrows point to the low level disulfide bond scrambling peaks.
- **[00109] FIG. 16.** Chromatograms of Lys-C digested aggregates with and without reduction. Samples were collected from SEC of A4B4L1FR-S28R formulated in 25 mM histidine-HCl, pH 6.0 and stored at 38-42 °C for 1 month. The arrows point to the low level disulfide bond scrambling peaks.
- **[00110] FIG. 17** summarizes the results of a RSV microneutralization assay using the anti-RSV antibodies A4B4L1FR-S28R and palivizumab, comparing the ability of both antibodies to inhibit the *in vitro* replication of RSV (Long) in the assay.
- [00111] FIG. 18 summarizes the results of a RSV microneutralization assay demonstrating the ability of A4B4L1FR-S28R to inhibit the *in vitro* replication of RSV (Long) in the microneutralization assay.
- [00112] FIG. 19 DSC thermograms of the full length palivizumab (top panel) and an overlay of the thermograms obtained from purified Fab and Fc fragments of palivizumab (bottom panel).

Two discrete peaks are seen for the Fc domain at approximately 68°C and 83°C. A single peak is seen for the Fab fragment at approximately 87°C.

[00113] FIG. 20 plot of the Tm and pI values of palivizumab and motavizumab.

[00114] FIG.21 plot of the viscosity of a 100 mg/ml solution of palivizumab and motavizumab at a range of temperatures from about 2 to 25 °C.

[00115] FIG. 22 plot of the aggregation rates of palivizumab and motavizumab against the Fab Tm for each antibody.

5. **DETAILED DESCRIPTION OF THE INVENTION**

5.1 Methods of Preparing Antibody Formulations

The present invention provides methods for preparing formulations of antibodies, [00116] or derivatives, analogues, or fragments thereof that immunospecifically bind to a an antigen of interest. Such antibodies may be purified according to any method known in the art for purification of antibodies. FIGS. 1 and 2 are schematic diagrams showing alternate outlines for preparing purified antibodies. In one embodiment, the methods for preparing liquid formulations of the present invention comprise: concentrating a fraction containing the purified antibody or a fragment to a final antibody or fragment concentration of from about 15 mg/ml, about 20 mg/ml, about 30 mg/ml, about 40 mg/ml, about 50 mg/ml, about 60 mg/ml, about 70 mg/ml, about 80 mg/ml, about 90 mg/ml, about 100 mg/ml, about 110 mg/ml, about 125 mg/ml, about 150 mg/ml, about 200 mg/ml, about 250 mg/ml, or about 300 mg/ml using a semipermeable membrane with an appropriate molecular weight (MW) cutoff (e.g., 30 kD cutoff for whole antibody molecules and F(ab')2 fragments; and 10 kD cutoff for antibody fragments, such as Fab fragments) and diffiltrating the concentrated antibody fraction into the formulation buffer using the same membrane. Antibodies are preferably expressed in myeloma cells, more preferably murine myeloma cells, most preferably NSO cells.

[00117] In the embodiment outlined by FIG. 1, conditioned medium containing antibody or a fragment thereof that immunospecifically binds to an antigen of interest is subjected to CUNO filtration and the filtered antibody is subjected to HS50 cation exchange chromatography. The fraction from the HS50 cation exchange chromatography is then subjected to rProtein A affinity chromatography followed by low pH treatment. Following low pH treatment, the antibody fraction is subject to super Q 650 anion exchange chromatography and then nanofiltration. The fraction of the antibody obtained after nanofiltration is then subjected to diafiltration to concentrate the antibody fraction into the formulation buffer using the same membrane.

[00118] Using the embodiment of FIG. 2, conditioned medium containing antibody or a fragment thereof that immunospecifically binds to an antigen of interest is subjected to CUNO filtration and the filtered antibody is subjected to Fractogel® S cation exchange chromatography. The fraction from the cation exchange chromatography is then subjected to super Q anion chromatography. followed by nanofiltration with a Planova® 20 N nanofilter. The antibody fraction recovered after nanofiltration is then subjected to low pH treatment followed by hydroxyapatite (HA) chromatography. The fraction of the antibody obtained after HA chromatography is then subjected to diafiltration to concentrate the antibody fraction into the formulation buffer using the same membrane.

[00119] The formulation buffer of the present invention preferably comprises histidine at a concentration ranging from about 1 mM to about 100 mM, about 10 mM to about 50 mM, about 20 mM to about 30 mM, or about 23 mM to about 27 mM. Preferably, the formulation buffer of the present invention comprises histidine at a concentration of about 25 mM. The formulations may further comprise glycine at a concentration of less than 100 mM, less than 50 mM, less than 3.0 mM, less than 2.0 mM, or less than 1.8 mM. Preferably, the formulations comprise glycine at a concentration of 1.6 mM. The amount of glycine in the formulation should not cause a significant buffering in order to avoid antibody precipitation at its isoelectric point. The pH of the formulation may range from about 5.0 to about 7.0, preferably about 5.5 to about 6.5, more preferably about 5.8 to about 6.2, and most preferably about 6.0. To obtain an appropriate pH for a particular antibody, it is preferable that histidine (and glycine, if added) is first dissolved in water to obtain a buffer solution with higher pH than the desired pH and then the pH is brought down to the desired level by adding HCl. This way, the formation of inorganic salts (e.g., formation of NaCl when, for example, histidine hydrochloride is used as histidine and pH is raised to a desired level by adding NaOH) can be avoided.

[00120] The formulations of the present invention can be prepared as unit dosage forms by preparing a vial containing an aliquot of the liquid formulation for a one-time use. For example, a unit dosage per vial may contain 1 ml, 2 ml, 3 ml, 4 ml, 5 ml, 6 ml, 7 ml, 8 ml, 9 ml, 10 ml, 15 ml, or 20 ml of different concentrations of an antibody or a fragment thereof that immunospecifically binds to the antigen of interest ranging from about 15 mg/ml to about 300 mg/ml. If necessary, these preparations can be adjusted to a desired concentration by adding a sterile diluent to each vial.

[00121] The formulations of the present invention may be sterilized by various sterilization methods, including sterile filtration, radiation, etc. In a most preferred embodiment,

the difiltrated antibody formulation is filter-sterilized with a presterilized 0.22-micron filter. In specific embodiments, sterilized liquid formulations of the present invention may be administered to a subject to prevent, treat, manage or ameliorate a RSV infection, one or more symptoms thereof, or a respiratory condition associated with, potentiated by, potentiating a RSV infection.

[00122] The formulations of the invention comprise labeled antibodies, derivatives and analogues thereof, that immunospecifically bind to an antigen of interest and can be used for diagnostic purposes to detect, diagnose, or monitor a disorder associated with and/or characterized by the presence or said antigen. In a specific embodiment, the formulations of the invention comprise labeled antibodies, derivatives and analogues thereof, that immunospecifically bind to a RSV antigen and can be used for diagnostic purposes to detect, diagnose, or monitor a RSV infection.

[00123] The invention encompasses both liquid and lyophilized forms of the formulations. Methods to produce lyophilized forms of liquid formulations are well-characterized in the art. In one embodiment, the ingredients of formulation of the invention are supplied either separately or mixed together in unit dosage form, for example, as a dry lyophilized powder or water free concentrate in a hermetically sealed container such as an ampoule or sachette indicating the quantity of active agent. Where the composition is to be administered by infusion, it can be dispensed with an infusion bottle containing sterile pharmaceutical grade water or saline. Where the composition is administered by injection, an ampoule of sterile water for injection or saline can be provided so that the ingredients may be mixed prior to administration.

[00124] The compositions of the invention can be formulated as neutral or salt forms. Pharmaceutically acceptable salts include those formed with anions such as those derived from hydrochloric, phosphoric, acetic, oxalic, tartaric acids, etc., and those formed with cations such as those derived from sodium, potassium, ammonium, calcium, ferric hydroxides, isopropylamine, triethylamine, 2-ethylamino ethanol, histidine, procaine, etc.

[00125] The formulation of the invention can be further processed into an oral or non-oral dosage form, for immediate or extended release. The formulation can additionally comprise inactive ingredients ordinarily used in pharmaceutical preparation such as diluents, fillers, disintegrants, sweeteners, lubricants and flavors. The formulation may also be processed for intravenous administration, either by bolus injection or sustained drip, or for release from an

implanted capsule. A typical formulation for intravenous administration utilizes physiological saline as a diluent.

5.2 Formulations of Antibodies

[00126] The invention provides formulations comprising antibodies of the invention for use in diagnosing, detecting, or monitoring a disorder, in preventing, treating, managing, or ameliorating of a disorder or one or more symptoms thereof, and/or in research. In a specific embodiment, the formulation of the invention comprises one or more antibodies. In another embodiment, the formulation of the invention comprises one or more antibodies and one or more prophylactic or therapeutic agents other than antibodies. Preferably, the prophylactic or therapeutic agents known to be useful for or having been or currently being used in the prevention, treatment, management, or amelioration of a disorder or one or more symptoms thereof. In accordance with these embodiments, the composition may further comprise of a carrier, diluent or excipient.

[00127] The formulations of the present invention provide antibody formulations which are substantially free of surfactant, inorganic salts, and/or other excipients and yet exhibit high stability during long periods of storage. In a specific embodiment, such antibody formulations are homogeneous. The formulations of the present invention comprise histidine at concentrations between 1 and 100 mM and an antibody which immunospecifically binds to a antigen of interest at concentrations of about 15 mg/ml to about 300 mg/ml. In one embodiment, the formulations of the invention do not comprise other ingredients except for water or suitable solvents. In a specific embodiment, the antibody immunospecifically binds to an RSV antigen and in preferred embodiments is not palivizumab or a fragment thereof.

[00128] In one embodiment, the antibody of the formulation of the invention is an antibody or antibody fragment conjugated to another moiety, including, but not limited to, a heterologous polypeptide, another antibody or another fragment, a marker sequence, a diagnostic agent, a therapeutic agent, a radioactive metal ion, a polymer, albumin, and a solid support. In another embodiment, formulations of the invention comprise two or more antibodies, or fragments thereof that immunospecifically binds to an antigen of interest. In a specific embodiment, formulations of the invention comprise two or more antibodies, or fragments thereof, that immunospecifically binds to a RSV antigen, wherein at least one of the antibodies or antibody fragments is not palivizumab or a fragment thereof.

The concentration of an antibody or a fragment thereof which is included in the formulations of the invention is at least 15 mg/ml, at least 20 mg/ml, at least 25 mg/ml, at least 30 mg/ml, at least 35 mg/ml, at least 40 mg/ml, at least 45 mg/ml, at least 50 mg/ml, at least 50 mg/ml, at least 50 mg/ml, at least 60 mg/ml, at least 65 mg/ml, at least 70 mg/ml, at least 75 mg/ml, at least 80 mg/ml, at least 85 mg/ml, at least 90 mg/ml, at least 95 mg/ml, at least 100 mg/ml, at least 105 mg/ml, at least 110 mg/ml, at least 115 mg/ml, at least 120 mg/ml, at least 125 mg/ml, at least 130 mg/ml, at least 300 mg/ml, at least 300 mg/ml, at least 250 mg/ml, or at least 300 mg/ml.

[00130] The concentration of histidine which is included in the formulations of the invention ranges from about 1 mM to about 100 mM, about 10 mM to about 50 mM, about 20 mM to about 30 mM, or about 23 mM to about 27 mM, and is most preferably about 25 mM. Histidine can be in the form of L-histidine, D-histidine, or a mixture thereof, but L-histidine is the most preferable. Histidine can also be in the form of hydrates. Histidine may be used in a form of pharmaceutically acceptable salt, such as hydrochloride (*e.g.*, monohydrochloride and dihydrochloride), hydrobromide, sulfate, acetate, etc. The purity of histidine should be at least 98%, preferably at least 99%, and most preferably at least 99.5%.

[00131] The pH of the formulation should not be equal to the isoelectric point of the particular antibody to be used in the formulation and may range from about 5.0 to about 7, preferably about 5.5 to about 6.5, more preferably about 5.8 to about 6.2, and most preferably about 6.0.

In addition to histidine and an antibody or a fragment thereof, the formulations of the present invention may further comprise glycine at a concentration of less than 100 mM, less than 50 mM, less than 3.0 mM, less than 2.0 mM, or less than 1.8 mM, and most preferably 1.6 mM. The amount of glycine in the formulation should not cause a significant buffering effect so that antibody precipitation at its isoelectric point can be avoided. Glycine may be also used in a form of pharmaceutically acceptable salt, such as hydrochloride, hydrobromide, sulfate, acetate, etc. The purity of glycine should be at least 98%, preferably at least 99%, and most preferably 99.5%. In a specific embodiment, glycine is included in the formulations of the present invention.

[00133] Optionally, the formulations of the present invention may further comprise other excipients, such as saccharides (e.g., sucrose, mannose, trehalose, etc.) and polyols (e.g., mannitol, sorbitol, etc.). In one embodiment, the other excipient is a saccharide. In a specific

embodiment, the saccharide is sucrose, which is at a concentration ranging from between about 1% to about 20%, preferably about 5% to about 15%, and more preferably about 8% to 10%. In another embodiment, the other excipient is a polyol. Preferably, however, the liquid formulations of the present invention do not contain mannitol. In a specific embodiment, the polyol is polysorbate (*e.g.*, Tween 20), which is at a concentration ranging from between about 0.001% to about 1%, preferably, about 0.01 to about 0.1.

[00134] The formulations of the present invention exhibit stability at the temperature ranges of 38°C-42°C for at least 60 days and, in some embodiments, at least 120 days, of 20°C-24°C for at least 1 year, of 2°C-8°C (in particular, at 4°C) for at least 3 years, at least 4 years, or at least 5 years and at -20°C for at least 3 years, at least 4 years, or at least 5 years, as assessed by AUC, LC-MS, size exclusion chromatography (SEC) or high performance size exclusion chromatography (HPSEC) or particle multisizer. Namely, the formulations of the present invention have low to undetectable levels of aggregation and/or fragmentation, as defined herein, after the storage for the defined periods as set forth above. Preferably, no more than 5%, no more than 4%, no more than 3%, no more than 2%, no more than 1%, and most preferably no more than 0.5% (but in certain embodiments, at least 0.1%) of the antibody or antibody fragment forms an aggregate or fragment (particularly of fragment I or fragment II) as measured by AUC, LC-MS, SEC or HPSEC, after the storage for the defined periods as set forth above. Furthermore, formulations of the present invention exhibit almost no loss in biological activities of the antibody or antibody fragment during the prolonged storage under the condition described above, as assessed by various immunological assays including, but not limited to, enzyme-linked immunosorbent assay (ELISA) and radioimmunoassay to measure the ability of the antibody or antibody fragment to immunospecifically bind to an antigen of interest, and by a C3a/C4a assay to measure the complement activating ability of the antibody. The formulations of the present invention retain after the storage for the above-defined periods more than 80%, more than 85%, more than 90%, more than 95%, more than 98%, more than 99%, or more than 99.5% of the initial biological activities of the formulation prior to the storage.

[00135] The formulations of the present invention can be prepared as unit dosage forms. For example, a unit dosage per vial may contain 1 ml, 2 ml, 3 ml, 4 ml, 5 ml, 6 ml, 7 ml, 8 ml, 9 ml, 10 ml, 15 ml, or 20 ml of different concentrations of an antibody or a fragment thereof that immunospecifically binds to a RSV antigen ranging from about 15 mg/ml to about 300 mg/ml. If necessary, these preparations can be adjusted to a desired concentration by adding a sterile diluent to each vial.

[00136] The invention encompasses stable liquid formulations comprising a single antibody or fragment thereof that immunospecifically binds to an antigen of interest. In a specific embodiment, the invention encompasses stable liquid formulations comprising a single antibody or fragment thereof that immunospecifically binds to a RSV antigen, with the proviso that said antibody is not palivizumab. The invention also encompasses stable liquid formulations comprising two or more antibodies or fragments thereof that immunospecifically bind to a RSV antigen. In one embodiment, a stable liquid formulation of the invention comprises two or more antibodies or fragments thereof that immunospecifically bind to a RSV antigen, wherein one of the antibodies or antibody fragments is not palivizumab or a fragment thereof.

5.3 Antibodies Useful in the Formulations of the Invention

[00137] The antibodies useful in the present invention include, but are not limited to, monoclonal antibodies, synthetic antibodies, multispecific antibodies (including bi-specific antibodies), human antibodies, humanized antibodies, chimeric antibodies, single-chain Fvs (scFv) (including bi-specific scFvs), single chain antibodies, Fab fragments, F(ab') fragments, disulfide-linked Fvs (sdFv), and epitope-binding fragments of any of the above. In particular, antibodies of the present invention include immunoglobulin molecules and immunologically active portions of immunoglobulin molecules, *i.e.*, molecules that contain an antigen binding site that immunospecifically binds to an antigen. The immunoglobulin molecules of the invention can be of any type (*e.g.*, IgG, IgE, IgM, IgD, IgA and IgY), class (*e.g.*, IgG₁, IgG₂, IgG₃, IgG₄, IgA₁ and IgA₂) or subclass of immunoglobulin molecule. Preferably, the antibodies of the invention are IgG, more preferably, IgG₁.

[00138] The antibodies useful in the present invention may be from any animal origin including birds and mammals (e.g., human, murine, donkey, sheep, rabbit, goat, guinea pig, camel, horse, or chicken). Preferably, the antibodies are human or humanized monoclonal antibodies. As used herein, "human" antibodies include antibodies having the amino acid sequence of a human immunoglobulin and include antibodies isolated from human immunoglobulin libraries or from mice or other animal that express antibodies from human genes.

[00139] The antibodies useful in the present invention may be monospecific, bispecific, trispecific or of greater multispecificity. Multispecific antibodies may immunospecifically bind to different epitopes of a polypeptide or may immunospecifically bind to both a polypeptide as well a heterologous epitope, such as a heterologous polypeptide or solid support material. See,

e.g., International Publication Nos. WO 93/17715, WO 92/08802, WO 91/00360, and WO 92/05793; Tutt, et al., 1991, *J. Immunol*. 147:60-69; U.S. Patent Nos. 4,474,893, 4,714,681, 4,925,648, 5,573,920, and 5,601,819; and Kostelny et al., 1992, *J. Immunol*. 148:1547-1553.

The antibodies useful in the present invention include derivatives of the [00140] antibodies. Standard techniques known to those of skill in the art can be used to introduce mutations in the nucleotide sequence encoding an antibody to be used with the methods of the invention, including, for example, site-directed mutagenesis and PCR-mediated mutagenesis which result in amino acid substitutions. Preferably, the derivatives include less than 25 amino acid substitutions, less than 20 amino acid substitutions, less than 15 amino acid substitutions, less than 10 amino acid substitutions, less than 5 amino acid substitutions, less than 4 amino acid substitutions, less than 3 amino acid substitutions, or less than 2 amino acid substitutions relative to the original molecule. In a preferred embodiment, the derivatives have conservative amino acid substitutions are made at one or more predicted non-essential amino acid residues. A "conservative amino acid substitution" is one in which the amino acid residue is replaced with an amino acid residue having a side chain with a similar charge. Families of amino acid residues having side chains with similar charges have been defined in the art. These families include amino acids with basic side chains (e.g., lysine, arginine, histidine), acidic side chains (e.g., aspartic acid, glutamic acid), uncharged polar side chains (e.g., glycine, asparagine, glutamine, serine, threonine, tyrosine, cysteine), nonpolar side chains (e.g., alanine, valine, leucine, isoleucine, proline, phenylalanine, methionine, tryptophan), beta-branched side chains (e.g., threonine, valine, isoleucine) and aromatic side chains (e.g., tyrosine, phenylalanine, tryptophan, histidine). Alternatively, mutations can be introduced randomly along all or part of the coding sequence, such as by saturation mutagenesis, and the resultant mutants can be screened for biological activity to identify mutants that retain activity. Following mutagenesis, the encoded protein can be expressed and the activity of the protein can be determined.

[00141] The antibodies useful in the present invention include derivatives that are modified, *i.e*, by the covalent attachment of any type of molecule to the antibody such that covalent attachment. For example, but not by way of limitation, the antibody derivatives include antibodies that have been modified, *e.g.*, by glycosylation, acetylation, pegylation, phosphorylation, amidation, derivatization by known protecting/blocking groups, proteolytic cleavage, linkage to a cellular ligand or other protein, etc. Any of numerous chemical modifications may be carried out by known techniques, including, but not limited to specific

chemical cleavage, acetylation, formylation, synthesis in the presence of tunicamycin, etc. Additionally, the derivative may contain one or more non-classical amino acids.

[00142] Antibodies useful in the present invention or fragments thereof can also comprise a framework region known to those of skill in the art. In certain embodiments, one or more framework regions, preferably, all of the framework regions, of an antibody to be used in the methods of the invention or fragment thereof are human. In certain other embodiments of the invention, the fragment region of an antibody of the invention or fragment thereof is humanized. In certain embodiments, the antibody to be used with the methods of the invention is a synthetic antibody, a monoclonal antibody, an intrabody, a chimeric antibody, a human antibody, a humanized chimeric antibody, a humanized antibody, a glycosylated antibody, a multispecific antibody, a human antibody, a single-chain antibody, or a bispecific antibody.

In certain embodiments of the invention, the antibodies useful in the present [00143] invention have half-lives in a mammal, preferably a human, of greater than 12 hours, greater than 1 day, greater than 3 days, greater than 6 days, greater than 10 days, greater than 15 days, greater than 20 days, greater than 25 days, greater than 30 days, greater than 35 days, greater than 40 days, greater than 45 days, greater than 2 months, greater than 3 months, greater than 4 months, or greater than 5 months. Antibodies or antigen-binding fragments thereof having increased in vivo half-lives can be generated by techniques known to those of skill in the art. For example, antibodies or antigen-binding fragments thereof with increased in vivo half-lives can be generated by modifying (e.g., substituting, deleting or adding) amino acid residues identified as involved in the interaction between the Fc domain and the FcRn receptor (see, e.g., PCT Publication No. WO 97/34631 and U.S. Patent Application No.: 10/020,354, entitled "Molecules with Extended Half-Lives, Compositions and Uses Thereof", filed December 12, 2001, by Johnson et al., which are incorporated herein by reference in their entireties). Such antibodies or antigen-binding fragments thereof can be tested for binding activity to RSV antigens as well as for in vivo efficacy using methods known to those skilled in the art, for example, by immunoassays described herein.

[00144] Further, antibodies or antigen-binding fragments thereof with increased *in vivo* half-lives can be generated by attaching to said antibodies or antibody fragments polymer molecules such as high molecular weight polyethyleneglycol (PEG). PEG can be attached to said antibodies or antibody fragments with or without a multifunctional linker either through site-specific conjugation of the PEG to the N- or C- terminus of said antibodies or antibody fragments or via epsilon-amino groups present on lysine residues. Linear or branched polymer

derivatization that results in minimal loss of biological activity will be used. The degree of conjugation will be closely monitored by SDS-PAGE and mass spectrometry to ensure proper conjugation of PEG molecules to the antibodies. Unreacted PEG can be separated from antibody-PEG conjugates by, *e.g.*, size exclusion or ion-exchange chromatography. PEG-derivatizated antibodies or antigen-binding fragments thereof can be tested for binding activity to RSV antigens as well as for *in vivo* efficacy using methods known to those skilled in the art, for example, by immunoassays described herein.

[00145] The antibodies useful in the present invention can be single-chain antibodies. The design and construction of a single-chain antibody is described in Marasco et al, 1993, Proc Natl Acad Sci 90:7889-7893, which is incorporated herein by reference in its entirety.

In certain embodiments, the antibodies useful in the present invention bind to an intracellular epitope, *i.e.*, are intrabodies. An intrabody comprises at least a portion of an antibody that is capable of immunospecifically binding an antigen and preferably does not contain sequences coding for its secretion. Such antibodies will bind its antigen intracellularly. In one embodiment, the intrabody comprises a single-chain Fv ("sFv"). sFv are antibody fragments comprising the V_H and V_L domains of antibody, wherein these domains are present in a single polypeptide chain. Generally, the Fv polypeptide further comprises a polypeptide linker between the V_H and V_L domains which enables the sFv to form the desired structure for antigen binding. For a review of sFv see Pluckthun in *The Pharmacology of Monoclonal Antibodies*, vol. 113, Rosenburg and Moore eds. Springer-Verlag, New York, pp. 269-315 (1994).

[00147] In a further embodiment, the intrabody preferably does not encode an operable secretory sequence and thus remains within the cell (see generally Marasco, WA, 1998, "Intrabodies: Basic Research and Clinical Gene Therapy Applications" Springer:New York).

5.3.1 Antibody Conjugates

[00148] The present invention also encompasses formulations comprising antibodies that are conjugated or fused to one or more moieties, including but not limited to, peptides, polypeptides, proteins, fusion proteins, nucleic acid molecules, small molecules, mimetic agents, synthetic drugs, inorganic molecules, and organic molecules.

[00149] The present invention encompasses formulations comprising antibodies that are recombinantly fused or chemically conjugated (including both covalent and non-covalent conjugations) to a heterologous protein or polypeptide (or fragment thereof, preferably to a

polypepetide of at least 10, at least 20, at least 30, at least 40, at least 50, at least 60, at least 70, at least 80, at least 90 or at least 100 amino acids) to generate fusion proteins. The fusion does not necessarily need to be direct, but may occur through linker sequences. For example, antibodies may be used to target heterologous polypeptides to particular cell types, either in vitro or in vivo, by fusing or conjugating the antibodies to antibodies specific for particular cell surface receptors. Antibodies fused or conjugated to heterologous polypeptides may also be used in in vitro immunoassays and purification methods using methods known in the art. See e.g., International publication No. WO 93/21232; European Patent No. EP 439,095; Naramura et al., 1994, Immunol. Lett. 39:91-99; U.S. Pat. No. 5,474,981; Gillies et al., 1992, PNAS 89:1428-1432; and Fell et al., 1991, J. Immunol. 146:2446-2452, which are incorporated by reference in their entireties.

[00150] The present invention further includes formulations comprising heterologous proteins, peptides or polypeptides fused or conjugated to antibody fragments. For example, the heterologous polypeptides may be fused or conjugated to a Fab fragment, Fd fragment, Fv fragment, F(ab)₂ fragment, a VH domain, a VL domain, a VH CDR, a VL CDR, or fragment thereof. Methods for fusing or conjugating polypeptides to antibody portions are well-known in the art. See, e.g., U.S. Pat. Nos. 5,336,603, 5,622,929, 5,359,046, 5,349,053, 5,447,851, and 5,112,946; European Patent Nos. EP 307,434 and EP 367,166; International publication Nos. WO 96/04388 and WO 91/06570; Ashkenazi et al., 1991, Proc. Natl. Acad. Sci. USA 88: 10535-10539; Zheng et al., 1995, J. Immunol. 154:5590-5600; and Vil et al., 1992, Proc. Natl. Acad. Sci. USA 89:11337-11341 (said references incorporated by reference in their entireties).

[00151] Additional fusion proteins may be generated through the techniques of gene-shuffling, motif-shuffling, exon-shuffling, and/or codon-shuffling (collectively referred to as "DNA shuffling"). DNA shuffling may be employed to alter the activities of antibodies of the invention or fragments thereof (e.g., antibodies or fragments thereof with higher affinities and lower dissociation rates). See, generally, U.S. Pat. Nos. 5,605,793; 5,811,238; 5,830,721; 5,834,252; and 5,837,458, and Patten et al., 1997, Curr. Opinion Biotechnol. 8:724-33; Harayama, 1998, Trends Biotechnol. 16(2):76-82; Hansson, et al., 1999, J. Mol. Biol. 287:265-76; and Lorenzo and Blasco, 1998, Biotechniques 24(2):308-313 (each of these patents and publications are hereby incorporated by reference in its entirety). Antibodies or fragments thereof, or the encoded antibodies or fragments thereof, may be altered by being subjected to random mutagenesis by error-prone PCR, random nucleotide insertion or other methods prior to recombination. One or more portions of a polynucleotide encoding an antibody or antibody

fragment may be recombined with one or more components, motifs, sections, parts, domains, fragments, etc. of one or more heterologous molecules.

[00152] Moreover, the antibodies or fragments thereof can be fused to marker sequences, such as a peptide to facilitate purification. In embodiments, the marker amino acid sequence is a hexa-histidine peptide, such as the tag provided in a pQE vector (QIAGEN, Inc., 9259 Eton Avenue, Chatsworth, Calif., 91311), among others, many of which are commercially available. As described in Gentz et al., 1989, Proc. Natl. Acad. Sci. USA 86:821-824, for instance, hexa-histidine provides for convenient purification of the fusion protein. Other peptide tags useful for purification include, but are not limited to, the hemagglutinin "HA" tag, which corresponds to an epitope derived from the influenza hemagglutinin protein (Wilson et al., 1984, Cell 37:767) and the "flag" tag.

In other embodiments, antibodies useful in the present invention or fragments, [00153] analogs or derivatives thereof can be conjugated to a diagnostic or detectable agent. Such antibodies can be useful for monitoring or prognosing the development or progression of a disorder as part of a clinical testing procedure, such as determining the efficacy of a particular therapy. Such diagnosis and detection can be accomplished by coupling the antibody to detectable substances including, but not limited to various enzymes, such as but not limited to horseradish peroxidase, alkaline phosphatase, beta-galactosidase, or acetylcholinesterase; prosthetic groups, such as but not limited to streptavidin/biotin and avidin/biotin; fluorescent materials, such as but not limited to, umbelliferone, fluorescein, fluorescein isothiocynate, rhodamine, dichlorotriazinylamine fluorescein, dansyl chloride or phycoerythrin; luminescent materials, such as but not limited to, luminol; bioluminescent materials, such as but not limited to, luciferase, luciferin, and aequorin; radioactive materials, such as but not limited to iodine (¹³¹I, ¹²⁵I, ¹²³I, ¹²¹I,), carbon (¹⁴C), sulfur (³⁵S), tritium (³H), indium (¹¹⁵In, ¹¹¹In, ¹¹¹In,), and technetium (⁹⁹Tc), thallium (²⁰¹Ti), gallium (⁶⁸Ga, ⁶⁷Ga), palladium (¹⁰³Pd), molybdenum (99Mo), xenon (133Xe), fluorine (18F), 153Sm, 177Lu, 159Gd, 149 Pm, 140La, 175Yb, 166Ho, 90Y, ⁴⁷Sc, ¹⁸⁶Re, ¹⁸⁸Re, ¹⁴²Pr, ¹⁰⁵Rh, ⁹⁷Ru, ⁶⁸Ge, ⁵⁷Co, ⁶⁵Zn, ⁸⁵Sr, ³²P, ¹⁵³Gd, ¹⁶⁹Yb, ⁵¹Cr, ⁵⁴Mn, ⁷⁵Se. ¹¹³Sn. and ¹¹⁷Tin; positron emitting metals using various positron emission tomographies, noradioactive paramagnetic metal ions, and molecules that are radiolabelled or conjugated to specific radioisotopes.

[00154] The present invention further encompasses formulations comprising antibodies that are conjugated to a therapeutic moiety. An antibody or fragment thereof may be conjugated to a therapeutic moiety such as a cytotoxin, e.g., a cytostatic or cytocidal agent, a therapeutic

agent or a radioactive metal ion, e.g., alpha-emitters. A cytotoxin or cytotoxic agent includes any agent that is detrimental to cells. Therapeutic moieties include, but are not limited to, antimetabolites (e.g., methotrexate, 6-mercaptopurine, 6-thioguanine, cytarabine, 5-fluorouracil decarbazine), alkylating agents (e.g., mechlorethamine, thioepa chlorambucil, melphalan, carmustine (BCNU) and lomustine (CCNU), cyclothosphamide, busulfan, dibromomannitol, streptozotocin, mitomycin C, and cisdichlorodiamine platinum (II) (DDP) cisplatin), anthracyclines (e.g., daunorubicin (formerly daunomycin) and doxorubicin), antibiotics (e.g., dactinomycin (formerly actinomycin), bleomycin, mithramycin, and anthramycin (AMC)). Auristatin molecules (e.g., auristatin PHE, bryostatin 1, and solastatin 10; see Woyke et al., Antimicrob. Agents Chemother. 46:3802-8 (2002), Woyke et al., Antimicrob. Agents Chemother, 45:3580-4 (2001), Mohammad et al., Anticancer Drugs 12:735-40 (2001), Wall et al., Biochem, Biophys, Res. Commun. 266:76-80 (1999), Mohammad et al., Int. J. Oncol. 15:367-72 (1999), all of which are incorporated herein by reference), hormones (e.g., glucocorticoids, progestins, androgens, and estrogens), DNA-repair enzyme inhibitors (e.g., etoposide or topotecan), kinase inhibitors (e.g., compound ST1571, imatinib mesylate (Kantaijian et al., Clin Cancer Res. 8(7):2167-76 (2002)), cytotoxic agents (e.g., paclitaxel, cytochalasin B, gramicidin D, ethidium bromide, emetine, mitomycin, etoposide, tenoposide, vincristine, vinblastine, colchicin, doxorubicin, daunorubicin, dihydroxy anthracin dione, mitoxantrone, mithramycin, actinomycin D, 1-dehydrotestosterone, procaine, tetracaine, lidocaine, propranolol, and puromycin and analogs or homologs thereof) and those compounds disclosed in U.S. Pat. Nos. 6,245,759, 6,399,633, 6,383,790, 6,335,156, 6,271,242, 6,242,196, 6,218,410, 6,218,372, 6,057,300, 6,034,053, 5,985,877, 5,958,769, 5,925,376, 5,922,844, 5,911,995, 5,872,223, 5,863,904, 5,840,745, 5,728,868, 5,648,239, 5,587,459), farnesyl transferase inhibitors (e.g., R115777, BMS-214662, and those disclosed by, for example, U.S. Pat. Nos. 6,458,935, 6,451,812, 6,440,974, 6,436,960, 6,432,959, 6,420,387, 6,414,145, 6,410,541, 6,410,539, 6,403,581, 6,399,615, 6,387,905, 6,372,747, 6,369,034, 6,362,188, 6,342,765, 6,342,487, 6,300,501, 6,268,363, 6,265,422, 6,248,756, 6,239,140, 6,232,338. 6,228,865, 6,228,856, 6,225,322, 6,218,406, 6,211,193, 6,187,786, 6,169,096, 6,159,984, 6,143,766, 6,133,303, 6,127,366, 6,124,465, 6,124,295, 6,103,723, 6,093,737, 6,090,948, 6,080,870, 6,077,853, 6,071,935, 6,066,738, 6,063,930, 6,054,466, 6,051,582, 6,051,574, and 6,040,305), topoisomerase inhibitors (e.g., camptothecin; irinotecan; SN-38; topotecan; 9aminocamptothecin; GG-211 (GI 147211); DX-895 If; IST-622; rubitecan; pyrazoloacridine; XR-5000; saintopin; UCE6; UCE1022; TAN-1518A; TAN-1518B; KT6006; KT6528; ED-110; NB-506; ED-110; NB-506; and rebeccamycin); bulgarein; DNA minor groove binders such as Hoescht dve 33342 and Hoechst dve 33258; nitidine; fagaronine; epiberberine; coralyne; beta-

lapachone; BC-4-1; bisphosphonates (e.g., alendronate, cimadronte, clodronate, tiludronate, etidronate, ibandronate, neridronate, olpandronate, risedronate, piridronate, pamidronate, zolendronate) HMG-CoA reductase inhibitors, (e.g., lovastatin, simvastatin, atorvastatin, pravastatin, fluvastatin, statin, cerivastatin, lescol, lupitor, rosuvastatin and atorvastatin) and pharmaceutically acceptable salts, solvates, clathrates, and prodrugs thereof. See, e.g., Rothenberg, M. L., Annals of Oncology 8:837-855(1997); and Moreau, P., et al., J. Med. Chem. 41:1631-1640(1998)), antisense oligonucleotides (e.g., those disclosed in the U.S. Pat. Nos. 6,277,832, 5,998,596, 5,885,834, 5,734,033, and 5,618,709), immunomodulators (e.g., antibodies and cytokines), antibodies, and adenosine deaminase inhibitors (e.g., Fludarabine phosphate and 2-Chlorodeoxyadenosine).

Further, an antibody or fragment thereof may be conjugated to a therapeutic [00155] moiety or drug moiety that modifies a given biological response. Therapeutic moieties or drug moieties are not to be construed as limited to classical chemical therapeutic agents. For example, the drug moiety may be a protein or polypeptide possessing a desired biological activity. Such proteins may include, for example, a toxin such as abrin, ricin A, pseudomonas exotoxin, cholera toxin, or diphtheria toxin; a protein such as tumor necrosis factor, .alpha.-interferon, .beta.-interferon, nerve growth factor, platelet derived growth factor, tissue plasminogen activator, an apoptotic agent, e.g., TNFα, TNFβ, AIM I (see, International publication No. WO 97/33899), AIM II (see, International Publication No. WO 97/34911), Fas Ligand (Takahashi et al., 1994, J. Immunol., 6:1567-1574), and VEGI (see, International publication No. WO 99/23105), a thrombotic agent or an anti-angiogenic agent, e.g., angiostatin, endostatin or a component of the coagulation pathway (e.g., tissue factor); or, a biological response modifier such as, for example, a lymphokine (e.g., interleukin-1 ("IL-1"), interleukin-2 ("IL-2"), interleukin-6 ("IL-6"), granulocyte macrophage colony stimulating factor ("GM-CSF"), and granulocyte colony stimulating factor ("G-CSF")), a growth factor (e.g., growth hormone ("GH")), or a coagulation agent (e.g., calcium, vitamin K, tissue factors, such as but not limited to, Hageman factor (factor XII), high-molecular-weight kininogen (HMWK), prekallikrein (PK), coagulation proteins-factors II (prothrombin), factor V, XIIa, VIII, XIIIa, XI, XIa, IX, IXa, X, phospholipid. fibrinopeptides A and B from the α and β chains of fibrinogen, fibrin monomer).

[00156] Moreover, an antibody can be conjugated to therapeutic moieties such as a radioactive metal ion, such as alph-emiters such as ²¹³Bi or macrocyclic chelators useful for conjugating radiometal ions, including but not limited to, ¹³¹In, ¹³¹LU, ¹³¹Y, ¹³¹Ho, ¹³¹Sm, to polypeptides. In certain embodiments, the macrocyclic chelator is 1,4,7,10-

tetraazacyclododecane-N,N',N",N"-tetraacetic acid (DOTA) which can be attached to the antibody via a linker molecule. Such linker molecules are commonly known in the art and described in Denardo et al., 1998, Clin Cancer Res. 4(10):2483-90; Peterson et al., 1999, Bioconjug. Chem. 10(4):553-7; and Zimmerman et al., 1999, Nucl. Med. Biol. 26(8):943-50, each incorporated by reference in their entireties.

[00157] Techniques for conjugating therapeutic moieties to antibodies are well known, see, e.g., Arnon et al., "Monoclonal Antibodies For Immunotargeting Of Drugs In Cancer Therapy", in Monoclonal Antibodies And Cancer Therapy, Reisfeld et al. (eds.), pp. 243-56 (Alan R. Liss, Inc. 1985); Hellstrom et al., "Antibodies For Drug Delivery", in Controlled Drug Delivery (2nd Ed.), Robinson et al. (eds.), pp. 623-53 (Marcel Dekker, Inc. 1987); Thorpe, "Antibody Carriers Of Cytotoxic Agents In Cancer Therapy: A Review", in Monoclonal Antibodies 84: Biological And Clinical Applications, Pinchera et al. (eds.), pp. 475-506 (1985); "Analysis, Results, And Future Prospective Of The Therapeutic Use Of Radiolabeled Antibody In Cancer Therapy", in Monoclonal Antibodies For Cancer Detection And Therapy, Baldwin et al. (eds.), pp. 303-16 (Academic Press 1985), and Thorpe et al., 1982, Immunol. Rev. 62:119-58.

[00158] Alternatively, an antibody can be conjugated to a second antibody to form an antibody heteroconjugate as described by Segal in U.S. Pat. No. 4,676,980, which is incorporated herein by reference in its entirety.

[00159] The therapeutic moiety or drug conjugated to an antibody or fragment thereof should be chosen to achieve the desired prophylactic or therapeutic effect(s) for a particular disorder in a subject. A clinician or other medical personnel should consider the following when deciding on which therapeutic moiety or drug to conjugate to an antibody or fragment thereof: the nature of the disease, the severity of the disease, and the condition of the subject.

[00160] Antibodies may also be attached to solid supports, which are particularly useful for immunoassays or purification of the target antigen. Such solid supports include, but are not limited to, glass, cellulose, polyacrylamide, nylon, polystyrene, polyvinyl chloride or polypropylene.

5.3.2 Formulations Comprising Purified Antibodies that Specifically Bind to a Particular Antigen

[00161] In further embodiments, the present invention encompasses formulations comprising isolated antibodies or compositions comprising antibodies, wherein said antibodies specifically bind to one or more particular antigens. In certain embodiments, the antibody of the

present invention specifically binds to an antigen of respiratory syncytial virus (RSV). In other embodiments, the antibody of the present invention specifically binds to an antigen of human metapneumovirus (hMPV). In some embodiments, the antibody is a humanized antibody that specifically binds to an antigen of hMPV. In certain embodiments, the antibody of the present invention specifically binds to integrin $\alpha_v \beta_3$. In some embodiments, the antibody is MEDI-522 (Vitaxin®). In certain embodiments, the antibody of the present invention specifically binds to CD2. In some embodiments, the antibody is siplizumab. In certain embodiments, the antibody of the present invention specifically binds to CD19. In some embodiments, the antibody is MT-103. In further embodiments, the antibody of the present invention specifically binds to EphA2. In some embodiments, the antibody is human or humanized EA2 or EA5. In certain embodiments, the antibody of the present invention specifically binds to EphA4. In some embodiments, the antibody is a humanized antibody that specifically binds to IL-9. In some embodiments, the antibody is a human or humanized antibody that specifically binds to IL-9. In some embodiments, the antibody is a human or humanized antibody that specifically binds to IL-9. In some embodiments, the antibody is a human or humanized antibody that specifically binds to IL-9. In some embodiments, the antibody is MEDI-528.

[00162] In some embodiments, the antibody is not palivizumab. In some embodiments, the antibody is not MEDI-522 (Vitaxin®). In some embodiments, the antibody is not siplizumab. In some embodiments, the antibody is not MT-103. In some embodiments, the antibody is not human or humanized EA2 or EA5. In some embodiments, the antibody is not MEDI-528.

[00163] The antibodies useful in the present invention may be high potency antibodies. High potency antibodies can be produced by genetically engineering appropriate antibody gene sequences and expressing the antibody sequences in a suitable host. The antibodies produced can be screened to identify antibodies with, e.g., high k_{on} values in a BIAcore assay.

[00164] In certain embodiments, the antibodies useful in the present invention have a high binding affinity for one or more antigens. In a specific embodiment, the antibodies of the present invention have an association rate constant or k_{on} rate (antibody (Ab) + antigen (Ag)^{k_{on}} \rightarrow Ab-Ag) of at least 10^5 M⁻¹s⁻¹, at least 5×10^5 M⁻¹s⁻¹, at least 10^6 M⁻¹s⁻¹, at least 10^7 M⁻¹s⁻¹, at least 5×10^7 M⁻¹s⁻¹, or at least 10^8 M⁻¹s⁻¹. In a preferred embodiment, the antibodies of the present invention have a k_{on} of at least 2×10^5 M⁻¹s⁻¹, at least 5×10^6 M⁻¹s⁻¹, at least 5×10^6 M⁻¹s⁻¹, at least 10^6 M⁻¹s⁻¹.

[00165] In another embodiment, the antibodies of the present invention have a k_{off} rate (antibody (Ab) + antigen) of less than 10^{-1} s⁻¹, less than 5×10^{-1} s⁻¹, less than 10^{-2} s⁻¹, less than 10^{-3} s⁻¹, less than 5×10^{-3} s⁻¹, less than 10^{-3} s⁻¹, less than 10^{-3} s⁻¹, less than 10^{-3} s⁻¹, less than 10^{-6} s⁻¹, less than 10^{-9} s⁻¹, less than 10^{-9} s⁻¹, or less than 10^{-10} s⁻¹. In a preferred embodiment, the antibodies of the present invention have a k_{on} of less than 10^{-6} s⁻¹, less than 10^{-

In certain embodiments, the antibodies of the present invention have an affinity constant or K_a (k_{on}/k_{off}) of at least 10^2 M⁻¹, at least 5×10^2 M⁻¹, at least 10^3 M⁻¹, at least 5×10^3 M⁻¹, at least 10^4 M⁻¹, at least 5×10^4 M⁻¹, at least 10^5 M⁻¹, at least 5×10^5 M⁻¹, at least 10^6 M⁻¹, at least 10^6 M⁻¹, at least 10^7 M⁻¹, at least 10^{10} M⁻¹, at least 10^{10} M⁻¹, at least 10^{11} M⁻¹, at least 10^{11} M⁻¹, at least 10^{12} M⁻¹, at least 10^{13} M⁻¹. The present invention also provides formulations comprising one or more antibodies which immunospecifically bind to an antigen with an affinity constant of at least 10^{10} M⁻¹, at

In yet another embodiment, the antibodies useful in the present invention have a dissociation constant or K_d (k_{off}/k_{on}) of less than 10^{-2} M, less than 5×10^{-2} M, less than 10^{-3} M, less than 10^{-4} M, less than 10^{-4} M, less than 10^{-5} M, less than 10^{-5} M, less than 10^{-5} M, less than 10^{-6} M, less than 10^{-10} M, less than 10^{-10} M, less than 10^{-10} M, less than 10^{-11} M, less than 10^{-11} M, less than 10^{-12} M, less than 10^{-12} M, less than 10^{-13} M, less than 10^{-13} M, less than 10^{-13} M, less than 10^{-13} M, less than 10^{-15} M, less than 10^{-15} M, or less than 10^{-15} M.

[00168] In certain embodiments, the antibodies useful in the present invention have a median effective concentration (EC₅₀) of less than 0.01 nM, less than 0.025 nM, less than 0.05 nM, less than 0.1 nM, less than 0.25 nM, less than 0.75 nM, less than 1 nM,

less than 1.25 nM, less than 1.5 nM, less than 1.75 nM, or less than 2 nM, in an *in vitro* microneutralization assay. The median effective concentration is the concentration of antibody or antibody fragments that neutralizes 50% of an antigen in an *in vitro* microneutralization assay. In a preferred embodiment, the antibodies of the present invention have an EC₅₀ of less than 0.01 nM, less than 0.025 nM, less than 0.05 nM, less than 0.1 nM, less than 0.25 nM, less than 1.5 nM, less than 1.75 nM, or less than 2 nM, in an *in vitro* microneutralization assay

The present invention also provides antibodies that immunospecifically bind to [00169] an antigen of interest, the antibodies comprising derivatives of the VH domains, VH CDRs, VL domains, and VL CDRs described herein that immunospecifically bind to antigens of interest. Standard techniques known to those of skill in the art can be used to introduce mutations in the nucleotide sequence encoding a molecule of the invention, including, for example, site-directed mutagenesis and PCR-mediated mutagenesis which results in amino acid substitutions. Preferably, the derivatives include less than 25 amino acid substitutions, less than 20 amino acid substitutions, less than 15 amino acid substitutions, less than 10 amino acid substitutions, less than 5 amino acid substitutions, less than 4 amino acid substitutions, less than 3 amino acid substitutions, or less than 2 amino acid substitutions relative to the original molecule. In a preferred embodiment, the derivatives have conservative amino acid substitutions are made at one or more predicted non-essential amino acid residues. A "conservative amino acid substitution" is one in which the amino acid residue is replaced with an amino acid residue having a side chain with a similar charge. Families of amino acid residues having side chains with similar charges have been defined in the art. These families include amino acids with basic side chains (e.g., lysine, arginine, histidine), acidic side chains (e.g., aspartic acid, glutamic acid), uncharged polar side chains (e.g., glycine, asparagine, glutamine, serine, threonine, tyrosine, cysteine), nonpolar side chains (e.g., alanine, valine, leucine, isoleucine, proline, phenylalanine, methionine, tryptophan), beta-branched side chains (e.g., threonine, valine, isoleucine) and aromatic side chains (e.g., tyrosine, phenylalanine, tryptophan, histidine). Alternatively, mutations can be introduced randomly along all or part of the coding sequence, such as by saturation mutagenesis, and the resultant mutants can be screened for biological activity to identify mutants that retain activity. Following mutagenesis, the encoded protein can be expressed and the activity of the protein can be determined.

5.3.3 Antibodies that Immunospecifically Bind to RSV antigen

[00170] It should be recognized that antibodies that immunospecifically bind to a RSV antigen are known in the art. For example, palivizumab is a humanized monoclonal antibody

presently used for the prevention of RSV infection in pediatric patients. The present invention provides formulations of antibodies that immunospecifically bind to one or more RSV antigens. Preferably, the antibodies useful in the invention immunospecifically bind to one or more RSV antigens regardless of the strain of RSV. The present invention also provides antibodies that differentially or preferentially bind to RSV antigens from one strain of RSV versus another RSV strain. In a specific embodiment, the formulations comprise antibodies that immunospecifically bind to the RSV F glycoprotein, G glycoprotein or SH protein. In a preferred embodiment, the formulations comprise antibodies that bind to the RSV F glycoprotein. In another preferred embodiment, the formulations comprise antibodies that bind to the A, B, or C antigenic sites of the RSV F glycoprotein.

[00171] The formulations of the invention comprise antibodies that immunospecifically bind to a RSV antigen and have a dissociation constant (K_D) of less than 3000 pM, less than 2500 pM, less than 2000 pM, less than 1500 pM, less than 1000 pM, less than 750 pM, less than 500 pM, less than 250 pM, less than 200 pM, less than 150 pM, less than 100 pM, less than 75 pM as assessed using an described herein or known to one of skill in the art (e.g., a BIAcore assay). In a specific embodiment, formulations of the invention comprise antibodies that immunospecifically bind to a RSV antigen and have a dissociation constant (K_D) of between 25 to 3400 pM, 25 to 3000 pM, 25 to 2500 pM, 25 to 2000 pM, 25 to 1500 pM, 25 to 1000 pM, 25 to 750 pM, 25 to 500 pM, 25 to 250 pM, 25 to 100 pM, 25 to 75 pM, 25 to 50 pM as assessed using an described herein or known to one of skill in the art (e.g., a BIAcore assay). In another embodiment, formulations of the invention comprise antibodies that immunospecifically bind to a RSV antigen and have a dissociation constant (K_D) of 500 pM, preferably 100 pM, more preferably 75 pM and most preferably 50 pM as assessed using an described herein or known to one of skill in the art (e.g., a BIAcore assay).

[00172] The present invention provides formulations that comprise antibodies that have a median inhibitory concentration (IC₅₀) of less than 5 nM, less than 4 nM, less than 3 nM, less than 2 nM, less than 1.75 nM, less than 1.5 nM, less than 1.25 nM, less than 1 nM, less than 0.75 nM, less than 0.5 nM, less than 0.25 nM, less than 0.01 nM, less than 0.05 nM, less than 0.025 nM, or less than 0.01 nM, in an *in vitro* microneutralization assay. The IC₅₀ is the concentration of antibody that neutralizes 50% of the RSV in an *in vitro* microneutralization assay. In a preferred embodiment, antibody of the invention has an IC₅₀ of less than 5 nM, less than 4 nM, less than 3 nM, less than 2 nM, less than 1.75 nM, less than 1.5 nM, less than 1.25 nM, less than

1 nM, less than 0.75 nM, less than 0.5 nM, less than 0.25 nM, less than 0.1 nM, less than 0.05 nM, less than 0.025 nM, or less than 0.01 nM, in an *in vitro* microneutralization assay.

In a specific embodiment, the formulations of the invention comprise an antibody [00173] that has approximately 20-fold, 25-fold, 30-fold, 35-fold, 40-fold, 45-fold, 50-fold, 55-fold, 60fold, 65-fold, 70-fold, 75-fold, 80-fold, 90-fold, 100-fold or higher affinity for a RSV F antigen than palivizumab or an antibody-binding fragment thereof as assessed by an assay known in the art or described herein (e.g., a BIAcore assay). In another embodiment, formulations of the invention comprise antibodies that have an approximately 1-fold, 1.5-fold, 2-fold, 3-fold, 4-fold, 5-fold, or more higher Ka than palivizumab or an antigen-binding fragment thereof as assessed by an assay known in the art or described herein. In another embodiment, a formulation of the invention comprises an antibody that is approximately 1-fold, 2-fold, 3-fold, 4-fold, 5-fold, 6fold, 7-fold, 8-fold, 9-fold, 10-fold, 11-fold 12-fold, 13-fold, 14-fold, 15-fold, 16-fold, 17-fold, 18-fold, 19-fold, or 20-fold or more potent than palivizumab or an antigen-binding fragment thereof in an in vitro microneutralization assay such as described herein. The amino acid sequence of palivizumab is disclosed, e.g., in Johnson et al., 1997, J. Infectious Disease 176:1215-1224, and U.S. Patent No. 5,824,307, each of which is incorporated herein by reference in its entirety. In a specific embodiment, a formulation of the invention comprise an antibody that is not palivizumab or a fragment of palivizumab or an antigen-binding fragment of palivizumab, e.g., is not an antibody comprising a VH domain of SEQ ID NO:7 and/or a VL domain of SEQ ID NO:8.

[00174] The present invention provides antibodies that immunospecifically bind to one or more RSV antigens, said antibodies comprising the amino acid sequence of palivizumab with one or more amino acid residue substitutions in the variable light (VL) domain and/or variable heavy (VH) domain depicted in FIG. 3. The present invention also provides antibodies that immunospecifically bind to one or more RSV antigens, said antibodies comprising the amino acid sequence of palivizumab with one or more amino acid residue substitutions in one or more VL CDRs and/or one or more VH CDRs. In a specific embodiment, an antibody comprises the amino acid sequence of palivizumab with one or more amino acid residue substitutions of the amino acid residues indicated in bold face and underlining in Table 1. In another embodiment, an antibody comprises the amino sequence of palivizumab with one or more amino acid residue substitutions of the amino acid residues indicated in bold face and underlining in Table 1 and one or more amino acid residue substitutions of the framework regions of the variable domains of palivizumab (e.g., mutations in framework region 4 of the heavy and/or light variable

domains). In accordance with these embodiments, the amino acid residue substitutions can be conservative or non-conservative. The antibody generated by introducing substitutions in the VH domain, VH CDRs, VL domain and/or VL CDRs of palivizumab can be tested *in vitro* and *in vivo*, for example, for its ability to bind to RSV F antigen, for its ability to neutralize RSV, or for its ability to prevent, treat or ameliorate an upper and/or lower respiratory tract RSV infection, otitis media, or a symptom or respiratory condition relating thereto (including, but not limited to, asthma, wheezing, RAD, or a combination thereof).

Table 1. CDR Sequences of palivizumab

CDR	Sequence	SEQ ID NO:
VH1	T <u>S</u> GMSVG	1
VH2	DIWWD D K KD YNPSLK S	. 2
VH3	$\underline{\mathbf{S}}$ MI $\underline{\mathbf{T}}$ N $\underline{\mathbf{W}}$ YFDV	3
VL1	KCQLS VGYMH	4
VL2	DT <u>SKLA</u> S	5
VL3	$FQGS\underline{\mathbf{G}}YP\underline{\mathbf{F}}T$	6

^{*} Bold faced & underlined amino acid residues are preferred residues which should be substituted.

[00175] The formulations of the present invention also comprise those antibodies and antigen-binding fragments of the antibodies referenced in Table 2 and the Examples Section of the application. In a specific embodiment, a formulation of the present invention comprises antibody AFFF, P12f2, P12f4, P11d4, Ale9, A12a6, A13c4, A17d4, A4B4, A8c7, 1X-493L1FR, H3-3F4, M3H9, Y10H6, DG, AFFF(1), 6H8, L1-7E5, L2-15B10, A13a11, A1h5, A4B4(1), A4B4L1FR-S28R (motavizumab), A4B4-F52S, A17d4(1), A3e2, A14a4, A16b4, A17b5, A17f5, or A17h4. In another embodiment, a formulation of the present invention comprises an antigen-binding fragment (*e.g.*, a Fab fragment of) AFFF, P12f2, P12f4, P11d4, Ale9, A12a6, A13c4, A17d4, A4B4, A8c7, 1X-493L1FR, H3-3F4, M3H9, Y10H6, DG, AFFF(1), 6H8, L1-7E5, L2-15B10, A13a11, A1h5, A4B4(1), A4B4L1FR-S28R (motavizumab), A4B4-F52S, A17d4(1), A3e2, A14a4, A16b4, A17b5, A17f5, or A17h4. In a preferred embodiment, a formulation of the present invention comprises antibody A4B4L1FR-S28R or an antigen-binding fragment thereof.

[00176] In some embodiments, AFFF, P12f2, P12f4, P11d4, Ale9, A12a6, A13c4, A17d4, A4B4, A8c7, 1X-493L1FR, H3-3F4, M3H9, Y10H6, DG, AFFF(1), 6H8, L1-7E5, L2-15B10, A13a11, A1h5, A4B4(1), A4B4L1FR-S28R, A4B4-F52S, A17d4(1), A3e2, A14a4, A16b4, A17b5, A17f5, and/or A17h4 comprise the framework region and constant regions of palivizumab (see FIG. 3). In preferred embodiments, AFFF, P12f2, P12f4, P11d4, Ale9, A12a6, A13c4, A17d4, A4B4, A8c7, 1X-493L1FR, H3-3F4, M3H9, Y10H6, DG, AFFF(1), 6H8, L1-7E5, L2-15B10, A13a11, A1h5, A4B4(1), A4B4L1FR-S28R (motavizumab), A4B4-F52S, A17d4(1), A3e2, A14a4, A16b4, A17b5, A17f5, and/or A17h4 comprise the framework region and constant regions of palivizumab with the exception that there is an amino acid substitution of an A105Q in the heavy chain framework 4 (FR4) (numbering used herein according to Kabat et al. (1991). Sequences of proteins of immunological interest. (U.S. Department of Health and Human Services, Washington, D.C.) 5th ed.) ("Kabat numbering") (i.e., position 112 in SEQ ID NO:7 (palivizumab VH domain)) and an L104V in the light chain FR4 (i.e., position 103 in SEO ID NO:8 (palivizumab VL domain)). An example of antibodies comprising a framework with these VH and VL single mutations is shown in FIG. 4 (1X-493L1FR) and in FIG. 5 (A4B4L1FR-S28R).

[00177] In a specific embodiment, the present invention provides one or more antibodies that immunospecifically bind to one or more RSV F antigens, said antibodies comprising a VH chain and/or VL chain having the amino acid sequence of a VH chain and/or VL chain of AFFF, P12f2, P12f4, P11d4, Ale9, A12a6, A13c4, A17d4, A4B4, A8c7, 1X-493L1FR, H3-3F4, M3H9, Y10H6, DG, AFFF(1), 6H8, L1-7E5, L2-15B10, A13a11, A1h5, A4B4(1), A4B4L1FR-S28R, A4B4-F52S, A17d4(1), A3e2, A14a4, A16b4, A17b5, A17f5, or A17h4. In a preferred embodiment, an antibody of the invention immunospecifically binds to a RSV F antigen, and said antibody comprises a VH chain and/or a VL chain having the amino acid sequence of the VH and/or VL chain of A4B4L1FR-S28 (VH chain, SEO ID NO:254; VL chain SEO ID NO:255). In another embodiment, the present invention provides one or more antibodies that immunospecifically bind to one or more RSV antigens, said antibodies comprising a VH domain and/or VL domain having the amino acid sequence of a VH domain and/or VL domain of AFFF, P12f2, P12f4, P11d4, Ale9, A12a6, A13c4, A17d4, A4B4, A8c7, 1X-493L1FR, H3-3F4, M3H9, Y10H6, DG, AFFF(1), 6H8, L1-7E5, L2-15B10, A13a11, A1h5, A4B4(1), A4B4L1FR-S28R, A4B4-F52S, A17d4(1), A3e2, A14a4, A16b4, A17b5, A17f5, or A17h4. In a preferred embodiment, an antibody of the invention immunospecifically binds to a RSV F antigen, and said antibody comprises a VH domain and/or VL domain having the amino acid sequence of the

VH domain and/or VL domain of A4B4L1FR-S28R (VH domain, SEQ ID NO:48; VL domain, SEQ ID NO:11).

In another embodiment, the present invention provides antibodies that [00178] immunospecifically bind to one or more RSV antigens, said antibodies comprising one, two, three, or more CDRs having the amino acid sequence of one, two, three, or more CDRs of AFFF, P12f2, P12f4, P11d4, Ale9, A12a6, A13c4, A17d4, A4B4, A8c7, 1X-493L1FR, H3-3F4, M3H9, Y10H6, DG, AFFF(1), 6H8, L1-7E5, L2-15B10, A13a11, A1h5, A4B4(1), A4B4L1FR-S28R (MEDI-524, motavizumab), A4B4-F52S, A17d4(1), A3e2, A14a4, A16b4, A17b5, A17f5, or A17h4. In a preferred embodiment, a formulation of the present invention comprises an antibody that immunospecifically binds to a RSV antigen, and said antibody comprises one, two, three, or more CDRs having the amino acid sequence of one, two, three, or more CDRs of A4B4L1FR-S28R. In yet another embodiment, the formulation of the present invention comprises an antibody that immunospecifically binds to one or more RSV F antigens, said antibodies comprising a combination of VH CDRs and/or VL CDRs having the amino acid sequence of VH CDRs and/or VL CDRs of AFFF, P12f2, P12f4, P11d4, Ale9, A12a6, A13c4, A17d4, A4B4, A8c7, 1X-493L1FR, H3-3F4, M3H9, Y10H6, DG, AFFF(1), 6H8, L1-7E5, L2-15B10, A13a11, A1h5, A4B4(1), A4B4L1FR-S28R (motavizumab), A4B4-F52S, A17d4(1), A3e2, A14a4, A16b4, A17b5, A17f5, and/or A17h4. In a preferred embodiment, a formulation of the present invention comprises an antibody that immunospecifically binds to a RSV F antigen and said antibody comprises a combination of VH CDRs and/or VL CDRs having the amino acid sequence of the VH CDRs and/or VL CDRs of A4B4L1FR-S28R.

[00179] The present invention provides antibodies that immunospecifically bind to one or more RSV antigens (e.g., RSV F antigen), said antibodies comprising a variable heavy ("VH") chain having an amino acid sequence of any one of the VH chains listed in Table 2. In certain embodiments, the antibody is not palivizumab and/or the VH chain is not the VH chain of palivizumab.

[00180] The invention also provides antibodies that immunospecifically bind to one or more RSV antigens (e.g., RSV F antigen), said antibodies comprising a VH domain having an amino acid sequence of any one of the VH domains listed in Table 2. In certain embodiments of the invention, the antibody is not palivizumab and/or the VH domain is not the VH domain of palivizumab.

[00181] The present invention also provides antibodies that immunospecifically bind to one or more RSV antigens, said antibodies comprising a VH complementarity determining region ("CDR") (e.g., VH CDR1, VH CDR2, and/or VH CDR3) having an amino acid sequence of any of the VH CDRs listed in Table 2 and/or Tables 3A-3C. In certain embodiments of the invention, an antibody comprising a VH CDR having an amino acid of any of one of the VH CDRs listed in Table 2 and/or Tables 3A-3C is not palivizumab. In some embodiments, the antibody or binding fragment thereof comprises one, two or three of the VH CDRs listed in Table 2 and/or Tables 3A-3C.

Table 2. Antibodies & Fragments Thereof

2007 ENG TA	700\dag	2543	FQ <u>F</u> S模型PFT (SEQ TD NO:확卷)	FQGSĞ¥PFT SEQ ID NO:6)	FQGSGYPFT SEQ ID NO:6)	FQGSGYPFT SEQ ID NO:6)	FQGSGYPET SEQ ID NO:6)	FQGSGYPET SEQ ID NO:6)	FQGSGYPFT SEQ ID NO:6)	FOGSGYPFT SEQ ID NO:6)	EQGSGYPET (SEQ ID NO:64	S20 CLEALES OSS) CLEALES OSS	GYPE19 D NO: (0/9)
ZA.	FOGS		FOES (SE NO:	FQGS (SEQ I	FQGS (SEQ I	EQGS (SEQ I	EQGS (SEQ I	I ČES)	FQGS (SEQ I	FQGS (SEQ I	FQGS (SEQ I	FQGS (SEQ I	
VL CDR2	DTSKTAS	(SEQ ID NO:5)	DTFKLAS (SEQ ID NO:15)	DTFYLSS (SEQ ID NO:23)	DTRGLPS (SEQ ID NO:27)	DTMRLAS (SEQ ID NO:32)	DTFKLSS (SEQ ID NO:35)	DTFKLSS (SEQ ID NO:35)	DTMYQSS (SEQ ID NO:43)	DIMYOSS (SEQ ID NO:43)	DTFFLDS (SEQ ID NO:50)	DTRYQSS (SEQ ID NO:53)	DTSKLAS
VL CDR1	KCOLSVCVMH	(SEQ ID NO:4)	SASSSVGYMH (SEQ ID NO:14)	SLSSRVGYMH (SEQ ID NO:22)	SISSRVGYMH (SEQ ID NO:22)	SPSSRVGYMH (SEQ ID NO:31)	SLSSRVGYMH (SEQ ID NO:22)	SASSRVGYMH (SEQ ID NO:39)	SLSSRVGYMH (SEQ ID NO:22)	LPSSRVGYMH (SEQ ID NO:47)	SASSRVGYMH (SRQ TD NO:39)	SPSSRVGYMH (SEQ ID NO:31)	SASSSVGYMH (SEC ID NO:14)
ZV.	OFO TD	8:0 8:0 8:0 8:0	SEQ ID NO:13	SEQ ID NO:21	SEQ ID NO:26	SEQ ID NO:30	SEQ ID NO:34	SEQ ID NO:38	SEQ ID NO:42	SEQ ID NO:46	SEQ ID NO:49	SEQ ID NO:52	SEQ ID
Δ. A.L.	Chain	SEX ID NO:209	SEQ ID NO:211	SEQ ID NO:213	SEQ ID NO:215	SEQ ID NO:217	SEQ ID NO:219	SEQ ID NO:221	SEQ ID NO:223	SEQ ID NO:225	SEQ ID NO:227	SEQ ID NO:229	SEQ ID
VH CDR3	OMTHUMAVERY	(SEQ ID NO:3)	SMITNEYEDV (SEQ ID NO:12)	DMIFNETDV (SEQ ID NO:20)	DMIFNEYEDV (SEQ ID NO:20)	DMIFNWYEDV (SEQ ID NO:29)	DMIENWYEDV (SEQ ID NO:29)	DMIFNEYEDV (SEQ ID NO:20)	DMIENEYEDV (SEQ ID NO:20)	DMLFNFYFDV (SEQ ID NO:20)	DMIENEYEDV (SEQ ID NO:20)	DMIENWYEDV (SEQ ID NO:29)	SMITHWWYFDV
VH CDR2	NYCHARACT	PSLKS (SEO ID NO:2)	DIWWDDKKDYN PSIKS (SEO ID NO:2)	DIWWDDKKHYN PSLKD (SEQ ID NO:19)	DIWWDGKKHYN PSLKD (SEO ID NO:25)	DIWWDGKKHYN PSIKD (SEO ID NO:25)	DIWWDGKKHYN PSIKD (SEO ID NO:25)	DIWWDGKKDYN PSLKD (SEQ ID NO:37)	DIWWDGKKSYN PSLKD (SEQ ID NO:41)	DIWWDDKKSYN PSLKD (SEQ ID NO:45)	DIWWDDKKHYN PSLKD (SEQ ID NO:19)	DIWWDDKKSYN PSLKD (SEQ ID NO:45)	DIWWDDKKDYN
VH CDR1	OTYONOOD	SEQ ID NO:1)	TAGMSVG (SEQ_ID NO:10)	TPGMSVG (SEQ_ID_NO:18)	TPGMSVG (SEQ_ID NO:18)	TPGMSVG (SEQ_ID NO:18)	TAGMSVG (SEQ_ID_NO:10)	TAGMSVG (SEQ_ID NO:10)	TAGMSVG (SEQ ID NO:10)	TACMSVG (SEQ_ID NO:10)	TAGMSVG (SEQ ID NO:10)	TAGMSVG (SEQ_ID_NO:10)	TSGMSVG
AH.	Domain	NO:7	SEQ ID NO:9	SEQ ID NO:17	SEQ ID NO:24	SEQ ID NO:28	SEQ ID NO:33	SEQ ID NO:36	SEQ ID NO:40	SEQ ID NO:44	SEQ ID NO:48	SEQ ID NO:51	SEQ ID
ΗΛ	Chain	SEQ 1D NO:208	SEQ ID NO:210	SEQ ID NO:212	SEQ ID NO:214	SEQ ID NO:216	SEQ ID NO:218	SEQ ID NO:220	SEQ ID NO:222	SEQ ID NO:224	SEQ ID NO:226	SEQ ID NO:228	SEQ ID
Antibody VH VH CI	Name	**palivizum ab	***AFFF	***P12f2	***P12f4	***P11d4	***A1e9	***A12a6	***A13c4	***A17d4	***A4B4	****A8c7	*1X-

VO_200	7/0 025 4	3 6 13	£T :6)	L	F. (9)	FT : 6)	ET:	ET: 6)	1. 1.6)	[6 P €	CT/US200	6/0 24 7
EQGSETPE (SEQ IN NO:	7/0 025 4 //O03:0N ⁻ -QI OSS)	EQGSG¥PEJG (SEQ ID NO:6)	EQGS(SYPET) (SEQ ID-NO:6)	FQGS <u>E</u> YPFT (SEQ ID NO:62)	FQGSGYPFT (SEQ ID NO:6)	FQGSGYPFT (SEQ ID NO:6)	FQGSGYPFT (SEQ ID NO:6)	FQGSGYPFT (SEQ ID NO:6)	FQGSGYPFT (SEQ ID NO:6)	FQGSGYPFT (SEQ ID NO:6)	CT/US200 T/US:00 CI CES)	FQGSGYPI (SEQ ID NO
DTFKLAS (SEQ ID NO:15)	DTYK QT S (SEQ ID NO:57)	DTRYLSS (SEQ ID NO:59)	DTFKLAS (SEQ ID NO:15)	DTFKLAS (SEQ ID NO:15)	DTFKLTS (SEQ ID NO:63)	DIFKLAS (SEQ ID NO:15)	DTFRLAS (SEQ ID NO:66)	DTYRHSS (SEQ ID NO:69)	DIFFHRS (SEQ ID NO:73)	DTILIDS (SEQ ID NO:75)	DTSKLAS (SEQ ID NO:5)	DISELDS (SEQ ID NO:77)
SASSSVGYMH (SEQ ID NO:14)	SASSSVGYMH (SEQ ID NO:14)	SASSSVGYMH (SEQ ID NO:14)	SASSSVGYMH (SEQ ID NO:14)	SASSSVGYMH (SEQ ID NO:14)	SASSSVGYMH (SEQ ID NO:14)	SASSRVGYMH (SEQ ID NO:39)	SASSSVGYMH (SEQ ID NO:14)	SPSSRVGYMH (SEQ ID NO:31)	SLSSSVGYMH (SEQ ID NO:72)	SASSRVGYMH (SEQ ID NO:39)	SASSEVGYMH (SEQ ID NO:39)	SASSRVGYMH (SEQ ID NO:39)
SEQ ID	SEQ ID	SEQ ID	SEQ ID	SEQ ID	SEQ ID	SEQ ID	SEQ ID	SEQ ID	SEQ ID	SEQ ID	SEQ ID	SEQ ID
NO:56	NO:70	NO:58	NO:56	NO:60	NO:62	NO:64	NO:65	NO:68	NO:71	NO:74	NO:11	NO:76
SEQ ID	SEQ ID	SEQ ID	SEQ ID	SEQ ID	SEQ ID	SEQ ID	SEQ ID	SEQ ID	SEQ ID	SEQ ID	SEQ ID	SEQ ID
NO:233	NO:235	NO:237	NO:239	NO:241	NO:243	NO:245	NO:247	NO:249	NO:251	NO:253	NO:255	NO:257
DMIFNWYEDV	DMIENWYEDV	DMIFNWYEDV	DMITNEYEDV	SMITNEYEDV	DMITNEYEDV	DMITNEYEDV	DMITNEYEDV	DMLENWYFDV	DMIENWYEDV	DMIENEYEDV	DMIENEYEDV	DMIENEYEDV
(SEQ ID NO:29)	(SEQ ID NO:29)	(SEQ ID NO:29)	(SEQ ID NO:79)	(SEQ ID NO:12)	(SEQ ID NO:79)	(SEQ ID NO:79)	(SEQ ID NO:79)	(SEQ ID NO:29)	(SEQ ID NO:29)	(SEQ ID NO:20)	(SEQ ID NO:20)	(SEQ ID NO:20)
DIWWDDKKDYN PSLKS (SEQ ID NO:2)	DIWWDDKKDYN PSLKS (SEQ ID NO:2)	DIWWDDKKDYN PSLKS (SEQ ID NO:2)	DIWWDDKKDYN PSLKS (SEQ ID NO:2)	DIWWDDKKDYN PSLKS (SEQ ID NO:2)	DIWWDDKKDYN PSLKS (SEQ ID NO:2)	DIWWDDKKDYN PSLKS (SEQ ID NO:2)	DIWWDDKKDYN PSLKS (SEQ ID NO:2)	DIWWDDKKHYN PSLKD (SEQ ID NO:19)	DIWWDGKKHYN PSLKD (SEQ ID NO:25)	DIWWDDKKHYN PSLKD (SEO ID NO:19)	DIWWDDKKHYN PSLKD (SEQ ID NO:19)	DIWWDDKKHYN PSIKD PSIKD
TAGMSVG	TAGMSVG	TAGMSVG	TAGMSVG	TAGMSVG	TAGMSVG	TAGMSVG	TAGMSVG	TAGMSVG	TAGMSVG	TAGMSVG	TAGMSVG	TAGMSVG
(SEQ ID NO:10)	(SEQ ID NO:10)	(SEQ ID NO:10)	(SEQ ID NO:10)	(SEQ_ID NO:10)	(SEQ ID NO:10)	(SEQ_ID_NO:10)	(SEQ_ID_NO:10)	(SEQ ID NO:10)	(SEQ_ID_NO:10)	(SEQ ID NO:10)	(SEQ ID NO:10)	(SEQ ID NO:10)
SEQ ID	SEQ ID	SEQ ID	SEQ ID	SEQ ID	SEQ ID	SEQ ID	SEQ ID	SEQ ID	SEQ ID	SEQ ID	SEQ ID	SEQ ID
NO:55	NO:55	NO:55	NO:78	NO:9	NO:78	NO:78	NO:78	NO:67	NO:33	NO:48	NO:48	NO:48
SEQ ID	SEQ ID	SEQ ID	SEQ ID	SEQ ID	SEQ ID	SEQ ID	SEQ ID	SEQ ID	SEQ ID	SEQ ID	SEQ ID	SEQ ID
NO:232	NO:234	NO:236	NO:238	NO:240	NO:242	NO:244	NO:246	NO:248	NO:250	NO:252	NO:254	NO:256
*H3-3E4	*M3H9	*Y10H6	*DG (aka D95/G93)	AFFF(1)	×6H8	*L1-7E5	*L2-15B10	*A13a11	*Alh5	A4B4(1)	***A4B4L1F R-S28R (aka motavizumab)	***A4B4- F52S

VO 20	07/ に		25 ₄	13 - E	o o		(6		E.J	(9:		ΕŢ	(9:		H H	(9:	
VIJAZOSOA	FOGSETET	(SEQ ID-NO:6	i testi ligar	FOGSCHPFTS	(P:ON TOTS)	, 10000	EQGSGYPT	(SEQ ID NO:6)		FQGSGYPET	(SEQ ID NO:6)		FQGSGYPFT	(SEQ ID NO:6)		FQGSGYPET	SEQ LD NO:67	
DTMYOSS (SEQ ID NO:43)	DIFYLHS	(SEQ ID	NO:308)	DIXXOIS	(SEQ 1D NO:315)		DIMYOAS	(SEQ 1D	MO. 321)	DIXXIBS	(SEQ ID	NO:326)	DTFRHTS	(SEQ ID	1755:ON	DTYYLAS	CEÇ ID	(acc:ON
LPSSRVGYMH (SEQ ID NO:47)	SASSSUGYMH	(SEQ ID NO:14)		LLSSRVGYMH	(SEQ ID NO:314)		LLSSRVGYMH	(SEQ ID NO:320)		SISSRVGYMH	(SEQ ID NO:22)		SLSSRVGYMH	(SEQ ID NO:22)		SPSSSVGYMH	(SEQ ID NO:335)	
SEQ ID NO:46	SEO TD	NO: 307	•	SEQ ID	NO:313		SEQ ID	NO:319		SEQ ID	NO:325		SEQ ID	NO:331		SEQ ID	NO:334	
SEQ ID NO:225	SEO TD	NO - 306		SEQ ID	NO:312		SEQ ID	NO:318		SEQ ID	NO:324		SEQ ID	NO:330		SEQ ID	NO:333	
DMIENEYFDV (SEQ ID NO:20)	DMTENMY	(SEO ID NO:29)	() () () () () () () () () ()	DMITMAKEDA	(SEQ ID NO:311)		DMIENWYFDV	(SEQ ID NO:29)		DMIENWYFDV	(SEQ ID NO:29)		DMIFNWYFDV	(SEQ ID NO:29)		DMIENWYFDV	(SEQ ID NO:29)	
DIWWDGKKSYN PSLKD	(SEQ ID NO:41)	TA ISO	(SEO ID NO:305)	DIWWDDKKSYN	PSLKD	(SEQ ID NO:45)	DIWWDDKKHYN	PSLKD	(SEQ ID NO:19)	DIWWDDKKHYN	PSTKD_	(SEQ ID NO:19)	DIWWDDKKDYN	PSLKD	(SEQ ID NO:329)	DIWWDGKKHYN	PSLKD	(SEQ ID NO:25)
TAGMSVG (SEQ ID NO:10)	OZYONO KIII	LAGMOVG	(סדיסא קד אקני)	TAGMSVG	(SEQ ID NO:10)		TAGMSVG	(SEQ ID NO:10)		TAGMSVG	(SEQ_ID NO:10)		TAGMSVG	(SEQ_ID NO:10)		TAGMSVG	(SEQ_ID NO:10)	
SEQ ID NO:40	6	OEQ ID	NO:304	SEQ ID	NO:310		SEO ID	NO:317		SEO ID	MO:323		SEO ID	NO:328		SEO ID	NO: 33	
SEQ ID NO:222	1	OT OHS	NO:303	SEO ID	NO:309		SEO ID	NO:316		SEO ID	NO 322		SEO TD	NO:327	1	SEO ID	NO:218	
***A17d4(1)		***A36Z		***A14a4			***A16b4			***A17h5	>		*** \\ \\ \	2		***A17h4		

Bold faced & underlined amino acid residues are the residues which differ from the amino acid sequence in palivizumab; Fab fragment produced (**); Monoclonal antibody produced (**); Fab fragment & monoclonal antibody produced (***)

Table 3A - VH CDR1 Sequences

NO:1)	NO:10)	NO:18)
NO:1)	7:	
	_	ID N(
QI (OI (
(SEQ	OES)	(SEO
SVG	SVG	SVG

Bold faced & underlined amino acid residues are the residues which differ from the amino acid sequence in palivizumab

Table 3B - VH CDR2 Sequences

Table 3D - VII CDIVE Schacifics			
DKKDYNPSLKS	DKKDYNPSLKS (SEQ ID NO:2)	GKKDYNPSLKS (SEQ ID NO:100)	SEQ ID NO:100)
DKKDYNPSLKD	DKKDYNPSIK D (SEQ ID NO:86)	GKKDYNPSLKD (SEQ ID NO:103)	SEQ ID NO:103)
DKKHYNPSLKS	DKKHYNPSLKS (SEQ ID NO:82)	GKKHYNPSLKS (SEQ ID NO:106)	SEQ ID NO:106)
DKKHYNPSLKD	DKKHYNPSIKD (SEQ ID NO:19)	GKKHYNPSLKD (SEQ ID NO:25)	SEQ ID NO:25)
DKKSYNPSLKS	DKKSYNPSLKS (SEQ ID NO:109)	GKKSYNPSLKS (SEQ ID NO:114)	SEQ ID NO:114)
DKKSYNPSIKD	DKKKSYNPSLKD (SEQ ID NO:111)	GKKSYNPSLKD (SEQ ID NO:41)	SEQ ID NO:41)
DKGDYNPSLKS	DKGDYNPSLKS (SEQ ID NO:384)	GKGDYNPSLKS (SEQ ID NO:390)	SEQ ID NO:390)
DKGDYNPSLKD	DDKGDYNPSLKD (SEQ ID NO:385)	GKGDYNPSLKD (SEQ ID NO:391)	SEQ ID NO:391)
DKGHYNPSLKS	DDKGHYNPSLKS (SEQ ID NO:386)	OGKGHYNPSLKS (SEQ ID NO:392)	SEQ ID NO:392)
DKGHYNPSLKD	DDKGHYNPSLKD (SEQ ID NO:387)	GKGHYNPSIKD (SEQ ID NO:393)	SEQ ID NO:393)
DKGSYNPSLKS	DDKGSYNPSLKS (SEQ ID NO:388)	GKGSYNPSLKS (SEQ ID NO:394)	SEQ ID NO:394)
DKGSYNPSIKD	DKGSYNPSLKD (SEQ ID NO:389)	GKGSYNPSLKD (SEQ ID NO:395	SEQ ID NO:395)

Bold faced & underlined amino acid residues are the residues which differ from the amino acid sequence in palivizumab

Table 3C - VH CDR3 Sequences

Laule	Table 3C - VII CDING Sequences		
NWYFDV	(SEQ ID NO:3)	NWYFDV	NWYEDV (SEQ ID NO:83)
NEYFDV	(SEQ ID NO:12)	NEYEDV	NEYFDV (SEQ ID NO:29)
NWYFDV	(SEQ ID NO:94)	NWYFDV	NWYFDV (SEQ ID NO:79)
NEYFDV	(SEQ ID NO:97)	NEYEDV	NEYEDV (SEQ ID NO:20)
			•

Bold faced & underlined amino acid residues are the residues which differ from the amino acid sequence in palivizumab

Table 3D - VL CDRI Sequences	1 Sequences			
KCOLSVGYMH (SEQ	Q ID NO:4)	SCQLSVGYMH (SEQ ID NO:127)	LCOLSVGYMH ((SEQ ID NO:204)
	0:87	RVGYMH (SEQ ID NO:132)	RVGYMH (SEQ I	ID NO:206)
KCQL F VGYMH (SEQ	Q ID NO:396)	SCQLFVGYMH (SEQ ID NO:436)	TCOTEACEMH ((SEQ ID NO:476)
KCOSSVGYMH (SEQ	Q ID NO:80)	SCOSSVGYMH (SEQ ID NO:129)	TCOSSAGAMH ((SEQ ID NO:205)
KCQSRVGYMH (SEQ	Q ID NO:84)	SCOSRVGYMH (SEQ ID NO:130)	LCQSRVGYMH (ID NO
KCQSFVGYMH (SEQ	Q ID NO:397)	SCOSFVGYMH (SEQ ID NO:437)	LCOSEVGYMH ((SEQ ID NO: 477)
1	Q ID NO:398)	SCQVSVGYMH (SEQ ID NO:438)		ID NO:47
	Q ID NO:399)	SCQVRVGYMH (SEQ ID NO: 439)		(SEQ ID NO: 479)
KCQVEVGYMH (SEQ	Q ID NO:400)	SCQVFVGYMH (SEQ ID NO:440)	LCQVFVGYMH ((SEQ ID NO: 480)
KCSLSVGYMH (SEQ	Q ID NO:112)	SCSLSVGYMH (SEQ ID NO:142)	ICSLSVGYMH ((SEQ ID NO:196)
	Q ID NO:119)	SCSLRVGYMH (SEQ ID NO:148)	$^{\prime}$	(SEQ ID NO:198)
KCSLFVGYMH (SEQ	Q ID NO:401)	SCSLFVGYMH (SEQ ID NO:441)	LCSLEVGYMH ((SEQ ID NO:481)
KCSSSVGYMH (SEQ	Q ID NO:115)	SCSSSVGYMH (SEQ ID NO:144)	LCSSSVGYMH ((SEQ ID NO:197)
	Q ID NO:117)	SCSSRVGYMH (SEQ ID NO:146)	LCSSRVGYMH ((SEQ ID NO:195)
1	Q ID NO:402)	SCSSFVGYMH (SEQ ID NO:442)	LCSSFVGYMH ((SEQ ID NO:482)
KCSVSVGYMH (SEQ	Q ID NO:403)	SCSVSVGYMH (SEQ ID NO:443)	TCSNSYMH ((SEQ ID NO:483)
1	Q ID NO:404)	SCSVRVGYMH (SEQ ID NO:444)	LCSVRVGYMH ((SEQ ID NO:484)
KCSVFVGYMH (SEQ	Q ID NO:405)	SCSVFVGYMH (SEQ ID NO:445)	LCSVEVGYMH ((SEQ ID NO:485)
	Q ID NO:182)	SAQLSVGYMH (SEQ ID NO:207)	TYOTSAGAMH	(SEQ ID NO:486)
	Q ID NO:180)	SAQLRVGYMH (SEQ ID NO:190)	LAQLRVGYMH ((SEQ ID NO:487)
KAQLFVGYMH (SEQ	Q ID NO:406)	SAQLEVGYMH (SEQ ID NO:446)	I.AQLEVGYMH	(SEQ ID NO:488)
KAQSSVGYMH (SEQ	Q ID NO:181)	SAQSSVGYMH (SEQ ID NO:191)		(SEQ ID NO: 489)
	Q ID NO:179)	SAQSRVGYMH (SEQ ID NO:189)		(SEQ ID NO:490)
KAQSFVGYMH (SEQ	Q ID NO:407)	SAQSEVGYMH (SEQ ID NO:447)	LAOSEVGYMH ((SEQ ID NO:491)
KAQVSVGYMH (SEQ	Q ID NO:408)	SAQVSVGYMH (SEQ ID NO:448)		(SEQ ID NO:492)
	Q ID NO:409)	SAQVRVGYMH (SEQ ID NO:449)	LAQVRVGYMH ((SEQ ID NO: 493)
KAQVEVGYMH (SEQ	Q ID NO:410)	SAQVEVGYMH (SEQ ID NO:450)	LAQVEVGYMH ((SEQ ID NO: 494)
KASLSVGYMH (SEQ	Q ID NO:186)	SASLSVGYMH (SEQ ID NO:188)	LASISVGYMH ((SEQ ID NO: 495)
1	Q ID NO:184)	SASLRVGYMH (SEQ ID NO:187)	LASLRVGYMH ((SEQ ID NO:496)
1	TD	SASLEVGYMH (SEQ ID NO:451)	LASLEVGYMH ((SEQ ID NO: 497)
KASSSVGYMH (SEQ	Q ID NO:185)	SASSSVGYMH (SEQ ID NO:14)	LASSSVGYMH ((SEQ ID NO:498)

		TMAUMAUMAU	(SE-ON OF CHS)	LASSRVGYMH (SEO LD NO:499)
KASSEVGYMH ((SEO ID NO: 412)		ID NO:4	LASSFVGYMH (SEQ ID NO:500)
1	Qi		(SEQ ID NO: 453)	LASVSVGYMH (SEQ ID NO:501)
	ID NO:41	SASVRVGYMH ((SEQ ID NO:454)	LASVRVGYMH (SEQ ID NO: 502)
	(SEQ ID NO:415)	SASVEVGYMH ((SEQ ID NO: 455)	LASVEVGYMH (SEQ ID NO:503)
	(SEQ ID NO:89)	STOLSVGYMH ((SEQ ID NO:134)	LLQLSVGYMH (SEQ ID NO:504)
	(SEQ ID NO:98)	SLOLRVGYMH ((SEQ ID NO:140)	LLQIRVGYMH (SEQ ID NO:505)
	(SEQ ID NO:416)	STOTEVGYMH ((SEQ ID NO: 456)	LLQLEVGYMH (SEQ ID NO:506)
	(SEQ ID NO:92)	SLOSSUGYMH ((SEQ ID NO:136)	LLQSSVGYMH (SEQ ID NO:507)
1	(SEQ ID NO:95)	SLOSRVGYMH ((SEQ ID NO:138)	ID
	(SEQ ID NO:417)	SLOSFVGYMH ((SEQ ID NO: 457)	LLQSEVGYMH (SEQ ID NO:509)
1	(SEQ ID NO: 418)		(SEQ ID NO:458)	LLQVSVGYMH (SEQ ID NO:510)
	QI		(SEQ ID NO:459)	LLQVRVGYMH (SEQ ID NO:511)
	(SEQ ID NO: 420)	SLQVFVGYMH ((SEQ ID NO:460)	LLQVFVGYMH (SEQ ID NO:512)
	(SEQ ID NO:101)	SLSUGYMH ((SEQ ID NO:120)	LLSLSVGYMH (SEQ ID NO:513)
	(SEQ ID NO:110)	SLSLRVGYMH ((SEQ ID NO:125)	LLSLRVGYMH (SEQ ID NO:514)
KLSLFVGYMH ((SEQ ID NO:421)	SLSLFVGYMH ((SEQ ID NO:461)	LLSLEVGYMH (SEQ ID NO:515)
	ID	SLSSVGYMH ((SEQ ID NO:122)	LLSSSVGYMH (SEQ ID NO:516)
KLSSRVGYMH ((SEQ ID NO:107)	SLSSRVGYMH ((SEQ ID NO:22)	LLSSRVGYMH (SEQ ID NO:517)
KLSSFVGYMH ((SEQ ID NO: 422)	SISSEVGYMH ((SEQ ID NO:462)	LLSSEVGYMH (SEQ ID NO:518)
	QI	SLSVGYMH ((SEQ ID NO:463)	LLSVSVGYMH (SEQ ID NO:519)
KLSVRVGYMH ((SEQ ID NO: 424)	SLSVRVGYMH ((SEQ ID NO: 464)	LLSVRVGYMH (SEQ ID NO:520)
KLSVFVGYMH ((SEQ ID NO: 425)	SLSVFVGYMH ((SEQ ID NO: 465)	LLSVEVGYMH (SEQ ID NO:521)
	(SEQ ID NO:163)	SPOLSVGYMH ((SEQ ID NO:177)	LPQLSVGYMH (SEQ ID NO:200)
K P QLRVGYMH ((SEQ ID NO:159)	SPOLRVGYMH ((SEQ ID NO:173)	LPQLRVGYMH (SEQ ID NO:202)
	(SEQ ID NO: 426)	SPOLEVGYMH ((SEQ ID NO:466)	LPQLFVGYMH (SEQ ID NO:522)
	(SEQ ID NO:161)	SPOSSVGYMH ((SEQ ID NO:176)	LPQSSVGYMH (SEQ ID NO:201)
KPQSRVGYMH ((SEQ ID NO:157)	SPQSRVGYMH ((SEQ ID NO:171)	
l	(SEQ ID NO:427)	SPQSFVGYMH ((SEQ ID NO:467)	LPQSFVGYMH (SEQ ID NO:523)
1	(SEQ ID NO: 428)	SPQVSVGYMH ((SEQ ID NO:468)	LPQVSVGYMH (SEQ ID NO:524)
KPQVRVGYMH ((SEQ ID NO: 429)	SPQVRVGYMH ((SEQ ID NO: 469)	LPQVRVGYMH (SEQ ID NO:525)

KDOVFVGYMH	KPOWEVGYMH (SEO ID NO:430)	SPOVEVGYMH	SPOVEVGYMH (SEQ ID NO: 470)	TPQVFVGYMH ((SEQ ID NO: 526)
KDCT GVICYMH	KDELSVEYMH (SEO TO NO:155)	SPSLSVGYMH	SPSI.SVGYMH (SEO ID NO: 169)	LPSLSVGYMH ((SEQ ID NO:192)
KDST. BVGYWH	KDELEVICYMH (SEO TD NO: 152)	SPSLRVGYMH	SPSLRVGYMH (SEQ ID NO: 166)	LPSLRVGYMH ((SEQ ID NO:194)
KDST. FVGYMH	KDGT. TVGVMH (SEO TD NO: 431)	SPSLFVGYMH	SPSLEVGYMH (SEQ ID NO:471)	LPSLEVGYMH ((SEQ ID NO:527)
KDOCACH	KDOGSVGVMH (SEO TD NO:153)	SPSSSVGYMH	SPSSSVGYMH (SEO ID NO:168)	LPSSSVGYMH ((SEQ ID NO:193)
HMV2Wdgady	MESSYGILMI (SEC TO NO.150)	SPSSRVGYMH	SPSSRVGYMH (SEO ID NO:31)	LPSSRVGYMH ((SEQ ID NO:47)
VDCCEVCVMH	PECETICOMH (SEC TD NO.432)	SPSSFVGYMH	(SEO ID NO:472)	LPSSFVGYMH ((SEQ ID NO:528)
MADON SOLVEN	VESTS VGILLI (SEX ED NO: 132)	SPSVGYMH	(SEO ID NO:473)	LPSVSVGYMH ((SEQ ID NO:529)
KDCVDVCVMH	KDENDYCYMH (SEO ID NO: 434)	SPSVRVGYMH	(SEO ID NO:474)	LPSVRVGYMH ((SEQ ID NO:530)
K DSVF VGYMH	KDSVINGYMH (SEO TD NO: 435)	SPSVEVGYMH	SPSVFVGYMH (SEQ ID NO:475)	LPSVFVGYMH (LPSVEVGYMH (SEQ ID NO:531)
11111 O A 1 1 1 1	(C) 10 10 10 (C)				

Bold faced & underlined amino acid residues are the residues which differ from the amino acid sequence in palivizumab

Table 3E VI, CDR2 Sequences	2 Sequences					
DISKLAS (SEQ TD NO:5)	DTFKLAS (SEQ ID NO:15)	DTXKLAS (SEQ ID NO:799)	DT R KLAS (SEQ ID	DTMKLAS (SEQ ID	DTKKLAS (SEQ ID NO:1211)	DTLKLAS (SEQ ID NO:135)
			NOS:113&174)	NOS:121&162)		
DISKLSS (SEQ	DIFKLSS (SEQ	DIXKLSS (SEQ	DTRKLSS (SEQ	DIMKLSS (SEQ	DTKKLSS (SEQ	DILKIS (SEQ
ID NO:165)	ID_NO:96)	ID NO:800)	ID NO:175)	ID NO:164)	ID NO:1212)	1D NO: L355)
DISKL K S (SEQ	DIFKLKS (SEQ	DT <u>Y</u> KLKS (SEQ	DTRKLKS (SEQ	DIMKLKS (SEQ	DTKKLKS (SEQ	DTLKLKS (SEQ
ID NO:532)	ID_NO:660)	ID NO:801)	ID NO:943)	ID NO:1076)	ID NO:1213)	1D NO:1356)
DISKLRS (SEQ	DTFKLRS (SEQ	DIYKLRS (SEQ	DTRKLRS (SEQ	DIMKLRS (SEQ	DTKKLRS (SEQ	DTLKLRS (SEQ
ID NO:533)	ID_NO:661)	ID_NO:802)	ID NO:944)	ID NO:1077)	ID NO:1214)	ID NO:1357)
DISKL H S (SEO	DTFKLHS (SEQ	DTYKLHS (SEQ	DTRKLHS (SEQ	DIMKLHS (SEQ	DIKKLHS (SEQ	DILKLHS (SEQ
ID NO:534)	ID_NO:662)	ID_NO:803)	ID_NO:945)	ID NO:1078)	ID NO:1215)	ID NO:1358)
DTSKLPS (SEQ	DIEKLPS (SEQ	DIXKLPS (SEQ	DTRKLPS (SEQ	DIMKIPS (SEG	DTKKLPS (SEQ	DTLKLPS (SEQ
ID NO:102)	ID NO:663)	ID NO:804)	ID NO:118)	ID NO:1079)	LD NO: LZI6)	TD NO: T30%)

			T			Ţ <u> </u>				
DTLKLTS (SEQ	DTLKLDS (SEQ	DT <u>raas</u> (SEQ	DTLKHSS (SEQ	DTLKHKS (SEQ	DTLKHRS (SEQ	DTLKHHS (SEQ	DTLKHPS (SEQ	DTLKHTS (SEQ	DT <u>r</u> k hd s (SEQ	DT <u>r</u> K o as (Seo
ID NO:1360)	ID NO:131)	ID NO:1361)	ID NO:1362)	ID NO:1363)	ID NO:1364)	ID NO:1365)	ID NO:1366)	ID NO:1367)	ID NO:1368)	ID NO:1369)
DTKKLTS (SEQ	DTKKLDS (SEQ	DTKKHAS (SEQ	DTKKHS (SEQ	DT K K HK S (SEQ	DTKKHRS (SEQ	DTKKHHS (SEQ	DTKKHPS (SEQ	DTKKHTS (SEQ	DTKKHDS (SEQ	DT <u>k</u> K <u>o</u> as (seo
ID NO:1217)	ID NO:1218)	ID NO:1219)	ID NO:1220)	ID NO:1221)	ID NO:1222)	ID NO:1223)	ID NO:1224)	ID NO:1225)	ID NO:1226)	ID NO:1227)
DIMKLIS (SEQ	DTMKLDS (SEQ	DTWKHAS (SEQ	DTWKHSS (SEQ	DTMKHKS (SEQ	DTMKHRS (SEQ	DT MKHH S (SEQ	DIMKHPS (SEQ	DTMKHTS (SEQ	DTMKHDS (SEQ	DT <u>w</u> K <u>o</u> AS (SEQ
ID NO:1080)	ID NO:1081)	ID NO:1082)	ID NO:1083)	ID NO:1084)	ID NO:1085)	ID NO:1086)	ID NO:1087)	ID NO:1088)	ID NO:1089)	ID NO:154)
DIRKLIS (SEQ	DTRKLDS (SEQ	DIRKHAS (SEQ	DTRKHSS (SEQ	DTRKHKS (SEQ	DTRKHRS (SEQ	DT R K HH S (SEQ	DTRKHPS (SEQ	DTRKHTS (SEQ	DT <u>r</u> K hd s (SEQ	DTRKQAS (SEQ
ID NO:946)	ID NO:947)	ID NO:948)	ID NO:949)	ID NO:950)	ID NO:951)	ID NO:952)	ID NO:953)	ID NO:954)	ID NO:955)	ID NO:170)
DT <u>Y</u> KL <u>T</u> S (SEQ	DT <u>Y</u> KL D S (SEQ	DTYKHAS (SEQ	DTYKHES (SEQ	DT <u>v</u> k <u>hk</u> s (seq	DT <u>v</u> khrs (seq	DT <u>Y</u> K HH S (SEQ	DT <u>Y</u> K HP S (SEQ	DT <u>x</u> K ht S (SEQ	DT <u>x</u> k hd s (Seq	DT <u>ykoa</u> s (SEQ
ID NO:805)	ID NO:806)	ID NO:807)	ID NO:808)	ID NO:809)	ID NO:810)	ID NO:811)	ID NO:812)	ID NO:813)	Id No:814)	ID NO:815)
DTEKLTS (SEQ	DT E KL D S (SEQ	DTEKHAS (SEQ	DTEKHSS (SEQ	DT e k hk s (Seq	DTEKHRS (SEQ	DTEKHHS (SEQ	DT <u>FKHP</u> S (SEQ	DT <u>FKHTS</u> (SEQ	DT <u>FKHD</u> S (SEQ	DTEKQAS (SEQ
ID NO:664)	ID NO:665)	ID NO:666)	ID NO:667)	ID NO:668)	ID NO: 669)	ID NO: 670)	ID NO: 671)	ID NO: 672)	ID NO:673)	ID NO:674)
DTSKL <u>t</u> s (SEQ	DTSKL D S (SEQ	DTSK H AS (SEQ	DTSKHSS (SEQ	DISKHKS (SEQ	DISK hr S (SEQ	DTSK hh S (SEQ	DTSK HP S (SEQ	DISK HI S (SEQ	DTSKHDS (SEQ	DTSK Q AS (SEQ
ID NO:535)	ID NO:128)	ID NO:536)	ID NO:537)	ID NO:538)	ID NO:539)	ID NO:540)	ID NO:541)	ID NO:542)	ID NO:543)	ID NO:139)

DT <u>rkos</u> s (SEQ	DT <u>rkor</u> s (SEQ	DT <u>r</u> K <u>Qr</u> S (SEQ	DT L K QH S (SEQ	DT <u>r</u> K <u>QP</u> S (SEQ	DT <u>rkot</u> s (SEQ	DT <u>rkod</u> s (SEQ	DTLGLAS (SEQ	DTIGES (SEQ	DTLGLKS (SEQ	DTLGLRS (SEQ
ID NO:1370)	ID NO:1371)	ID NO:1372)	ID NO:1373)	ID NO:1374)	ID NO:1375)	ID NO:1376)	ID NO:1377)	ID NO:1378)	ID NO:1379)	ID NO:1380)
DTKKQSS (SEQ	DTKKQKS (SEQ	DIKKQRS (SEQ	DTKK QH S (SEQ	DTKK QP S (SEQ ID NO:1232)	DTKK QT S (SEQ	DTKKQDS (SEQ	DTKGLAS (SEQ	DTKGLSS (SEQ	DTKGLKS (SEQ	DTKGLRS (SEQ
ID NO:1228)	ID NO:1229)	ID NO:1230)	ID NO:1231)		ID NO:1233)	ID NO:1234)	ID NO:1235)	ID NO:1236)	ID NO:1237)	ID NO:1238)
DIMK QS S (SEQ	DTMKQKS (SEQ	DTMKQRS (SEQ	DTMKQHS (SEQ	DTMKQPS (SEQ	DTMKQTS (SEQ	DTMKQDS (SEQ	DTMGLAS (SEQ	DTMGLSS (SEQ	DIMGLKS (SEQ	DTMGLES (SEQ
ID NO:156)	ID NO:1090)	ID NO:1091)	ID NO:1092)	ID NO:1093)	ID NO:1094)	ID NO:1095)	ID NO:1096)	ID NO:1097)	ID NO:1098)	ID NO:1099)
DTRKQSS(SEQ	DTRKQKS (SEQ	DTRKQRS (SEQ	DTRKQHS (SEQ	DTRK <u>QP</u> S (SEQ	DTRKQTS (SEQ	DTRKQDS (SEQ	DTRGLAS (SEQ	DTRGLSS (SEQ	DTRGLKS (SEQ	DTRGLRS (SEQ
ID NO:172)	ID NO:956)	ID NO:957)	ID NO:958)	ID NO:959)	ID NO:960)	ID NO:961)	ID NO:116)	ID NO:962)	ID NO:963)	ID NO:964)
DTXKQSS (SEQ	DT <u>Y</u> K <u>QK</u> S (SEQ	DTYKQRS (SEQ	DTXKQHS (SEQ	DTYKQPS (SEQ	DT <u>x</u> K <u>QT</u> S (SEQ	DT <u>y</u> K <u>od</u> s (SEQ	DT <u>¥G</u> LAS (SEQ	DTYGLSS (SEQ	DTYGLKS (SEQ	DTYGLRS (SEQ
ID NO:816)	ID NO:817)	ID NO:818)	ID NO:819)	ID NO:820)	ID NO:821)	ID NO:822)	ID NO:823)	ID NO:824)	ID NO:825)	ID NO:826)
DTEKQSS (SEQ	DTFKQKS (SEQ	DTFKQRS (SEQ	DT <u>e</u> K <u>Qa</u> S (SEQ	DTEKQES (SEQ	DTEKQTS (SEQ	DTEKQDS (SEQ	DTFGLAS (SEQ	DT FG LSS (SEQ	DTFGLKS (SEQ	DTFGLRS (SEQ
ID NO:675)	ID NO:676)	ID NO:677)	ID NO:678)	ID NO:679)	ID NO:680)	ID NO:681)	ID NO:682)	ID NO: 683)	ID NO:684)	ID NO: 685)
DTSK QS S (SEQ	DTSK <u>ok</u> s (SEQ	DTSK <u>OR</u> S (SEQ	DTSK QH S (SEQ	DTSK QP S (SEQ	DTSK <u>QT</u> S (SEQ	DTSK OD S (SEQ	DTSGLAS (SEQ	DTSGLSS (SEQ	DTSGLKS (SEQ	DTSGLRS (SEQ
ID NO:141)	ID NO:544)	ID NO:545)	ID NO:546)	ID NO:547)	ID NO:548)	ID NO:549)	ID NO:105)	ID NO:550)	ID NO:551)	ID NO:552)

DTSGLHS (SEQ	DTFGLHS (SEQ	DT <u>YG</u> LHS (SEQ	DTRGLHS (SEQ	DTMCLHS (SEQ	DT <u>kg</u> l h s (SEQ	DTLGLES (SEQ
ID NO:553)	ID NO:686)	ID NO:827)	ID NO:965)	ID NO:1100)	ID NO:1239)	ID NO:1381)
DTSGLPS (SEQ	DTEGLES (SEQ	DTYGLPS (SEQ	DTRGLPS(SEQ	DTMGLPS (SEQ ID NO:1101)	DTKGLPS (SEQ	DT <u>LG</u> LPS (SEQ
ID NO:108)	ID NO: 687)	ID NO:828)	ID NO:27)		ID NO:1240)	ID NO:1382)
DTSGLTS (SEQ	DIFCLTS (SEQ	DT <u>YG</u> LTS (SEQ	DTRGLTS (SEQ	DTMGLTS (SEQ	DTKGLTS (SEQ	DTLGLTS (SEQ
ID NO:554)	ID NO: 688)	ID NO:829)	ID NO:966)	ID NO:1102)	ID NO:1241)	ID NO:1383)
DTSGLDS (SEQ	DTFGLDS (SEQ	DT <u>YG</u> LDS (SEQ	DIRGLDS (SEQ	DTMGLDS (SEQ	DTKGLDS (SEQ	DTLGLDS (SEQ
ID NO:555)	ID NO:689)	ID NO:830)	ID NO:967)	ID NO:1103)	ID NO:1242)	ID NO:1384)
DTS GH AS (SEQ	DT FGH AS (SEQ	DT YGH AS (SEQ	DTRGHAS (SEQ	DTMGHAS (SEQ	DTKGHAS (SEQ	DTLGHAS (SEQ
ID NO:556)	ID NO: 690)	ID NO:831)	ID NO:968)	ID NO:1104)	ID NO:1243)	ID NO:1385)
DTSGHSS (SEQ	DIFGHSS (SEQ	DTYGHSS (SEQ	DIRGHS (SEQ	DTMGHS (SEQ	DTKGHSS (SEQ	DTLGHSS (SEQ
ID NO:557)	ID NO:691)	ID NO:832)	ID NO:969)	ID NO:1105)	ID NO:1244)	ID NO:1386)
DTS GHK S (SEQ	DTFGHKS (SEQ	DTYGHKS (SEQ	DTRGHKS (SEQ	DTMGHKS (SEQ	DT KGHK S (SEQ	DTLGHKS (SEQ
ID NO:558)	ID NO:692)	ID NO:833)	ID NO:970)	ID NO:1106)	ID NO:1245)	ID NO:1387)
DTSGHRS (SEQ	DT FGHR S (SEQ	DT <u>YGHR</u> S (SEQ	DTRGHRS (SEQ	DTMGHRS (SEQ	DT KGHR S (SEQ	DT <u>LGHR</u> S (SEQ
ID NO:559)	ID NO: 693)	ID NO:834)	ID NO:971)	ID NO:1107)	ID NO:1246)	ID NO:1388)
DTSGHHS (SEQ	DT FGHH S (SEQ	DTYGHHS (SEQ	DTRGHHS (SEQ	DTMGHHS (SEQ	DT KGHH S (SEQ	DT <u>LGHH</u> S (SEQ
ID NO:560)	ID NO: 694)	ID NO:835)	ID NO:972)	ID NO:1108)	ID NO:1247)	ID NO:1389)
DTSGHPS (SEQ	DTEGHPS (SEQ	DT <u>YGHP</u> S (SEQ	DTRGHPS (SEQ	DTMGHPS (SEQ	DTKGHPS (SEQ	DTLGHPS (SEQ
ID NO:561)	ID NO: 695)	ID NO:836)	ID NO:973)	ID NO:1109)	ID NO:1248)	ID NO:1390)
DTSGHTS (SEQ	DTFGHTS (SEQ	DTYGHTS (SEQ	DTRGHTS (SEQ	DTMGHTS (SEQ	DTKGHTS (SEQ	DTLGHTS (SEQ
ID NO:562)	ID NO:696)	ID NO:837)	ID NO:974)	ID NO:1110)	ID NO:1249)	ID NO:1391)

DT <u>LGHD</u> S (SEQ	DT <u>reo</u> AS (SEQ	DT <u>LGQS</u> S (SEQ	DT LGQK S (SEQ	DT LGQR S (SEQ	DTLGQHS (SEQ	DT <u>LGQPS</u> (SEQ	DT <u>LGQT</u> S (SEQ	DTIGODS (SEQ	DTLRLAS (SEQ	DTLRLS (SEQ
ID NO:1392)	ID NO:1393)	ID NO:1394)	ID NO:1395)	ID NO:1396)	ID NO:1397)	ID NO:1398)	ID NO:1399)	ID NO:1400)	ID NO:1401)	ID NO:1402)
DT KGHD S (SEQ	DT KGQ AS (SEQ	DT KGQS S (SEQ	DT kgok s (SEQ	DT KGQR S (SEQ	DT KGQH S (SEQ	DT KGQP S (SEQ	DT KGQT S (SEQ	DT KGQD S (SEQ	DTKRLAS (SEQ	DTKRLS (SEQ
ID NO:1250)	ID NO:1251)	ID NO:1252)	ID NO:1253)	ID NO:1254)	ID NO:1255)	ID NO:1256)	ID NO:1257)	ID NO:1258)	ID NO:1259)	ID NO:1260)
DIMGHDS (SEQ	DTMGQAS (SEQ	DTMGQS (SEQ	DIMGORS (SEQ	DIMGORS (SEQ	DIMGQHS (SEQ	DTMGQPS (SEQ	DTMGQTS (SEQ	DTMGQDS (SEQ	DTMRLAS (SEQ	DTMRLSS (SEQ
ID NO:1111)	ID NO:1112)	ID NO:1113)	ID NO:1114)	ID NO:1115)	ID NO:1116)	ID NO:1117)	ID NO:1118)	ID NO:1119)	ID NO:32)	ID NO:1120)
DTRGHDS (SEQ	DT RGQ AS (SEQ	DTRGOSS (SEQ	DT RGQK S (SEQ	DT RGQR S (SEQ	DT RGQH S (SEQ	DT RGQP S (SEQ	DTRGQTS (SEQ	DTRGODS (SEQ	DTRLAS (SEQ	DTRRLSS (SEQ
ID NO:975)	ID NO:976)	ID NO:977)	ID NO:978)	ID NO:979)	ID NO:980)	ID NO:981)	ID NO:982)	ID NO:983)	ID NO:984)	ID NO:985)
DT <u>YGHD</u> S (SEQ	DT YGQ AS (SEQ	DT <u>YGOS</u> S (SEQ	DT YGQK S (SEQ	DT YGQR S (SEQ	DT YGQH S (SEQ	DT <u>YGQP</u> S (SEQ	DT YGQT S (SEQ	DT <u>YGQD</u> S (SEQ	DTYRLAS (SEQ	DT <u>YR</u> LSS (SEQ
ID NO:838)	ID NO:839)	ID NO:840)	ID NO:841)	ID NO:842)	ID NO:843)	ID NO:844)	ID NO:845)	ID NO:846)	ID NO:847)	ID NO:848)
DTEGHDS (SEQ	DT FGQ AS (SEQ	DT FGQS S (SEQ	DTFGQKS (SEQ	DTFGQRS (SEQ	DT FGQH S (SEQ.	DT EGQP S (SEQ	DT EGQT S (SEQ	DTEGODS (SEQ	DTERLAS (SEQ	DTERLSS (SEQ
ID NO:697)	ID NO:698)	ID NO:699)	ID NO:700)	ID NO:701)	ID NO:702)	ID NO:703)	ID NO:704)	ID NO:705)	ID NO:706)	ID NO:707)
DTS GHD S (SEQ	DTS GQA S (SEQ	DTS GQS S (SEQ	DTS GQK S (SEQ	DTSGORS (SEQ	DIS GQH S (SEQ	DIS GQP S (SEQ	DTSG <u>QT</u> S (SEQ	DISGODS (SEQ	DTSRLAS (SEQ	DTSKLSS (SEQ
ID NO:563)	ID NO:564)	ID NO:565)	ID NO:566)	ID NO:567)	ID NO:568)	ID NO:569)	ID NO:570)	ID NO:571)	ID NO:123)	ID NO:572)

DTKRLKS (SEQ DTLRLKS (SEQ ID NO:1261) ID NO:1403)	DTKRIRS (SEQ DTLRIRS (SEQ ID NO:1262) ID NO:1264)	DTKRLHS (SEQ DTLRLHS (SEQ ID NO:1263) ID NO:1405)	DTKRIPS (SEQ DTLRIPS (SEQ ID NO:1264) ID NO:1406)	DTKRLTS (SEQ DTLRLTS (SEQ ID NO:1265) ID NO:1407)	DTKRIDS (SEQ DTLRIDS (SEQ ID NO:1266) ID NO:1408)	DIKRHAS (SEQ DILRHAS (SEQ ID NO:1267) ID NO:1409)	DIKRHS (SEQ DILRHS (SEQ ID NO:1268) ID NO:1410)	DIKRHKS (SEQ DILRHKS (SEQ ID NO:1269) ID NO:1411)	DIKRHRS (SEQ DILRHRS (SEQ ID NO:1270) ID NO:1412)	ORS (SEO OFFICE SEO
DTMRLKS (SEQ DTK ID NO:1121) ID	DTMRLRS (SEQ DTW ID NO:1122) ID	DTWRLHS (SEQ DTK ID NO:1123) ID	DTWRLPS (SEQ DTW ID NO:1124) ID	DTMRLTS (SEQ DTM ID NO:1125) ID	DTMRLDS (SEQ DTK ID NO:1126) ID	DTMRHAS (SEQ DTK ID NO:1127) ID	DIMRHSS (SEQ DIWIND ID ID	DTMRHKS (SEQ DTK ID NO:1129) ID	DIMRHRS (SEQ DIK ID NO:1130) ID	DTMRHHS (SEO DTK
DTRELKS (SEQ	DTRELES (SEQ	DTRELES (SEQ	DTRELES (SEQ	DTRELTS (SEQ	DTRRLDS (SEQ	DTRRHAS (SEQ	DTRRHSS (SEQ	DTRRHKS (SEQ	DTRRHRS (SEQ	DES (SEO
ID NO:986)	ID NO:987)	ID NO:988)	ID NO:989)	ID NO:990)	ID NO:991)	ID NO:992)	ID NO:993)	ID NO:994)	ID NO:995)	
DT <u>YR</u> LKS (SEQ	DTYRLRS (SEQ	DTYRLES (SEQ	DTXRLPS (SEQ	DTXRLTS (SEQ	DTYRLDS (SEQ	DT <u>YRH</u> AS (SEQ	DTYRHSS (SEQ	DTYRHKS (SEQ	DT <u>YRHR</u> S (SEQ	CHS/ SHHGVIFT
ID NO:849)	ID NO:850)	ID NO:851)	ID NO:852)	ID NO:853)	ID NO:854)	ID NO:855)	ID NO:856)	ID NO:857)	ID NO:858)	
DTFRLKS (SEQ	DTFRLRS (SEQ	DTERLHS (SEQ	DTERLES (SEQ	DTERLES (SEQ	DTERLDS (SEQ	DTERHAS (SEQ	DTFRHSS (SEQ	DTFRHKS (SEQ	DTFRHRS (SEQ	OTO / OTO
ID NO:708)	ID NO:709)	ID NO:710)	ID NO:711)	ID NO:712)	ID NO:713)	ID NO:714)	ID NO:715)	ID NO:716)	ID NO:717)	
DTSRLKS (SEQ	DTSRLRS (SEQ	DTSRLHS (SEQ	DTSRLPS (SEQ	DTSRLTS (SEQ	DTSRLDS (SEQ	DTSRHAS (SEQ	DTSRHSS (SEQ	DTSRHKS (SEQ	DIS RHR S (SEQ	1000 / CEO
ID NO:573)	ID NO:574)	ID NO:575)	ID NO:576)	ID NO:577)	ID NO:578)	ID NO:579)	ID NO:580)	ID NO:581)	ID NO:582)	

DTYRHPS (SEQ DTRRHPS (S ID NO:860) ID NO:997)
DTYRHTS (SEQ ID NO:861)
DTXRHDS (SEQ ID NO:862)
DT <u>yrq</u> as (SEQ ID NO:863)
DT <u>YRQS</u> S (SEQ ID NO:864)
DT YRQK S (SEQ ID NO:865)
DTYRQRS (SEQ ID NO:866)
DT yroh s (SEQ ID NO:867)
DTYRQPS (SEQ ID NO:868)
DTYRQTS (SEQ ID NO:869)
DT <u>xrod</u> s (SEQ ID NO:870)

(SEQ DT <u>LY</u> LAS (SEQ 10 NO:1425)	(SEQ DILYLS (SEQ 84) ID NO:1426)	SEQ 5)	(SEQ DILYLRS 86) ID NO:14	(SEQ DTLYLES (SEQ .287) ID NO:1429)	(SEQ DTLYLES 88) ID NO:14	(SEQ DTLYLTS (SEQ 89) ID NO:1431)	(SEQ DTLYIDS (SEQ 90) ID NO:1432)	(SEQ DTLYHAS (SEQ 91) ID NO:1433)	(SEQ DTLYHSS (SEQ 92) ID NO:1434)	(SEQ DTLYHKS (SEQ 93) ID NO:1435)
DTKYLAS (SE	DTKYLSS (SE	0	DTKKIRS	DTKYLHS	DTKYLPS	DTKYLTS (DTKYLDS	DTWHAS (SE	DTKYHSS (SE	DTKYHKS (SE
ID NO:1283)	ID NO:1284)		ID NO:12	ID NO:12	ID NO:12	ID NO:128	ID NO:12	ID NO:1291)	ID NO:1292)	ID NO:1293)
DTMYLAS (SEQ	DTMYLSS (SEQ	DTMYLKS (SEQ	DTMYLRS (SEQ	DT <u>MY</u> LHS (SEQ	DTMYLPS (SEQ	DIMKLTS (SEQ	DTMYLDS (SEQ	DTWTHAS (SEQ	DTMYHSS (SEQ	DTWYHKS (SEQ
ID NO:158)	ID NO:160)	ID NO:1143)	ID NO:1144)	ID NO:1145)	ID NO:1146)	ID NO:1147)	ID NO:1148)	ID NO:1149)	ID NO:1150)	ID NO:1151)
DTRYLAS (SEQ	DTRYLSS (SEQ	DTRYLKS (SEQ	DTRYLRS (SEQ	DTRYLHS (SEQ	DTRYLPS (SEQ	DTRYLTS (SEQ	DTRYLDS (SEQ	DTRYHAS (SEQ	DTRYHSS (SEQ	DTRYHKS (SEQ
ID NO:178)	ID NO:59)	ID NO:1008)	ID NO:1009)	ID NO:1010)	ID NO:1011)	ID NO:1012)	ID NO:1013)	ID NO:1014)	ID NO:1015)	ID NO:1016)
DT <u>YY</u> LAS (SEQ	DT <u>YY</u> LSS (SEQ	DTYYLKS (SEQ	DT <u>YY</u> LRS (SEQ	DT <u>YY</u> LHS (SEQ	DT <u>YY</u> LPS (SEQ	DT <u>YY</u> L <u>T</u> S (SEQ	DT <u>YY</u> LDS (SEQ	DT <u>YYH</u> AS (SEQ	DT <u>YYHS</u> S (SEQ	DT <u>YYHK</u> S (SEQ
ID NO:871)	ID NO:872)	ID NO:873)	ID NO:874)	ID NO:875)	ID NO:876)	ID NO:877)	ID NO:878)	ID NO:879)	ID NO:880)	ID NO:881)
DI FY LAS (SEQ	DTEYLSS (SEQ	DTFYLKS (SEQ	DIFYLRS (SEQ	DI fy l h s (Seq	DTEYLES (SEQ	DTEYLES (SEQ	DTEYLDS (SEQ	DTEYHAS (SEQ	DTEYHSS (SEQ	DTEYHKS (SEQ
ID NO:99)	ID NO:90)	ID NO:730)	ID NO:731)	ID NO:732)	ID NO:733)	ID NO:734)	ID NO:735)	ID NO:736)	ID NO:737)	ID NO:738)
DTS <u>Y</u> LAS (SEQ ID NOS-81&143)	(S)	DIS <u>Y</u> IKS (SEQ ID NO:595)	DTS <u>Y</u> LRS (SEQ ID NO:596)	DTS <u>Y</u> LHS (SEQ ID NO:597)	DTS <u>Y</u> L <u>P</u> S (SEQ ID NO:598)	DTS <u>Y</u> LTS (SEQ ID NO:599)	DTS <u>Y</u> L <u>D</u> S (SEQ ID NO:600)	DTS <u>vh</u> AS (SEQ ID NO:601)	DTS <u>YHS</u> S (SEQ ID NO:602)	DTS <u>vhk</u> s (SEQ ID NO:603)

	DTYYHRS (SEQ DTRYHRS (SEQ DTMYHRS ID NO:1017) ID NO:111
DTRYHES ID NO: 1(ID NO: 5(ID	(SEQ DTYYHRS (SEQ DTRYHRS (SEQ DTYYHRS (SEQ DTRYHRS 0) ID NO:883) ID NO:10 (SEQ DTYYHPS (SEQ DTRYHPS 1) ID NO:884) ID NO:10 (SEQ DTYYHTS (SEQ DTRYHTS 2) ID NO:885) ID NO:10 (SEQ DTYYHDS (SEQ DTRYHDS 3) ID NO:885) ID NO:10 (SEQ DTYYQS (SEQ DTRYDS 4) ID NO:887) ID NO:10 (SEQ DTYYQS (SEQ DTRYQS 5) ID NO:887) ID NO:16 (SEQ DTYYQS (SEQ DTRYQS 5) ID NO:888) ID NO:55
DTYYHRS (SEQ ID NO:882) DTYHHS (SEQ ID NO:883) DTYHHS (SEQ ID NO:884) DTYHTS (SEQ ID NO:885) DTYYHDS (SEQ ID NO:885) DTYYORS (SEQ ID NO:887) DTYYORS (SEQ ID NO:888)	(SEQ DTYYHRS (SEQ DTYYHRS (SEQ DTYYHRS 1) ID NO:88 (SEQ DTYYHTS 2) ID NO:88 (SEQ DTYYHTS 3) ID NO:88 (SEQ DTYYHDS 3) ID NO:88 (SEQ DTYYDAS 4) ID NO:88 5) ID NO:88 650 DTYYOSS 5) ID NO:88
	(SEQ (SEQ (SEQ (SEQ (SEQ (SEQ (SEQ (SEQ

DTS YQT S (SEQ	DTEYQTS (SEQ	DT <u>YYQT</u> S (SEQ	DTRYQTS (SEQ	DTMYQTS (SEQ	DT <u>KYQT</u> S (SEQ	DTLYQTS (SEQ
ID NO:613)	ID NO:750)	ID NO:893)	ID NO:1026)	ID NO:1161)	ID NO:1305)	ID NO:1447)
DTS <u>YQD</u> S (SEQ	DTFYQDS (SEQ	DT <u>YYQD</u> S (SEQ	DT RYQD S (SEQ	DTMYQDS (SEQ	DT KYQD S (SEQ	DTLYQDS (SEQ
ID NO:614)	ID NO:751)	ID NO:894)	ID NO:1027)	ID NO:1162)	ID NO:1306)	ID NO:1448)
DTS <u>F</u> LAS (SEQ	DTFELAS (SEQ	DT <u>YF</u> LAS (SEQ	DTRELAS (SEQ	DTMFLAS (SEQ	DTKFLAS (SEQ	DTLFLAS (SEQ
ID NO:615)	ID NO:752)	ID NO:895)	ID NO:1028)	ID NO:1163)	ID NO:1307)	ID NO:1449)
DTSFLSS (SEQ	DTFFLSS (SEQ	DTX#LSS (SEQ	DTRFLSS (SEQ	DTMFLSS (SEQ	DTKFLSS (SEQ	DTLFLSS (SEQ
ID NO:616)	ID NO:753)	ID NO:896)	ID NO:1029)	ID NO:1164)	ID NO:1308)	ID NO:1450)
DTSFLKS (SEQ	DTETKS (SEQ	DT <u>YF</u> LKS (SEQ	DTRELKS (SEQ	DTMFLKS (SEQ	DTKFLKS (SEQ	DTLFLKS (SEQ
ID NO:617)	ID NO:754)	ID NO:897)	ID NO:1030)	ID NO:1165)	ID NO:1309)	ID NO:1451)
DTSELES (SEQ	DTFFIRS (SEQ	DTYFLRS (SEQ	DTRELES (SEQ	DTMFLRS (SEQ	DT KF LRS (SEQ	DT <u>lf</u> lrs (SEQ
ID NO:618)	ID NO:755)	ID NO:898)	ID NO:1031)	ID NO:1166)	ID NO:1310)	ID NO:1452)
DTSFLHS (SEQ	DTETHS (SEQ	DIXFLES (SEQ	DTRELHS (SEQ	DIMELHS (SEQ	DTKFLHS (SEQ	DTLFLHS (SEQ
ID NO:619)	ID NO:756)	ID NO:899)	ID NO:1032)	ID NO:1167)	ID NO:1311)	ID NO:1453)
DTSFLPS (SEQ	DTFIPS (SEQ	DTXFLPS (SEQ	DTRFLPS (SEQ	DTMFLPS (SEQ	DTKFLPS (SEQ	DTLFLPS (SEQ
ID NO:620)	ID NO:757)	ID NO:900)	ID NO:1033)	ID NO:1168)	ID NO:1312)	ID NO:1454)
DTSFLTS (SEQ	DTFELTS (SEQ	DTXFLTS (SEQ	DTRFLTS (SEQ ID NO:1034)	DTMFLTS (SEQ	DTKFLTS (SEQ	DT <u>LF</u> LTS (SEQ
ID NO:621)	ID NO:758)	ID NO:901)		ID NO:1169)	ID NO:1313)	ID NO:1455)
DTS <u>F</u> LDS (SEQ	DTFFLDS (SEQ	DTXFLDS (SEQ	DTRELDS (SEQ	DTMFLDS (SEQ	DTKFLDS (SEQ	DT <u>LF</u> LDS (SEQ
ID NO:77)	ID NO:50)	ID NO:902)	ID NO:1035)	ID NO:1170)	ID NO:1314)	ID NO:1456)
DTS FH AS (SEQ	DTFFHAS (SEQ	DI <u>YFH</u> AS (SEQ	DTRFHAS (SEQ	DTMEHAS (SEQ	DTKEHAS (SEQ	DT <u>LFH</u> AS (SEQ
ID NO:622)	ID NO:759)	ID NO:903)	ID NO:1036)	ID NO:1171)	ID NO:1315)	ID NO:1457)

DTSFHSS (SEQ	DTEFHSS (SEQ	DT <u>YFHS</u> S (SEQ	DTRFHSS (SEQ	DTMFHSS (SEQ	DT kfhs s (SEQ	DT <u>LFHS</u> S (SEQ
ID NO:623)	ID NO:760)	ID NO:904)	ID NO:1037)	ID NO:1172)	ID NO:1316)	ID NO:1458)
DTS FHK S (SEQ	DTFFHKS (SEQ	DTYFHKS (SEQ	DTRFHKS (SEQ	DIMEHKS (SEQ	DTKFHKS (SEQ	DTLFHKS (SEQ
ID NO:624)	ID NO:761)	ID NO:905)	ID NO:1038)	ID NO:1173)	ID NO:1317)	ID NO:1459)
DTS FHR S (SEQ	DT efhr S (SEQ	DT <u>YFHR</u> S (SEQ	DTRFHRS (SEQ	DIMEHRS (SEQ	DTKFHRS (SEQ	DTLEHRS (SEQ
ID NO:625)	ID NO:762)	ID NO:906)	ID NO:1039)	ID NO:1174)	ID NO:1318)	ID NO:1460)
DTS FHH S (SEQ	DTEFHHS (SEQ	DTYFHHS (SEQ	DIRFHHS (SEQ	DTMFHHS (SEQ	DTKFHHS (SEQ	DTLEHHS (SEQ
ID NO:626)	ID NO:763)	ID NO:907)	ID NO:1040)	ID NO:1175)	ID NO:1319)	ID NO:1461)
DTS FHP S (SEQ	DTFFHPS (SEQ	DT YFHP S (SEQ	DTREHES (SEQ	DIMEHPS (SEQ	DT KFHP S (SEQ	DTLFHPS (SEQ
ID NO:627)	ID NO:764)	ID NO:908)	ID NO:1041)	ID NO:1176)	ID NO:1320)	ID NO:1462)
DTS FHT S (SEQ	DIFFHTS (SEQ	DT YFHT S (SEQ	DTREHTS (SEQ	DIMEHIS (SEQ	DT KFHT S (SEQ	DTLEHTS (SEQ
ID NO:628)	ID NO:765)	ID NO:909)	ID NO:1042)	ID NO:1177)	ID NO:1321)	ID NO:1463)
DTS FHD S (SEQ	DT FFHD S (SEQ	DTXFHDS (SEQ	DTRFHDS (SEQ	DTWFHDS (SEQ	DT KFHD S (SEQ	DTLEHDS (SEQ
ID NO:629)	ID NO:766)	ID NO:910)	ID NO:1043)	ID NO:1178)	ID NO:1322)	ID NO:1464)
DTS FQ AS (SEQ	DT FFQ AS (SEQ	DT <u>YFQ</u> AS (SEQ	DTRFQAS (SEQ	DTMFQAS (SEQ	DT KFQ AS (SEQ	DTLFQAS (SEQ
ID NO:630)	ID NO:767)	ID NO:911)	ID NO:1044)	ID NO:1179)	ID NO:1323)	ID NO:1465)
DTS FQS S (SEQ	DIFFQSS (SEQ	DT <u>YFQS</u> S (SEQ	DTRFQS (SEQ	DTMFQS (SEQ	DTKFQSS (SEQ	DTLEQS (SEQ
ID NO:631)	ID NO:768)	ID NO:912)	ID NO:1045)	ID NO:1180)	ID NO:1324)	ID NO:1466)
DTS FOK S (SEQ	DT FFQK S (SEQ	DT <u>yfok</u> s (SEQ	DIREOKS (SEQ	DTMFQKS (SEQ	DTKFQKS (SEQ	DT LFQK S (SEQ
ID NO:632)	ID NO:769)	ID NO:913)	ID NO:1046)	ID NO:1181)	ID NO:1325)	ID NO:1467)
DTS FQR S (SEQ	DT FFQR S (SEQ	DT <u>YFQR</u> S (SEQ	DTREORS (SEQ	DIMEQRS (SEQ	DT KFQR S (SEQ	DT LFOR S (SEQ
ID NO:633)	ID NO:770)	ID NO:914)	ID NO:1047)	ID NO:1182)	ID NO:1326)	ID NO:1468)

DT leoh s (SEQ	DI leop s (SEQ	DT leot s (SEQ	DTLEQDS (SEQ	DT <u>LL</u> LAS (SEQ	DTILLSS (SEQ ID NO:1473)	DT <u>re</u> lks (SEQ	DT <u>LL</u> LRS (SEQ	DT <u>LL</u> LHS (SEQ	DT <u>LL</u> LES (SEQ	DTLLTS (SEQ
ID NO:1469)	ID NO:1470)	ID NO:1471)	ID NO:1472)	ID NO:133)		ID NO:1474)	ID NO:1475)	ID NO:1476)	ID NO:1477)	ID NO:1478)
DT KFQH S (SEQ D'	DT KFQP S (SEQ DT	DT KFQT S (SEQ D'	DT KFQD S (SEQ D'	DTKLLAS (SEQ D	DTKLLSS (SEQ D	DTKLLKS (SEQ D	DTKLLRS (SEQ D	DTKLLES (SEQ DID NO:1335) I	DIKLLPS (SEQ D	DTKLLTS (SEQ D
ID NO:1327) I	ID NO:1328) ID	ID NO:1329) I	ID NO:1330) I	ID NO:1331) I	ID NO:1332) I	ID NO:1333) I	ID NO:1334) I		ID NO:1336) I	ID NO:1337) I
DIMEQHS (SEQ	DTMFQPS (SEQ	DTMFQTS (SEQ	DTMEQDS (SEQ	DTMLLAS (SEQ	DTMILS (SEQ	DTMLLKS (SEQ	DTMILRS (SEQ	DTMLLHS (SEQ	DTMILPS (SEQ	DTMLLTS (SEQ
ID NO:1183)	ID NO:1184)	ID NO:1185)	ID NO:1186)	ID NO:1187)	ID NO:1188)	ID NO:1189)	ID NO:1190)	ID NO:1191)	ID NO:1192)	ID NO:1193)
DT RFQH S (SEQ	DTREQES (SEQ	DIREQTS (SEQ	DTRFQDS (SEQ	DTRLLAS (SEQ	DTRLISS (SEQ	DTRLIKS (SEQ	DTRLLRS (SEQ	DTRLLES (SEQ	DTRLLPS (SEQ	DTRLLTS (SEQ
ID NO:1048)	ID NO:1049)	ID NO:1050)	ID NO:1051)	ID NO:1052)	ID NO:1053)	ID NO:1054)	ID NO:1055)	ID NO:1056)	ID NO:1057)	ID NO:1058)
DTXFQHS (SEQ	DT <u>YFQP</u> S (SEQ	DTXFQTS (SEQ	DT <u>veod</u> s (Seq	DT <u>YL</u> LAS (SEQ	DT <u>YL</u> LSS (SEQ	DT <u>YL</u> LKS (SEQ	DT <u>vi</u> lrs (SEQ	DT <u>YL</u> LHS (SEQ	DT <u>YLLP</u> S (SEQ	DT <u>YL</u> LTS (SEQ
ID NO:915)	ID NO:916)	ID NO:917)	ID NO:918)	ID NO:919)	ID NO:920)	ID NO:921)	ID NO:922)	ID NO:923)	ID NO:924)	ID NO:925)
DT feqh s (Seq	DT FFQP S (SEQ	DTEFQTS (SEQ	DTFFQDS (SEQ	DTFLLAS (SEQ	DTFLLSS (SEQ	DTELLES (SEQ	DTFLIRS (SEQ	DTELLES (SEQ	DTFLLPS (SEQ	DTFLLTS (SEQ
ID NO:771)	ID NO:772)	ID NO:773)	ID NO:774)	ID NO:775)	ID NO:776)	ID NO:777)	ID NO:778)	ID NO:779)	ID NO:780)	ID NO:781)
DTS FQH S (SEQ	DTS FQP S (SEQ	DTS FQT S (SEQ	DTS FQD S (SEQ	DTS <u>r</u> LAS (SEQ	DTS <u>L</u> LSS (SEQ	DTS <u>L</u> LKS (SEQ	DTS <u>I</u> LRS (SEQ	DTSLLHS (SEQ	DTSLLPS (SEQ	DTS <u>t</u> l <u>t</u> s(Seq
ID NO:634)	ID NO: 635)	ID NO:636)	ID NO:637)	ID NO:124)	ID NO:638)	ID NO:639)	ID NO:640)	ID NO:641)	ID NO:642)	Id No:643)

DTS <u>t</u> l d s (SEQ	DTELLDS (SEQ	DT <u>ve</u> los (SEQ	DTRILDS (SEQ	DTMILDS (SEQ	DT KL L D S (SEQ	DT <u>LL</u> LDS (SEQ
ID NO:126)	ID NO:782)	ID NO:926)	ID NO:1059)	ID NO:1194)	ID NO:1338)	ID NO:75)
DTSLHAS (SEQ	DTFLHAS (SEQ	DT <u>YLH</u> AS (SEQ	DTRLHAS (SEQ	DTMLHAS (SEQ	DT KLH AS (SEQ	DT <u>LLH</u> AS (SEQ
ID NO:644)	ID NO:783)	ID NO:927)	ID NO:1060)	ID NO:1195)	ID NO:1339)	ID NO:1479)
DTSLHSS (SEQ	DT ELHS S (SEQ	DTYLHSS (SEQ	DTRLHSS (SEQ	DTMLHSS (SEQ	DTKLHS (SEQ	DTLLHSS (SEQ
ID NO:645)	ID NO:784)	ID NO:928)	ID NO:1061)	ID NO:1196)	ID NO:1340)	ID NO:1480)
DTS LHK S (SEQ	DT FLHK S (SEQ	DT <u>YLHK</u> S (SEQ	DTRLHKS (SEQ	DTMLHKS (SEQ	DT KLHK S (SEQ	DTLLHKS (SEQ
ID NO:646)	ID NO:785)	ID NO:929)	ID NO:1062)	ID NO:1197)	ID NO:1341)	ID NO:1481)
DTS LHR S (SEQ	DT FLHR S (SEQ	DT <u>vihr</u> S (SEQ	DTRLHRS (SEQ	DTMLHRS (SEQ	DTKLHRS (SEQ	DT LIHR S (SEQ
ID NO:647)	ID NO:786)	ID NO:930)	ID NO:1063)	ID NO:1198)	ID NO:1342)	ID NO:1482)
DTSLHHS (SEQ	DTFLHHS (SEQ	DT <u>vlhh</u> s (SEQ	DTRLHHS (SEQ	DTMLHHS (SEQ	DTKLHHS (SEQ	DT LLHH S (SEQ
ID NO:648)	ID NO:787)	ID NO:931)	ID NO:1064)	ID NO:1199)	ID NO:1343)	ID NO:1483)
DTS LHP S (SEQ	DTELHPS (SEQ	DT <u>YLHP</u> S (SEQ	DIRLHPS (SEQ ID NO:1065)	DTMLHPS (SEQ	DT KLHP S (SEQ	DTLLHPS (SEQ
ID NO:649)	ID NO:788)	ID NO:932)		ID NO:1200)	ID NO:1344)	ID NO:1484)
DTSLHTS (SEQ	DT FLHT S (SEQ	DT <u>YLHT</u> S (SEQ	DTRLHTS (SEQ	DTMLHTS (SEQ ID NO:1201)	DTKLHTS (SEQ	DTLLHTS (SEQ
ID NO:650)	ID NO:789)	ID NO:933)	ID NO:1066)		ID NO:1345)	ID NO:1485)
DTSLHDS (SEQ	DTFLHDS (SEQ	DT <u>YLHD</u> S (SEQ	DTRLHDS (SEQ ID NO:1067)	DTMLHDS (SEQ	DT KLHD S (SEQ	DT <u>LLHD</u> S (SEQ
ID NO:651)	ID NO:790)	ID NO:934)		ID NO:1202)	ID NO:1346)	ID NO:1486)
DTS <u>LQ</u> AS (SEQ	DT <u>FLQ</u> AS (SEQ	DT <u>YLQ</u> AS (SEQ	DTRLQAS (SEQ	DTMLQAS (SEQ	DTKLQAS (SEQ	DT <u>LLQ</u> AS (SEQ
ID NO:652)	ID NO:791)	ID NO:935)	ID NO:1068)	ID NO:1203)	ID NO:1347)	ID NO:1487)
DTS <u>LQS</u> S (SEQ	DT FLQS S (SEQ	DT <u>YLQS</u> S (SEQ	DTRLQS (SEQ	DTMIQS (SEQ	DT KLQS S (SEQ	DT <u>LLQS</u> S (SEQ
ID NO:653)	ID NO:792)	ID NO:936)	ID NO:1069)	ID NO:1204)	ID NO:1348)	ID NO:1488)

DTS LQK S (SEQ ID NO:654)	DTFLOKS (SEQ ID NO:793)	DT <u>YLOK</u> S (SEQ ID NO:937)	DTRIQKS (SEQ DTMIQKS (SEQ ID NO:1070) ID NO:1205)	DT MLQK S (SEQ ID NO:1205)	DTKLOKS (SEQ DTLLOKS (SEQ ID NO:1349)	DT <u>LLOK</u> S (SEQ ID NO:1489)
DTS <u>LOR</u> S (SEQ	DTFLQRS (SEQ	DT <u>vior</u> S (SEQ	DTRLQRS (SEQ	DTMLQRS (SEQ	DT KLQR S (SEQ	DTLLORS (SEQ
ID NO:655)	ID NO:794)	ID NO:938)	ID NO:1071)	ID NO:1206)	ID NO:1350)	ID NO:1490)
DTSLQHS (SEQ	DTFLQHS (SEQ	DT <u>YLQH</u> S (SEQ	DTRLQHS (SEQ	DIMIQHS (SEQ	DT KLQH S (SEQ	DT LLQH S (SEQ
ID NO:656)	ID NO:795)	ID NO:939)	ID NO:1072)	ID NO:1207)	ID NO:1351)	ID NO:1491)
DTS <u>LQP</u> S (SEQ	DTFLQPS (SEQ	DT <u>YLQP</u> S (SEQ	DTRLQPS (SEQ	DIMIQPS (SEQ	DTKLQPS (SEQ	DT LLQP S (SEQ
ID NO:657)	ID NO:796)	ID NO:940)	ID NO:1073)	ID NO:1208)	ID NO:1352)	ID NO:1492)
DTS <u>LOT</u> S (SEQ	DIFILOTS (SEQ	DT <u>YLQT</u> S (SEQ	DTRLQTS (SEQ	DTMIQIS (SEQ	DTKLQTS (SEQ	DT <u>LLQT</u> S (SEQ
ID NO:658)	ID NO:797)	ID NO:941)	ID NO:1074)	ID NO:1209)	ID NO:1353)	ID NO:1493)
DTS <u>LOD</u> S (SEQ	DTELODS (SEQ	DT <u>YLQD</u> S (SEQ	DTRLODS (SEQ	DTMLQDS (SEQ	DTKLQDS (SEQ	DTLLODS (SEQ
ID NO:659)	ID NO:798)	ID NO:942)	ID NO:1075)	ID NO:1210)	ID NO:1354)	ID NO:1494)
				,		

Bold faced & underlined amino acid residues are the residues which differ from the amino acid sequence in palivizumab

Table 3F - VL CDR3 Sequences

			((
2	(9	61)	1495	1496
	NO: 6	NO: 61	NO:	NO:
	ID	ID	ID	ΠI
	OES)	Õ∃S)	ÕES)	ÕES)
TO OTO T	FQGSGYPFT	FOGSEYPFT	FOGSTYPFT	FOGSWYPFT

Bold faced and underlined amino acid residues are the residues which differ from the amino acid sequence in palivizumab

[00182] "Iff one embodiment, formulations of the present invention comprise antibodies that comprise a VH CDR1 having the amino acid sequence of SEQ ID NO:1, SEQ ID NO:10 or SEQ ID NO:18. In another embodiment, formulations of the present invention comprise antibodies that comprise a VH CDR2 having the amino acid sequence of SEQ ID NO:2, SEQ ID NO:19, SEQ ID NO:25, SEQ ID NO:37, SEQ ID NO:41, SEQ ID NO:45, SEQ ID NO:305, or SEO ID NO:329. In another embodiment, formulations of the present invention comprise antibodies that comprise a VH CDR3 having the amino acid sequence of SEQ ID NO:3, SEQ ID NO:12, SEQ ID NO:20, SEQ ID NO:29, SEQ ID NO:79, or SEQ ID NO:311. In another embodiment, formulations of the present invention comprise antibodies that comprise a VH CDR1 having the amino acid sequence of SEQ ID NO:1, SEQ ID NO:10 or SEQ ID NO:18, a VH CDR2 having the amino acid sequence of SEQ ID NO:2, SEQ ID NO:19, SEQ ID NO:25, SEQ ID NO:37, SEQ ID NO:41, SEQ ID NO:45, SEQ ID NO:305, or SEQ ID NO:329, and a VH CDR3 having the amino acid sequence of SEQ ID NO:3, SEQ ID NO:12, SEQ ID NO:20, SEQ ID NO:29, SEQ ID NO:79, or SEQ ID NO:311. In a preferred embodiment, formulations of the present invention comprise antibodies that comprise a VH CDR1 having the amino acid sequence of SEQ ID NO:10, a VH CDR2 having the amino acid sequence of SEQ ID NO:19, and a VH CDR3 having the amino acid sequence of SEQ ID NO:20. In accordance with these embodiments, the antibodies immunspecifically bind to a RSV F antigen. In specific embodiments, the antibodies are not palivizumab, a Fab fragment of palivizumab, or an antigenbinding fragment thereof. In specific embodiments, the antibodies have a high affinity for a RSV antigen (e.g., RSV F antigen).

[00	183]		In o	ne e	nbod	imen	t, the	amir	io aci	d sec	luenc	e of 1	the V	H domain is
Q	V	T	L	R	E	ន	G	P	A	L	V	K	P	T
Q	Т	L	${f T}$	L	T	С	т	F	S	G	F	S	L	S
<u>T</u>	A	G	M	S	V	G	W	I	R	Q	P	P	G	К
A	L	E	W	Ŀ	A	D_	I	W	W	D	D_	K	K	H
<u>Y</u>	N	P	S	L	K	D	R	L	T	I	S	K	D	T
s	K	N	Q	V	V	L	K	V	T	N	M	D	P	A
D	т	A	т	Y	Y	С	A	R	D	M	I	F	N	F
<u>Y</u>	_F_	D	V	W	G	Q*	G	T	T	V	Т	V	S	S

(SEQ ID NO:48), wherein the three underlined regions indicate the VH CDR1, CDR2, and CDR3 regions, respectively; the four non-underlined regions correlate with the VL FR1, FR2, FR3, FR4, respectively; and the asterisk indicates the position of an A→Q mutation in VH FR4

as compared to the VH FR4 of palivizumab shown in Figure 1B(SEQ ID NO:7). This VH domain (SEQ ID NO:48) is identical to that of the motavizumab antibody described elsewhere herein and shown in Figure 13A. In some embodiments, this VH FR can be used in combination with any of the VH CDRs identified in Table 1 and/or Tables 3A-C. In one embodiment, the motavizumab antibody comprises the VH domain of Figure 13A (SEQ ID NO:208) and the C-gamma-1 (nG1m) constant domain described in Johnson et al. (1997) *J. Infect. Dis.* 176, 1215-1224 and U.S. Patent No. 5,824,307. In one embodiment, an antibody of the invention comprises a VH chain having the amino acid sequence of SEQ ID NO:208.

[00184] The present invention provides antibodies that immunospecifically bind to one or more RSV antigens (e.g., RSV F antigen), said antibodies comprising a VL chain having an amino acid sequence of any one of the VL chain listed in Table 2. In certain embodiments, the antibody is not palivizumab and/or the VL chain is not the VL chain of palivizumab.

[00185] The present invention also provides antibodies that immunospecifically bind to one or more RSV antigens (e.g., RSV F antigens), said antibodies comprising a variable light ("VL") domain having an amino acid sequence of any one of the VL domains listed in Table 2. In certain embodiments, the antibody is not palivizumab and/or the VH domain is not the VH domain of palivizumab. The present invention also provides antibodies that immunospecifically bind to one or more RSV antigens (e.g., RSV F antigens), said antibodies comprising a VL CDR having an amino acid sequence of any one of the VL CDRs listed in Table 2 and/or Tables 3D-3F. In certain embodiments, the antibody is not palivizumab. In some embodiments, the antibody comprises one, two or three of the VL CDRs listed in Table 2 and/or Tables 3D-3F.

In one embodiment of the present invention, antibodies comprise a VL CDR1 having the amino acid sequence of SEQ ID NO:4, SEQ ID NO:14, SEQ ID NO:22, SEQ ID NO:31, SEQ ID NO:39, SEQ ID NO:47, SEQ ID NO:72, SEQ ID NO:314, SEQ ID NO:320, or SEQ ID NO:335. In another embodiment, formulations of the invention comprise antibodies that comprise a VL CDR2 having the amino acid sequence of SEQ ID NO:5, SEQ ID NO:15, SEQ ID NO:23, SEQ ID NO:27, SEQ ID NO:32, SEQ ID NO:35, SEQ ID NO:43, SEQ ID NO:50, SEQ ID NO:53, SEQ ID NO:57, SEQ ID NO:59, SEQ ID NO:63, SEQ ID NO:66, SEQ ID NO:69, SEQ ID NO:73, SEQ ID NO:75, SEQ ID NO:77, SEQ ID NO:308, SEQ ID NO:315, SEQ ID NO:321, SEQ ID NO:326, SEQ ID NO:332, or SEQ ID NO:336. In another embodiment, formulations of the invention comprise antibodies that comprise a VL CDR3 having the amino acid sequence of SEQ ID NO:6, SEQ ID NO:16 or SEQ ID NO:61. In another embodiment, formulations of the invention comprise antibodies that comprise a VL CDR1

having the amino acid sequence of SEQ ID NO:4, SEQ ID NO:14, SEQ ID NO:22, SEQ ID NO:31, SEQ ID NO:39, SEQ ID NO:47, SEQ ID NO:72, SEQ ID NO:314, SEQ ID NO:320, or SEQ ID NO:335, a VL CDR2 having the amino acid sequence of SEQ ID NO:5, SEQ ID NO:43, SEQ ID NO:15, SEQ ID NO:23, SEQ ID NO:27, SEQ ID NO:32, SEQ ID NO:35, SEQ ID NO:43, SEQ ID NO:50, SEQ ID NO:53, SEQ ID NO:57, SEQ ID NO:59, SEQ ID NO:63, SEQ ID NO:66, SEQ ID NO:69, SEQ ID NO:73, SEQ ID NO:75, SEQ ID NO:77, SEQ ID NO:308, SEQ ID NO:315, SEQ ID NO:321, SEQ ID NO:326, SEQ ID NO:332, or SEQ ID NO:336, and a VL CDR3 having the amino acid sequence of SEQ ID NO:6, SEQ ID NO:16 or SEQ ID NO:61. In a preferred embodiment, formulations of the invention comprise antibodies that comprise a VL CDR1 having the amino acid sequence of SEQ ID NO:39, a VLCDR2 having the amino acid sequence of SEQ ID NO:5, and a VLCDR3 having the amino acid sequence of SEQ ID NO:6. In accordance with these embodiments, the antibodies immunospecifically bind to a RSV F antigen. In specific embodiments, the antibodies are not palivizumab or an antigen-binding fragment thereof (e.g., a Fab fragment of palivizumab). In another specific embodiment, the antibodies have a high affinity for RSV antigen (e.g., RSV F antigen).

[00187]	In one embodiment the amino acid sequence of the VL domain is
---------	---

D	I	Q	M	T	Q	S	P	S	Т	L	S	A	S	V	
G	D	R	V	T	I	T	С	s	A	S_	S	R	V	G	
<u>Y</u>	_M	H	W	Y	Q	Q	ĸ	P	G	K	A	P	K	L	
L	I	Y	<u>D</u>	Т	S	Ķ	L	A	S	G	v	P	s	R	
F	S	G	S	G	S	G	T	E	F	Т	L	T	I	S	
ន	L	Q	P	D	D	F	A	Т	Y	Y	С	F	0	G	
S	G	Y	P	F	<u>T</u>	F	G	G	G	т	K	V*	E	I	

K

(SEQ ID NO:8), wherein the three underlined regions indicate the VL CDR1, CDR2, and CDR3 regions, respectively; the four non-underlined regions correlate with the VL FR1, FR2, FR3, FR4, respectively; and the asterisk indicates the position of an L→V mutation in VL FR4 as compared to the VL FR4 of palivizumab shown in Figure 1A. This VL domain (SEQ ID NO:8) is identical to that of the motavizumab antibody described elsewhere herein and shown in Figure 13B. In some embodiments, this VL framework can be used in combination with any of the VL CDRs identified in Table 1 and/or Tables 3D-3F. In one embodiment, the motavizumab

antibody comprises the VL domain of Figure 13B (SEQ ID NO:209) and the C-kappa constant domain described in Johnson et al. (1997) *J. Infect. Dis.* 176, 1215-1224 and U.S. Patent No. 5,824,307. In one embodiment, an antibody of the invention comprises a VL chain having the amino acid sequence of SEQ ID NO:209.

[00188] The present invention further provides antibodies that immunospecifically bind to one or more RSV antigens (e.g., RSV F antigen), wherein the antibody comprises a VH chain disclosed herein combined with a VL chain disclosed herein, or other VL chain. The present invention also provides antibodies that immunospecifically bind to one or more RSV antigens (e.g., RSV F antigen), wherein the antibody comprises a VL chain disclosed herein combined with a VH chain disclosed herein, or other VH chain.

[00189] The present invention also provides antibodies that immunospecifically bind to one or more RSV antigens (e.g., RSV F antigens), said antibodies comprising a VH domain disclosed herein combined with a VL domain disclosed herein, or other VL domain. The present invention further provides antibodies that immunospecifically bind to one or more RSV antigens (e.g., RSV F antigens), said antibodies comprising a VL domain disclosed herein combined with a VH domain disclosed herein, or other VH domain.

[00190] In a specific embodiment, antibodies that immunospecifically bind to a RSV antigen (e.g., RSV F antigens) comprise a VH domain having the amino acid sequence of SEO ID NO:7, SEQ ID NO:9, SEQ ID NO:17, SEQ ID NO:24, SEQ ID NO:28, SEQ ID NO:33, SEQ ID NO:36, SEQ ID NO:40, SEQ ID NO:44, SEQ ID NO:48, SEQ ID NO:51, SEQ ID NO:55, SEQ ID NO:67, SEQ ID NO:78, SEQ ID NO:304, SEQ ID NO:310, SEQ ID NO:317, SEQ ID NO:323, or SEQ ID NO:328, and a VL domain having the amino acid sequence of SEO ID NO:8, SEQ ID NO:13, SEQ ID NO:21, SEQ ID NO:26, SEQ ID NO:30, SEQ ID NO:34, SEQ ID NO:38, SEQ ID NO:42, SEQ ID NO:46, SEQ ID NO:49, SEQ ID NO:52, SEQ ID NO:54, SEO ID NO:56, SEO ID NO:58, SEQ ID NO:60, SEQ ID NO:62, SEQ ID NO:64, SEQ ID NO:65, SEQ ID NO:68, SEQ ID NO:70, SEQ ID NO:71, SEQ ID NO:74, SEQ ID NO:76, SEQ ID NO:307, SEQ ID NO:313, SEQ ID NO:319, SEQ ID NO:325, SEQ ID NO:331, or SEQ ID NO:334. In a preferred embodiment, antibodies that immunospecifically bind to a RSV F antigen comprise a VH domain having the amino acid sequence of SEQ ID NO:48 and a VL domain comprising the amino acid sequence of SEQ ID NO:11. In specific embodiments, the antibodies are not palivizumab or an antigen-binding fragment thereof (e.g., a Fab fragment). In another specific embodiment, the antibodies of the invention have a high affinity for a RSV antigen (e.g., RSV F antigen).

The present invention further provides antibodies that specifically bind to an RSV [00191]antigen (e.g., RSV F antigen), wherein the antibody comprises any VH CDR1 disclosed herein, optionally in combination with any VH CDR2 disclosed herein (or other VH CDR2), and/or optionally in combination with any VH CDR3 disclosed herein (or other VH CDR3)), and/or optionally in combination with any VL CDR1 disclosed herein (or other VL CDR1), and/or optionally in combination with any VL CDR2 disclosed herein (or other VL CDR2), and/or optionally in combination with any VL CDR3 disclosed herein (or other VL CDR3). The present invention also provides antibodies that specifically bind to an RSV antigen (e.g., RSV F antigen), wherein the antibody comprises any VH CDR2 disclosed herein, optionally in combination with any VH CDR1 disclosed herein (or other VH CDR1), and/or optionally in combination with any VH CDR3 disclosed herein (or other VH CDR3)), and/or optionally in combination with any VL CDR1 disclosed herein (or other VL CDR1), and/or optionally in combination with any VL CDR2 disclosed herein (or other VL CDR2), and/or optionally in combination with any VL CDR3 disclosed herein (or other VL CDR3). The present invention also provides antibodies that specifically bind to an RSV antigen (e.g., RSV F antigen), wherein the antibody comprises any VH CDR3 disclosed herein, optionally in combination with any VH CDR1 disclosed herein (or other VH CDR1), and/or optionally in combination with any VH CDR2 disclosed herein (or other VH CDR3)), and/or optionally in combination with any VL CDR1 disclosed herein (or other VL CDR1), and/or optionally in combination with any VL CDR2 disclosed herein (or other VL CDR2), and/or optionally in combination with any VL CDR3 disclosed herein (or other VL CDR3). The present invention also provides antibodies that specifically bind to an RSV antigen (e.g., RSV F antigen), wherein the antibody comprises any VL CDR1 disclosed herein, optionally in combination with any VH CDR1 disclosed herein (or other VH CDR1), and/or optionally in combination with any VH CDR2 disclosed herein (or other VH CDR2)), and/or optionally in combination with any VH CDR3 disclosed herein (or other VH CDR3), and/or optionally in combination with any VL CDR2 disclosed herein (or other VL CDR2), and/or optionally in combination with any VL CDR3 disclosed herein (or other VL CDR3). The present invention further provides antibodies that specifically bind to an RSV antigen (e.g., RSV F antigen), wherein the antibody comprises any VL CDR2 disclosed herein, optionally in combination with any VH CDR1 disclosed herein (or other VH CDR1), and/or optionally in combination with any VH CDR2 disclosed herein (or other VH CDR2)), and/or optionally in combination with any VH CDR3 disclosed herein (or other VH CDR3), and/or optionally in combination with any VL CDR1 disclosed herein (or other VL CDR1), and/or optionally in combination with any VL CDR3 disclosed herein (or other VL CDR3). The present invention also provides antibodies that specifically bind to an RSV antigen (e.g., RSV F

antigen), wherein the antibody comprises any VL CDR3 disclosed herein, optionally in combination with any VH CDR1 disclosed herein (or other VH CDR1), and/or optionally in combination with any VH CDR2 disclosed herein (or other VH CDR2)), and/or optionally in combination with any VH CDR3 disclosed herein (or other VH CDR3), and/or optionally in combination with any VL CDR1 disclosed herein (or other VL CDR1), and/or optionally in combination with any VL CDR2 disclosed herein (or other VL CDR2).

[00192] The present invention also provides antibodies comprising one or more VH CDRs and one or more VL CDRs listed in Table 2 and/or Tables 3A-3F. In particular, the invention provides for an antibody comprising a VH CDR1 and a VL CDR1; a VH CDR1 and a VL CDR2; a VH CDR1 and a VL CDR3; a VH CDR2 and a VL CDR1; VH CDR2 and VL CDR2; a VH CDR2 and a VL CDR3; a VH CDR3 and a VH CDR1; a VH CDR3 and a VL CDR2; a VH CDR3 and a VL CDR3; a VH1 CDR1, a VH CDR2 and a VL CDR1; a VH CDR1, a VH CDR2 and a VL CDR2; a VH CDR1, a VH CDR2 and a VL CDR3; a VH CDR2, a VH CDR3 and a VL CDR1, a VH CDR2, a VH CDR3 and a VL CDR2; a VH CDR2, a VH CDR2 and a VL CDR3; a VH CDR1, a VL CDR1 and a VL CDR2; a VH CDR1, a VL CDR1 and a VL CDR3; a VH CDR2, a VL CDR1 and a VL CDR2; a VH CDR2, a VL CDR1 and a VL CDR3; a VH CDR3, a VL CDR1 and a VL CDR2; a VH CDR3, a VL CDR1 and a VL CDR3; a VH CDR1, a VH CDR2, a VH CDR3 and a VL CDR1; a VH CDR1, a VH CDR2, a VH CDR3 and a VL CDR2; a VH CDR1, a VH CDR2, a VH CDR3 and a VL CDR3; a VH CDR1, a VH CDR2, a VL CDR1 and a VL CDR2; a VH CDR1, a VH CDR2, a VL CDR1 and a VL CDR3; a VH CDR1, a VH CDR3, a VL CDR1 and a VL CDR2; a VH CDR1, a VH CDR3, a VL CDR1 and a VL CDR3; a VH CDR2, a VH CDR3, a VL CDR1 and a VL CDR2; a VH CDR2, a VH CDR3, a VL CDR1 and a VL CDR3; a VH CDR2, a VH CDR3, a VL CDR2 and a VL CDR3; a VH CDR1, a VH CDR2, a VH CDR3, a VL CDR1 and a VL CDR2; a VH CDR1, a VH CDR2, a VH CDR3, a VL CDR1 and a VL CDR3; a VH CDR1, a VH CDR2, a VL CDR1, a VL CDR2, and a VL CDR3; a VH CDR1, a VH CDR3, a VL CDR1, a VL CDR2, and a VL CDR3; a VH CDR2, a VH CDR3, a VL CDR1, a VL CDR2, and a VL CDR3; or any combination thereof of the VH CDRs and VL CDRs listed in Table 2 and/or Tables 3A-3F. In a specific embodiment, the formulations of the invention comprise antibodies that have a high affinity for a RSV antigen (e.g., RSV F antigen).

[00193] The invention also provides an antibody that immunospecifically binds to a RSV F antigen, comprising a VH CDR1 and a VL CDR1, a VH CDR1 and a VL CDR2, a VH CDR1 and a VL CDR3, a VH CDR1 and a VL CDR1 and a VL CDR2; a VH CDR1 and

a VL CDR3; a VH CDR2 and a VL CDR1; VH CDR2 and VL CDR2; a VH CDR2 and a VL CDR3; a VH CDR3 and a VH CDR1; a VH CDR3 and a VL CDR2; a VH CDR3 and a VL CDR3; a VH1 CDR1, a VH CDR2 and a VL CDR1; a VH CDR1, a VH CDR2 and a VL CDR2; a VH CDR1, a VH CDR2 and a VL CDR3; a VH CDR2, a VH CDR3 and a VL CDR1, a VH CDR2, a VH CDR3 and a VL CDR2; a VH CDR2, a VH CDR2 and a VL CDR3; a VH CDR1, a VL CDR1 and a VL CDR2; a VH CDR1, a VL CDR1 and a VL CDR3; a VH CDR2, a VL CDR1 and a VL CDR2; a VH CDR2, a VL CDR1 and a VL CDR3; a VH CDR3, a VL CDR1 and a VL CDR2; a VH CDR3, a VL CDR1 and a VL CDR3; a VH CDR1, a VH CDR2, a VH CDR3 and a VL CDR1; a VH CDR1, a VH CDR2, a VH CDR3 and a VL CDR2; a VH CDR1, a VH CDR2, a VH CDR3 and a VL CDR3; a VH CDR1, a VH CDR2, a VL CDR1 and a VL CDR2; a VH CDR1, a VH CDR2, a VL CDR1 and a VL CDR3; a VH CDR1, a VH CDR3, a VL CDR1 and a VL CDR2; a VH CDR1, a VH CDR3, a VL CDR1 and a VL CDR3; a VH CDR2, a VH CDR3, a VL CDR1 and a VL CDR2; a VH CDR2, a VH CDR3, a VL CDR1 and a VL CDR3; a VH CDR2, a VH CDR3, a VL CDR2 and a VL CDR3; a VH CDR1, a VH CDR2, a VH CDR3, a VL CDR1 and a VL CDR2; a VH CDR1, a VH CDR2, a VH CDR3, a VL CDR1 and a VL CDR3; a VH CDR1, a VH CDR2, a VL CDR1, a VL CDR2, and a VL CDR3; a VH CDR1, a VH CDR3, a VL CDR1, a VL CDR2, and a VL CDR3; a VH CDR2, a VH CDR3, a VL CDR1, a VL CDR2, and a VL CDR3; or any combination thereof of the VH CDRs and VL CDRs listed in Table 2 and/or Tables 3A-3F, supra. In another specific embodiment, the formulations of the invention comprise antibodies that have a high affinity for a RSV antigen (e.g., RSV F antigen).

[00194] In one embodiment, a formulation of the invention comprises an antibody that comprises a VH CDR1 having the amino acid sequence of SEQ ID NO:1, SEQ ID NO:10 or SEQ ID NO:18 and a VL CDR1 having the amino acid sequence of SEQ ID NO:4, SEQ ID NO:14, SEQ ID NO:22, SEQ ID NO:31, SEQ ID NO:39, SEQ ID NO:47, SEQ ID NO:314, SEQ ID NO:320, or SEQ ID NO:335. In another embodiment, a formulation of the invention comprises an antibody that comprises a VH CDR1 having the amino acid sequence of SEQ ID NO:1, SEQ ID NO:10 or SEQ ID NO:18 and a VL CDR2 having the amino acid sequence of SEQ ID NO:5, SEQ ID NO:15, SEQ ID NO:23, SEQ ID NO:27, SEQ ID NO:32, SEQ ID NO:35, SEQ ID NO:43, SEQ ID NO:50, SEQ ID NO:53, SEQ ID NO:57, SEQ ID NO:59, SEQ ID NO:63, SEQ ID NO:66, SEQ ID NO:69, SEQ ID NO:73, SEQ ID NO:75, SEQ ID NO:77, SEQ ID NO:308, SEQ ID NO:315, SEQ ID NO:321, SEQ ID NO:326, SEQ ID NO:332, or SEQ ID NO:336. In another embodiment, a formulation of the invention comprises an antibody that comprises a VH CDR1 having the amino acid sequence of SEQ ID NO:1, SEQ ID NO:10 or

SEQ ID NO:18 and a VL CDR3 having the amino acid sequence of SEQ ID NO:6, SEQ ID NO:16 or SEQ ID NO:61. In accordance with these embodiments, the antibody immunospecifically binds to a RSV F antigen.

In another embodiment, a formulation of the invention comprises an antibody [00195] that comprises a VH CDR2 having the amino acid sequence of SEQ ID NO:2, SEQ ID NO:19, SEO ID NO:25, SEQ ID NO:37, SEQ ID NO:41, SEQ ID NO:45, SEQ ID NO:305, or SEQ ID NO:329, and a VL CDR1 having the amino acid sequence of SEQ ID NO:4, SEQ ID NO:14, SEQ ID NO:22, SEQ ID NO:31, SEQ ID NO:39, SEQ ID NO:47, SEQ ID NO:314, SEQ ID NO:320, or SEO ID NO:335. In another embodiment, an antibody of the invention comprises a VH CDR2 having the amino acid sequence of SEQ ID NO:2, SEQ ID NO:19, SEQ ID NO:25, SEO ID NO:37, SEO ID NO:41, SEQ ID NO:45, SEQ ID NO:305, or SEQ ID NO:329, and a VL CDR2 having the amino acid sequence of SEQ ID NO:5, SEQ ID NO:15, SEQ ID NO:23, SEQ ID NO:27, SEQ ID NO:32, SEQ ID NO:35, SEQ ID NO:43, SEQ ID NO:50, SEQ ID NO:53, SEQ ID NO:57, SEQ ID NO:59, SEQ ID NO:63, SEQ ID NO:66, SEQ ID NO:69, SEQ ID NO:73, SEO ID NO:75, SEQ ID NO:77, SEQ ID NO:308, SEQ ID NO:315, SEQ ID NO:321, SEQ ID NO:326, SEQ ID NO:332, or SEQ ID NO:336. In another embodiment, an antibody of the invention comprises a VH CDR2 having the amino acid sequence of SEQ ID NO:2, SEQ ID NO:19, SEQ ID NO:25, SEQ ID NO:37, SEQ ID NO:41, SEQ ID NO:45, SEQ ID NO:305, or SEO ID NO:329, and a VL CDR3 having the amino acid sequence of SEQ ID NO:6, SEQ ID NO:16, or SEQ ID NO:61. In accordance with these embodiments, the antibody immunospecifically binds to a RSV F antigen.

In another embodiment, a formulation of the invention comprises an antibody that comprises a VH CDR3 having the amino acid sequence of SEQ ID NO:3, SEQ ID NO:12, SEQ ID NO:20, SEQ ID NO:29, SEQ ID NO:79, or SEQ ID NO:311, and a VL CDR1 having the amino acid sequence of SEQ ID NO:4, SEQ ID NO:14, SEQ ID NO:22, SEQ ID NO:31, SEQ ID NO:39, SEQ ID NO:47, SEQ ID NO:314, SEQ ID NO:320, or SEQ ID NO:335. In another embodiment, a formulation of the invention comprises an antibody that comprises a VH CDR3 having the amino acid sequence of SEQ ID NO:3, SEQ ID NO:12, SEQ ID NO:20, SEQ ID NO:29, SEQ ID NO:79, or SEQ ID NO:311, and a VL CDR2 having the amino acid sequence of SEQ ID NO:3, SEQ ID NO:27, SEQ ID NO:32, SEQ ID NO:35, SEQ ID NO:43, SEQ ID NO:50, SEQ ID NO:53, SEQ ID NO:57, SEQ ID NO:59, SEQ ID NO:66, SEQ ID NO:69, SEQ ID NO:73, SEQ ID NO:75, SEQ ID NO:77, SEQ ID NO:308, SEQ ID NO:315, SEQ ID NO:321, SEQ ID NO:326, SEQ ID NO:308, SEQ ID NO:315, SEQ ID NO:321, SEQ ID NO:326, SEQ ID NO:308, SEQ ID NO:315, SEQ ID NO:321, SEQ ID NO:326, SEQ ID NO:308, SEQ ID NO:315, SEQ ID NO:321, SEQ ID NO:326, SEQ ID

NO:332, or SEQ ID NO:336. In a preferred embodiment, an antibody of the invention comprises a VH CDR3 having the amino acid sequence of SEQ ID NO:3, SEQ ID NO:12, SEQ ID NO:20, SEQ ID NO:29, SEQ ID NO:79, or SEQ ID NO:311, and a VL CDR3 having the amino acid sequence of SEQ ID NO:6, SEQ ID NO:16, or SEQ ID NO:61. In accordance with these embodiments, the antibody immunospecifically binds to a RSV F antigen.

The present invention provides antibodies that immunospecifically bind to a RSV [00197] F antigen, said antibodies comprising the amino acid sequence of the variable heavy domain and/or variable light domain or an antigen-binding fragment thereof of AFFF, P12f2, P12f4, P11d4, Ale9, A12a6, A13c4, A17d4, A4B4, A8C7, 1X-493L1FR, H3-3F4, M3H9, Y10H6, DG, AFFF(1), 6H8, L1-7E5, L2-15B10, A13a11, A1h5, A4B4(1), A4B4L1FR-S28R (motavizumab), A4B4-F52S, A17d4(1), A3e2, A14a4, A16b4, A17b5, A17f5, or A17h4 with one or more amino acid residue substitutions in the variable heavy domain and/or variable light domain or antigenbinding fragment. The present invention also provides antibodies that immunospecifically bind to a RSV antigen, said antibodies comprising the amino acid sequence of the variable heavy domain and/or variable light domain or an antigen-binding fragment thereof of AFFF, P12f2, P12f4, P11d4, Ale9, A12a6, A13c4, A17d4, A4B4, A8C7, 1X-493L1FR, H3-3F4, M3H9, Y10H6, DG, AFFF(1), 6H8, L1-7E5, L2-15B10, A13a11, A1h5, A4B4(1), A4B4L1FR-S28R (motavizumab), A4B4-F52S, A17d4(1), A3e2, A14a4, A16b4, A17b5, A17f5, or A17h4 with one or more amino acid residue substitutions in one or more VH CDRs and/or one or more VL CDRs. Non-limiting examples of amino acid residues in the VH CDRs and VL CDRs of AFFF, P12f2, P12f4, P11d4, Ale9, A12a6, A13c4, A17d4, A4B4, A8C7, 1X-493L1FR, H3-3F4, M3H9, Y10H6, DG, AFFF(1), 6H8, L1-7E5, L2-15B10, A13a11, A1h5, A4B4(1), A4B4L1FR-S28R (motavizumab), A4B4-F52S, A17d4(1), A3e2, A14a4, A16b4, A17b5, A17f5, or A17h4, which may be substituted, are shown in bold in Table 2. The present invention also provides antibodies that immunospecifically bind to a RSV antigen, said antibodies comprising the amino acid sequence of the variable heavy domain and/or variable light domain or an antigen-binding fragment thereof of AFFF, P12f2, P12f4, P11d4, Ale9, A12a6, A13c4, A17d4, A4B4, A8C7, 1X-493L1FR, H3-3F4, M3H9, Y10H6, DG, AFFF(1), 6H8, L1-7E5, L2-15B10, A13a11, A1h5, A4B4(1), A4B4L1FR-S28R (motavizumab), A4B4-F52S, A17d4(1), A3e2, A14a4, A16b4, A17b5, A17f5, or A17h4 with one or more amino acid residue substitutions in one or more VH frameworks and/or one or more VL frameworks. The antibody generated by introducing substitutions in the VH domain, VH CDRs, VL domain, VL CDRs and/or frameworks of AFFF, P12f2, P12f4, P11d4, Ale9, A12a6, A13c4, A17d4, A4B4, A8C7, 1X-493L1FR, H3-3F4, M3H9, Y10H6, DG, AFFF(1), 6H8, L1-7E5, L2-15B10, A13a11, A1h5, A4B4(1), A4B4L1FR-

S28R (motavizumab), A4B4-F52S, A17d4(1), A3e2, A14a4, A16b4, A17b5, A17f5, or A17h4 can be tested *in vitro* and/or *in vivo*, for example, for its ability to bind to a RSV antigen, or for its ability to prevent, treat and/or ameliorate a an upper and/or lower respiratory tract RSV infection, otitis media, or one or more symptoms thereof.

[00198] In a specific embodiment, an antibody that immunospecifically binds to a RSV F antigen comprises an amino acid sequence encoded by a nucleotide sequence that hybridizes to the nucleotide sequence(s) encoding palivizumab, AFFF, P12f2, P12f4, P11d4, Ale9, A12a6, A13c4, A17d4, A4B4, A8C7, 1X-493L1FR, H3-3F4, M3H9, Y10H6, DG, AFFF(1), 6H8, L1-7E5, L2-15B10, A13a11, A1h5, A4B4(1), A4B4L1FR-S28R (MEDI-524, motavizumab), A4B4-F52S, A17d4(1), A3e2, A14a4, A16b4, A17b5, A17f5, A17h4, or an antigen-binding fragment thereof under stringent conditions, *e.g.*, hybridization to filter-bound DNA in 6x sodium chloride/sodium citrate (SSC) at about 45° C followed by one or more washes in 0.2xSSC/0.1% SDS at about 50-65° C, under highly stringent conditions, *e.g.*, hybridization to filter-bound nucleic acid in 6xSSC at about 45° C followed by one or more washes in 0.1xSSC/0.2% SDS at about 68° C, or under other stringent hybridization conditions which are known to those of skill in the art (see, for example, Ausubel, F.M. et al., eds., 1989, Current Protocols in Molecular Biology, Vol. I, Green Publishing Associates, Inc. and John Wiley & Sons, Inc., New York at pages 6.3.1-6.3.6 and 2.10.3).

[00199] In another embodiment, an antibody that immunospecifically binds to a RSV Fantigen comprises an amino acid sequence that is at least 35%, at least 40%, at least 45%, at least 50%, at least 55%, at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, or at least 99% identical to the amino acid sequence of AFFF, P12f2, P12f4, P11d4, Ale9, A12a6, A13c4, A17d4, A4B4, A8C7, 1X-493L1FR, H3-3F4, M3H9, Y10H6, DG, AFFF(1), 6H8, L1-7E5, L2-15B10, A13a11, A1h5, A4B4(1), A4B4L1FR-S28R (motavizumab), A4B4-F52S, A17d4(1), A3e2, A14a4, A16b4, A17b5, A17f5, or A17h4, or an antigen-binding fragment thereof. In preferred embodiment, an antibody that immunospecifically binds to a RSV F antigen comprises an amino acid sequence that is at least 35%, at least 40%, at least 45%, at least 50%, at least 55%, at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 99% identical to an amino acid sequence of A4B4L1FR-S28R (motavizumab), or an antigen-binding fragment thereof.

[00200] In a specific embodiment, an antibody that immunospecifically binds to a RSV F antigen comprises an amino acid sequence of a VH domain and/or an amino acid sequence a VL

domain encoded by a nucleotide sequence that hybridizes to the nucleotide sequence encoding any one of the VH and/or VL domains listed in Table 2 under stringent conditions, e.g., hybridization to filter-bound DNA in 6x sodium chloride/sodium citrate (SSC) at about 45° C followed by one or more washes in 0.2xSSC/0.1% SDS at about 50-65° C, under highly stringent conditions, e.g., hybridization to filter-bound nucleic acid in 6xSSC at about 45° C followed by one or more washes in 0.1xSSC/0.2% SDS at about 68° C, or under other stringent hybridization conditions which are known to those of skill in the art (see, for example, Ausubel, F.M. et al., eds., 1989, Current Protocols in Molecular Biology, Vol. I, Green Publishing Associates, Inc. and John Wiley & Sons, Inc., New York at pages 6.3.1-6.3.6 and 2.10.3). In another embodiment, an antibody that immunospecifically binds to a RSV antigen comprises an amino acid sequence of a VH CDR or an amino acid sequence of a VL CDRs encoded by a nucleotide sequence that hybridizes to the nucleotide sequence encoding any one of the VH CDRs or VL CDRs listed in Table 2 and/or Tables 3A-3F under stringent conditions e.g., hybridization to filter-bound DNA in 6X sodium chloride/sodium citrate (SSC) at about 45° C followed by one or more washes in 0.2X SSC/0.1% SDS at about 50-65° C, under highly stringent conditions, e.g., hybridization to filter-bound nucleic acid in 6X SSC at about 45° C followed by one or more washes in 0.1X SSC/0.2% SDS at about 68° C, or under other stringent hybridization conditions which are known to those of skill in the art. In yet another embodiment, an antibody that immunospecifically binds to a RSV F antigen comprises an amino acid sequence of a VH CDR and an amino acid sequence of a VL CDR encoded by nucleotide sequences that hybridizes to the nucleotide sequences encoding any one of the VH CDRs and VL CDRs, respectively, listed in Table 2 and/or Tables 3A-3F under stringent conditions, e.g., hybridization to filter-bound DNA in 6X sodium chloride/sodium citrate (SSC) at about 45° C followed by one or more washes in 0.2X SSC/0.1% SDS at about 50-65° C, under highly stringent conditions, e.g., hybridization to filter-bound nucleic acid in 6X SSC at about 45° C followed by one or more washes in 0.1X SSC/0.2% SDS at about 68° C, or under other stringent hybridization conditions which are known to those of skill in the art.

[00201] In another embodiment, an antibody that immunospecifically binds to a RSV F antigen comprises an amino acid sequence of a VH domain that is at least 35%, at least 40%, at least 45%, at least 50%, at least 55%, at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, or at least 99% identical to any one of the VH domains listed in Table 2. In another embodiment, an antibody that immunospecifically binds to a RSV antigen comprises an amino acid sequence of one or more VH CDRs that are at least 35%, at least 40%, at least 45%, at least 50%, at least 55%, at least 65%, at least 65%, at

least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, or at least 99% identical to any of the VH CDRs listed in Table 2 and/or Tables 3A-3C. In another embodiment, an antibody that immunospecifically binds to a RSV F antigen comprises an amino acid sequence of a VL domain that is at least 35%, at least 40%, at least 45%, at least 50%, at least 55%, at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, or at least 99% identical to any one of the VL domains listed in Table 2. In another embodiment, an antibody that immunospecifically binds to a RSV F antigen comprises an amino acid sequence of one or more VL CDRs that are at least 35%, at least 40%, at least 45%, at least 50%, at least 55%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, or at least 99% identical to any of the VL CDRs listed in Table 2 and/or Tables 3D-3F.

[00202] The present invention also provides antibodies that compete with an antibody or Fab fragment listed in Table 2 for binding to a RSV F antigen. The present invention also encompasses polypeptides, proteins and peptides comprising VL domains and/or VH domains that compete with a polypeptide, protein or peptide comprising a VL domain and/or a VH domain listed in Table 2 for binding to a RSV F antigen. Further, the present invention encompasses polypeptides, proteins and peptides comprising VL CDRs and/or VH CDRs that compete with a polypeptide, protein or peptide comprising a VL CDR and/or VH CDR listed in Table 2 and/or Tables 3A-3F for binding to a RSV F antigen.

[00203] The formulations of the present invention comprise antibodies that include derivatives that are modified, *i.e.*, by the covalent attachment of any type of molecule to the antibody such that covalent attachment. For example, but not by way of limitation, the antibody derivatives include antibodies that have been modified, *e.g.*, by glycosylation, acetylation, pegylation, phosphorylation, amidation, derivatization by known protecting/blocking groups, proteolytic cleavage, linkage to a cellular ligand or other protein, etc. Any of numerous chemical modifications may be carried out by known techniques, including, but not limited to specific chemical cleavage, acetylation, formylation, metabolic synthesis of tunicamycin, etc. Additionally, the derivative may contain one or more non-classical amino acids.

[00204] The present invention also provides antibodies that immunospecifically bind to a RSV antigen (e.g., RSV F antigen) which comprise a framework region known to those of skill in the art (e.g., a human or non-human fragment). The framework region may be naturally occurring or consensus framework regions. Preferably, the framework region of an antibody of the invention is human (see, e.g., Chothia et al., 1998, J. Mol. Biol. 278:457-479 for a listing of

human framework regions, which is incorporated by reference herein in its entirety). In a specific embodiment, an antibody of the invention comprises the framework region of A4B4L1FR-S28R (motavizumab).

[00205] In a specific embodiment, the present invention provides antibodies that immunospecifically bind to a RSV F antigen, said antibodies comprising the amino acid sequence of one or more of the CDRs of an antibody listed in Table 2 (i.e., AFFF, P12f2, P12f4, P11d4, Ale9, A12a6, A13c4, A17d4, A4B4, A8C7, 1X-493L1FR, H3-3F4, M3H9, Y10H6, DG, AFFF(1), 6H8, L1-7E5, L2-15B10, A13a11, A1h5, A4B4(1), A4B4L1FR-S28R (motavizumab), A4B4-F52S, A17d4(1), A3e2, A14a4, A16b4, A17b5, A17f5, or A17h4) and/or one or more of the CDRs in Table 3A-3F, and human framework regions with one or more amino acid substitutions at one, two, three or more of the following residues: (a) rare framework residues that differ between the murine antibody framework (i.e., donor antibody framework) and the human antibody framework (i.e., acceptor antibody framework); (b) Venier zone residues when differing between donor antibody framework and acceptor antibody framework; (c) interchain packing residues at the VH/VL interface that differ between the donor antibody framework and the acceptor antibody framework; (d) canonical residues which differ between the donor antibody framework and the acceptor antibody framework sequences, particularly the framework regions crucial for the definition of the canonical class of the murine antibody CDR loops; (e) residues that are adjacent to a CDR; (g) residues capable of interacting with the antigen; (h) residues capable of interacting with the CDR; and (i) contact residues between the VH domain and the VL domain.

The present invention encompasses formaultions that comprise antibodies that immunospecifically bind to a RSV F antigen, said antibodies comprising the amino acid sequence of the variable heavy domain and/or variable light domain or an antigen-binding fragment thereof of AFFF, P12f2, P12f4, P11d4, Ale9, A12a6, A13c4, A17d4, A4B4, A8C7, 1X-493L1FR, H3-3F4, M3H9, Y10H6, DG, AFFF(1), 6H8, L1-7E5, L2-15B10, A13a11, A1h5, A4B4(1), A4B4L1FR-S28R (motavizumab), A4B4-F52S, A17d4(1), A3e2, A14a4, A16b4, A17b5, A17f5, or A17h4 with mutations (*e.g.*, one or more amino acid substitutions) in the framework regions. In certain embodiments, antibodies that immunospecifically bind to a RSV antigen comprise the amino acid sequence of the variable heavy domain and/or variable light domain or an antigen-binding fragment thereof of AFFF, P12f2, P12f4, P11d4, Ale9, A12a6, A13c4, A17d4, A4B4, A8C7, 1X-493L1FR, H3-3F4, M3H9, Y10H6, DG, AFFF(1), 6H8, L1-7E5, L2-15B10, A13a11, A1h5, A4B4(1), A4B4L1FR-S28R (motavizumab), A4B4-F52S,

A17d4(1), A3e2, A14a4, A16b4, A17b5, A17f5, or A17h4 with one or more amino acid residue substitutions in the framework regions of the VH and/or VL domains.

[00207] The present invention also encompasses formulations that comprise antibodies which immunospecifically bind to one or more RSV F antigens, said antibodies comprising the amino acid sequence of A4B4L1FR-S28R (motavizumab) with mutations (e.g., one or more amino acid substitutions) in the framework regions. In certain embodiments, antibodies which immunospecifically bind to one or more RSV F antigens comprise the amino acid sequence of A4B4L1FR-S28R (motavizumab) with one or more amino acid residue substitutions in the framework regions of the VH and/or VL domains. In a specific embodiment, antibodies which immunospecifically bind to one or more RSV F antigens comprise the framework regions depicted in Figure 2 or Figure 13.

[00208] The present invention also provides antibodies that immunospecifically bind to a RSV antigen, said antibodies comprising the amino acid sequence of the variable heavy domain and/or variable light domain of an antibody in Table 2 (*i.e.*, AFFF, P12f2, P12f4, P11d4, Ale9, A12a6, A13c4, A17d4, A4B4, A8C7, 1X-493L1FR, H3-3F4, M3H9, Y10H6, DG, AFFF(1), 6H8, L1-7E5, L2-15B10, A13a11, A1h5, A4B4(1), A4B4L1FR-S28R (motavizumab), A4B4-F52S, A17d4(1), A3e2, A14a4, A16b4, A17b5, A17f5, or A17h4) with mutations (*e.g.*, one or more amino acid residue substitutions) in the hypervariable and framework regions. Preferably, the amino acid substitutions in the hypervariable and framework regions improve binding of the antibody to a RSV antigen.

[00209] The present invention also provides antibodies which immunospecifically bind to one or more RSV F antigens, said antibodies comprising the amino acid sequence of A4B4L1FR-S28R (motavizumab) with mutations (*e.g.*, one or more amino acid residue substitutions) in the variable and framework regions.

[00210] The present invention also provides antibodies that immunospecifically bind to a RSV antigen (*e.g.*, RSV F antigen) which comprise constant regions known to those of skill in the art. Preferably, the constant regions of an antibody of the invention are human. In a specific embodiment, an antibody of the invention comprises the constant regions of A4B4L1FR-S28R (motavizumab).

[00211] The present invention also provides fusion proteins comprising an antibody that immunospecifically binds to a RSV antigen and a heterologous polypeptide. Preferably, the

heterologous polypeptide that the antibody is fused to is useful for targeting the antibody to respiratory epithelial cells.

[00212] The present invention also encompasses formulations that comprise panels of antibodies that immunospecifically bind to a RSV antigen. In specific embodiments, the invention provides panels of antibodies having different association rate constants different dissociation rate constants, different affinities for a RSV antigen, and/or different specificities for a RSV antigen. The invention provides panels of at least 10, preferably at least 25, at least 50, at least 75, at least 100, at least 125, at least 150, at least 175, at least 200, at least 250, at least 300, at least 350, at least 400, at least 450, at least 500, at least 550, at least 600, at least 650, at least 700, at least 750, at least 800, at least 850, at least 900, at least 950, or at least 1000 antibodies. Panels of antibodies can be used, for example, in 96 well plates for assays such as ELISAs.

[00213] The present invention further provides one or more antibodies for use in the prevention, treatment, and/or amelioration of an upper and/or lower respiratory tract RSV infection, otitis media, or a symptom or respiratory condition relating thereto (including, but not limited to, asthma, wheezing, RAD, or a combination thereof). In a specific embodiment, a formulation for use in the prevention, treatment, and/or amelioration of an upper and/or lower respiratory tract RSV infection, otitis media, or a symptom or respiratory condition relating thereto (including, but not limited to, asthma, wheezing, RAD, or a combination thereof) comprises AFFF, P12f2, P12f4, P11d4, Ale9, A12a6, A13c4, A17d4, A4B4, A8C7, 1X-493L1FR, H3-3F4, M3H9, Y10H6, DG, AFFF(1), 6H8, L1-7E5, L2-15B10, A13a11, A1h5, A4B4(1), A4B4L1FR-S28R (motavizumab), A4B4-F52S, A17d4(1), A3e2, A14a4, A16b4, A17b5, A17f5, and/or A17h4. In another specific embodiment, a formulation for use in the prevention, treatment, and/or amelioration of an upper and/or lower respiratory tract RSV infection, otitis media, or a symptom or respiratory condition relating thereto (including, but not limited to, asthma, wheezing, RAD, or a combination thereof) comprises an antigen-binding fragment of AFFF, P12f2, P12f4, P11d4, Ale9, A12a6, A13c4, A17d4, A4B4, A8C7, 1X-493L1FR, H3-3F4, M3H9, Y10H6, DG, AFFF(1), 6H8, L1-7E5, L2-15B10, A13a11, A1h5, A4B4(1), A4B4L1FR-S28R (motavizumab), or A4B4-F52S, A17d4(1), A3e2, A14a4, A16b4, A17b5, A17f5, or A17h4.

[00214] In another embodiment, a formulation for use in the prevention, treatment, and/or amelioration of an upper and/or lower respiratory tract RSV infection, otitis media, or a symptom or respiratory condition relating thereto (including, but not limited to, asthma,

wheezing, RAD, or a combination thereof) comprises one or more antibodies comprising one or more VH domains having an amino acid sequence of any one of the VH domains listed in Table 2. In another embodiment, a formulation for use in the prevention, treatment, and/or amelioration of an upper and/or lower respiratory tract RSV infection, otitis media, or a symptom or respiratory condition relating thereto (including, but not limited to, asthma, wheezing, RAD, or a combination thereof) comprises one or more antibodies comprising one or more VH CDR1s having an amino acid sequence of any one of the VH CDR1s listed in Table 2 and/or Table 3A. In another embodiment, a formulation for use in the prevention, treatment, and/or amelioration of an upper and/or lower respiratory tract RSV infection, otitis media, or a symptom or respiratory condition relating thereto (including, but not limited to, asthma, wheezing, RAD, or a combination thereof) comprises one or more antibodies comprising one or more VH CDR2s having an amino acid sequence of any one of the VH CDR2s listed in Table 2 and/or Table 3B. In a preferred embodiment, a formulation for use in the prevention, treatment, and/or amelioration of an upper and/or lower respiratory tract RSV infection, otitis media, or a symptom or respiratory condition relating thereto (including, but not limited to, asthma, wheezing, RAD, or a combination thereof) comprises one or more antibodies comprising one or more VH CDR3s having an amino acid sequence of any one of the VH CDR3s listed in Table 2 and/or Table 3C.

In another embodiment, a formulation for use in the prevention, treatment, and/or [00215] amelioration of an upper and/or lower respiratory tract RSV infection, otitis media, or a symptom or respiratory condition relating thereto (including, but not limited to, asthma, wheezing, RAD, or a combination thereof) comprises one or more antibodies comprising one or more VL domains having an amino acid sequence of any one of the VL domains listed in Table 2. In another embodiment, a formulation for use in the prevention, treatment, and/or amelioration of an upper and/or lower respiratory tract RSV infection, otitis media, or a symptom or respiratory condition relating thereto (including, but not limited to, asthma, wheezing, RAD, or a combination thereof) comprises one or more antibodies comprising one or more VL CDR1s having an amino acid sequence of any one of the VL CDR1s listed in Table 2 or Table 3D. In another embodiment, a formulation for use in the prevention, treatment, and/or amelioration of an upper and/or lower respiratory tract RSV infection, otitis media, or a symptom or respiratory condition relating thereto (including, but not limited to, asthma, wheezing, RAD, or a combination thereof) comprises one or more antibodies comprising one or more VL CDR2s having an amino acid sequence of any one of the VL CDR2s listed in Table 2 and/or Table 3E. In a preferred embodiment, a formulation for use in the prevention, treatment,

and/or amelioration of an upper and/or lower respiratory tract RSV infection, otitis media, or a symptom or respiratory condition relating thereto (including, but not limited to, asthma, wheezing, RAD, or a combination thereof) comprises one or more antibodies comprising one or more VL CDR3s having an amino acid sequence of any one of the VL CDR3s listed in Table 2 and/or Table 3F.

In another embodiment, a formulation for use in the prevention, treatment, and/or [00216] amelioration of an upper and/or lower respiratory tract RSV infection, otitis media, or a symptom or respiratory condition relating thereto (including, but not limited to, asthma, wheezing, RAD, or a combination thereof) comprises one or more antibodies comprising one or more VH domains having an amino acid sequence of any one of the VH domains listed in Table 2 and one or more VL domains having an amino acid sequence of any one of the VL domains listed in Table 2. In another embodiment, a formulation for use in the prevention, treatment, and/or amelioration of an upper and/or lower respiratory tract RSV infection, otitis media, or a symptom or respiratory condition relating thereto (including, but not limited to, asthma, wheezing, RAD, or a combination thereof) comprises one or more antibodies comprising one or more VH CDR1s having an amino acid sequence of any one of the VH CDR1s listed in Table 2 and/or Table 3A and one or more VL CDR1s having an amino acid sequence of any one of the VL CDR1s listed in Table 2 and/or Table 3D. In another embodiment, a formulation for use in the prevention, treatment, and/or amelioration of an upper and/or lower respiratory tract RSV infection, otitis media, or a symptom or respiratory condition relating thereto (including, but not limited to, asthma, wheezing, RAD, or a combination thereof) comprises one or more antibodies comprising one or more VH CDR1s having an amino acid sequence of any one of the VH CDR1s listed in Table 2 and/or Table 3A and one or more VL CDR2s having an amino acid sequence of any one of the VL CDR2s listed in Table 2 and/or Table 3E. In another embodiment, a formulation for use in the prevention, treatment, and/or amelioration of an upper and/or lower respiratory tract RSV infection, otitis media, or a symptom or respiratory condition relating thereto (including, but not limited to, asthma, wheezing, RAD, or a combination thereof) comprises one or more antibodies comprising one or more VH CDR1s having an amino acid sequence of any one of the VH CDR1s listed in Table 2 and/or Table 3A and one or more VL CDR3s having an amino acid sequence of any one of the VL CDR3s listed in Table 2 and/or Table 3F.

[00217] In another embodiment, a formulation for use in the prevention, treatment, and/or amelioration of an upper and/or lower respiratory tract RSV infection, otitis media, or a

symptom or respiratory condition relating thereto (including, but not limited to, asthma, wheezing, RAD, or a combination thereof) comprises one or more antibodies comprising one or more VH CDR2s having an amino acid sequence of any one of the VH CDR2s listed in Table 2 and/or Table 3B and one or more VL CDR1s having an amino acid sequence of any one of the VL CDR1s listed in Table 2 and/or Table 3D. In another embodiment, a formulation for use in the prevention, treatment, and/or amelioration of an upper and/or lower respiratory tract RSV infection, otitis media, or a symptom or respiratory condition relating thereto (including, but not limited to, asthma, wheezing, RAD, or a combination thereof) comprises one or more antibodies comprising one or more VH CDR2s having an amino acid sequence of any one of the VH CDR2s listed in Table 2 and/or Table 3B and one or more VL CDR2s having an amino acid sequence of any one of the VL CDR2s listed in Table 2 and/or Table 3E. In another embodiment, a formulation of the present invention comprises one or more antibodies comprising one or more VH CDR2s having an amino acid sequence of any one of the VH CDR2s listed in Table 2 and/or Table 3B and one or more VL CDR3s having an amino acid sequence of any one of the VH CDR2s listed in Table 2 and/or Table 3B and one or more VL CDR3s having an amino acid sequence of any one of the VH CDR2s listed in Table 2 and/or Table 3B.

In another embodiment, a formulation for use in the prevention, treatment, and/or [00218] amelioration of an upper and/or lower respiratory tract RSV infection, otitis media, or a symptom or respiratory condition relating thereto (including, but not limited to, asthma, wheezing, RAD, or a combination thereof) comprises one or more antibodies comprising one or more VH CDR3s having an amino acid sequence of any one of the VH CDR3s listed in Table 2 and/or Table 3C and one or more VL CDR1s having an amino acid sequence of any one of the VL CDR1s listed in Table 2 and/or Table 3D. In another embodiment, a formulation for use in the prevention, treatment, and/or amelioration of an upper and/or lower respiratory tract RSV infection, otitis media, or a symptom or respiratory condition relating thereto (including, but not limited to, asthma, wheezing, RAD, or a combination thereof) comprises one or more antibodies comprising one or more VH CDR3s having an amino acid sequence of any one of the VH CDR3s listed in Table 2 and/or Table 3C and one or more VL CDR2s having an amino acid sequence of any one of the VL CDR2s listed in Table 2 and/or Table 3E. In a preferred embodiment, a formulation for use in the prevention, treatment, and/or amelioration of an upper and/or lower respiratory tract RSV infection, otitis media, or a symptom or respiratory condition relating thereto (including, but not limited to, asthma, wheezing, RAD, or a combination thereof) comprises one or more antibodies comprising one or more VH CDR3s having an amino acid sequence of any one of the VH CDR3s listed in Table 2 and/or Table 3C and one or more VL CDR3s having an amino acid sequence of any one of the VL CDR3s listed in Table 2 and/or

Table 3F. In a preferred embodiment, a formulation for use in the prevention, treatment, and/or amelioration of an upper and/or lower respiratory tract RSV infection, otitis media, or a symptom or respiratory condition relating thereto (including, but not limited to, asthma, wheezing, RAD, or a combination thereof) comprises A4B4L1FR-S28R (motavizumab) or an antigen-binding fragment thereof. In yet another embodiment, a formulation of the present invention comprises one or more fusion proteins of the invention.

[00219] As discussed in more detail below, a formulation of the invention may be used either alone or in combination with other compositions. The antibodies may further be recombinantly fused to a heterologous polypeptide at the N – or C-terminus or chemically conjugated (including covalently and non-covalently conjugations) to polypeptides or other compositions. For example, antibodies of the present invention may be recombinantly fused or conjugated to molecules useful as labels in detection assays and effector molecules such as heterologous polypeptides, drugs, radionucleotides, or toxins. See, *e.g.*, PCT publications WO 92/08495; WO 91/14438; WO 89/12624; U.S. Patent No. 5,314,995; and EP 396,387.

[00220] Antibodies of the present invention may be used, for example, to purify, detect, and target RSV antigens, in both *in vitro* and *in vivo* diagnostic and therapeutic methods. For example, the antibodies have use in immunoassays for qualitatively and quantitatively measuring levels of the RSV in biological samples such as sputum. See, *e.g.*, Harlow et al., Antibodies: A Laboratory Manual, (Cold Spring Harbor Laboratory Press, 2nd ed. 1988) (incorporated by reference herein in its entirety).

[00221] The invention provides an antibody comprising a Fab fragment, which immunospecifically binds to an RSV antigen (e.g., the F protein epitope NSELLSLINDMPITNDQKKLMSNN (SEQ ID NO: 337)), wherein the Tm of the Fab fragment is at least about 87 °C, and wherein said antibody is not any one of palivizumab, AFFF, P12f2, P12f4, P11d4, Ale9, A12a6, A13c4, A17d4, A4B4, A8c7, 1X-493L1FR, H3-3F4, M3H9, Y10H6, DG, AFFF(1), 6H8, L1-7E5, L2-15B10, A13a11, A1h5, A4B4(1), A4B4L1FR-S28R (motavizumab), A4B4-F52S, A17d4(1), A3e2, A14a4, A16b4, A17b5, A17f5, and A17h4. In a specific embodiment, the Fab in such an antibody is different from the Fab of palivizumab. In another embodiment, such an antibody comprises a VH or VL domain that is different from the VH or VL domain of palivizumab. In preferred embodiment, the Tm of the Fab fragment is at least about 90 °C or at least about 93 °C. In another preferred embodiment, the pI of the antibody is between about 8.5 to 9.5 or between about 9.0 to 9.5.

[00222] In another specific embodiment, the antibody comprises a VH domain of the antibody A4B4L1FR-S28R (SEQ ID NO:48). In still another embodiment, the antibody comprises a VL domain of the antibody A4B4L1FR-S28R (SEQ ID NO:11). In still another embodiment, the Fab of the antibody is the Fab of antibody A4B4L1FR-S28R, preferably having one or more amino acid modifications in this constant domain.

[00223] The invention also provides an antibody formulation comprising the above described antibody, said formulation having a viscosity of less than 10.00 cP or less than 5.00cP at any temperature in the range of 1 to 26 °C, or in the range of 5 to 25 °C, or in the range of 10 to 25 °C.

[00224] The invention also provides an antibody formulation comprising the above described antibody, said formulation having an aggregration rate of less than about 5%, 10%, or 15% per day at any temperature in the range of 38 to 42 °C.

[00225] The above described antibodies can be generated by a method described in U.S. Provisional Patent Application No.: 60/696,113, by Christian B. Allan, filed on July 1, 2005, which is incorporated by reference herein in its entirety. In a specific embodiment, such an antibody is generated by a method comprising screening a plurality of candidate antibody domains (e.g., Fab, Fc and Fv) that have high binding affinity to a target (e.g., RSV antigen) for their solubility and thermal stability. Any method known in the art for screening protein domains for their solubility and thermal stability can be used. One or more antibody domains having high solubility and/or thermal stability are then selected and used for constructing the full antibodies by combining them with the appropriate domain(s) to generate a full antibody. In one embodiment, one or more candidate Fab domains that have a Tm value higher than at least 87°C, 90°C, 95°C, 100°C, 105°C, 110°C, 115°C, or 120°C are selected for construction of the full antibody. In another embodiment, one or more candidate domains that have a pI value higher than about 6.5, 7.0, 7.5, 8.0, 8.5 or 9.0 are selected for construction of the full antibody. In a specific embodiment, the plurality of candidate Fab domains comprises Fab domains containing one or more amino acid residue substitutions to the Fab domain of the following antibodies palivizumab, AFFF, P12f2, P12f4, P11d4, Ale9, A12a6, A13c4, A17d4, A4B4, A8c7, 1X-493L1FR, H3-3F4, M3H9, Y10H6, DG, AFFF(1), 6H8, L1-7E5, L2-15B10, A13a11, A1h5, A4B4(1), A4B4L1FR-S28R (motavizumab), A4B4-F52S, A17d4(1), A3e2, A14a4, A16b4, A17b5, A17f5, and/or A17h4.

[00226] A plurality of antigen binding domains (e.g., Fab, scFv, etc.) that bind a RSV antigen with an affinity above a chosen threshold may be obtained, e.g., by affinity screening of a phage display library. One or more metrics characterizing the antigen binding domains' formulation properties are then evaluated for each of the antigen binding domains. The plurality of antigen binding domains are ranked according to the one or more metrics. In one embodiment, the plurality of antigen binding domains are ranked according to their Tm values, and one or more antigen binding domains are selected from the top of the ranked list. In another embodiment, the plurality of antigen binding domains are ranked according to their pI values, and one or more antigen binding domains are selected from the top of the ranked list. In still another embodiment, the plurality of antigen binding domains are ranked according to a combined Tm and pI rank, and one or more antigen binding domains are selected from the top of the ranked list. The selected antigen binding domains are then used for construction of the full anti-RSV antibody molecule (e.g., antibodies, diabodies, etc.).

[00227] In another embodiment, a plurality of antibody constant region domains (e.g., Fc, CH2, CH3, etc) is screened for solubility and thermal stability. In one embodiment, one or more candidate antibody constant region domains that have a Tm value higher than at least 50°C, 55°C, 60°C, 65°C, 70°C, 75°C, 80°C, 85°C, 90°C, 95°C, 100°C, 105°C, 110°C, 115°C, or 120°C are selected for construction of the full antibody. In another embodiment, one or more candidate antibody constant region domains that have a pI value higher than about 6.5, 7.0, 7.5, 8.0, 8.5 or 9.0 are selected for construction of the full antibody (e.g., antibody, Fc-fusion protein, etc.).

[00228] Such an antibody can also be generated by a method for engineering a protein for preferred formulation characteristics and/or properties including but not limited to, Tm, pI, solubility, stability. In one embodiment, the method comprises engineering one or more domains to improve the antibody's formulation characteristics. In a preferred embodiment, the engineered domain exhibits improved formulation characteristics without reducing significantly the antibody's pharmacological characteristics including but not limited to, the antibody's binding specificity, binding affinity and/or avidity to its target, or the antibody's Fc effector functions, *e.g.*, Fc-receptor (FcR) binding, antibody dependent cellular cytotoxicity (ADCC), complement dependent cytotoxicity (CDC), and/or serum half life. In a more preferred embodiment, the engineered domain exhibits improved formulation characteristics without substantially affecting the antibody's pharmacological characteristics.

In a preferred embodiment, a domain is engineered by substituting one or more amino acid residues in the domain such that the stability of the domain is increased. In one embodiment, a domain is engineered such that its Tm value is increased. In one embodiment, a domain is engineered such that it has a Tm greater than a predetermined threshold value. In some preferred embodiments, the predetermined Tm threshold value is at least 50°C, 55°C, 60°C, 65°C, 70°C, 75°C, 80°C, 85°C, 90°C, 95°C, 100°C, 105°C, 110°C, 115°C, or 120°C. In a specific embodiment, such an engineered Fab domain is generated by substituting one or more amino acid residues in the Fab domain of palivizumab, AFFF, P12f2, P12f4, P11d4, Ale9, A12a6, A13c4, A17d4, A4B4, A8c7, 1X-493L1FR, H3-3F4, M3H9, Y10H6, DG, AFFF(1), 6H8, L1-7E5, L2-15B10, A13a11, A1h5, A4B4(1), A4B4L1FR-S28R (motavizumab), A4B4-F52S, A17d4(1), A3e2, A14a4, A16b4, A17b5, A17f5, or A17h4.

[00230] In another preferred embodiment, a domain is engineered by substituting one or more amino acid residues in the domain such that the solubility of the domain is increased. In one embodiment, a domain is engineered such that its pI value is increased. In one embodiment, a domain is engineered such that it has a pI greater than a predetermined threshold value. In some preferred embodiments, the predetermined pI threshold value is about 6.5, 7.0, 7.5, 8.0, 8.5, or 9.0.

[00231] In one embodiment, the antigen binding (e.g., Fab) and/or constant region (e.g., Fc) domains are engineered to improve the protein's formulation characteristics, e.g., Tm, pI, or stability. In preferred embodiments, the engineered antibody exhibits improved formulation characteristics without reducing significantly the antibody's pharmacological characteristics, e.g., the antibody's binding specificity, binding affinity and/or avidity to its target, or the antibody's Fc effector functions, e.g., Fc-receptor (FcR) binding, antibody dependent cellular cytotoxicity (ADCC), complement dependent cytotoxicity (CDC), and/or serum half life. In another embodiment, the engineered antibody exhibits improved formulation characteristics and improved pharmacological characteristics, e.g., the antibody's binding specificity, binding affinity and/or avidity to its target, or the antibody's Fc effector functions, e.g., FcR binding, ADCC, CDC, and/or serum half life.

[00232] The solubility of a protein may be optimized by altering the number and location of ionizable residues in the protein to adjust the pI. For example the pI of a polypeptide can be manipulated by making the appropriate amino acid substitutions (e.g., by substituting a charged amino acid such as a lysine, for an uncharged residue such as alanine). Without wishing to be bound by any particular theory, amino acid substitutions of a protein that result in changes of the

pI of said protein may improve solubility and/or the stability of the protein. One skilled in the art would be able to determine amino acid substitutions that is most appropriate for a particular protein to achieve a desired pI. The pI of a protein may be determined by a variety of methods including but not limited to isoelectric focusing. It can also be estimated using any one of the various computer algorithms (see for example Bjellqvist *et al.*, 1993, *Electrophoresis* 14:1023, which is incorporated herein by reference in its entirety).

[00233] In one embodiment, the pI of an engineered antibody binding domain is between pH 6.2 and pH 10.0. In one embodiment, substitutions resulting in alterations in the pI of the antigen binding domain will not significantly diminish its binding affinity for an antigen. In one embodiment, the pI of an engineered antibody constant region domain is between pH 6.2 and pH 10.0. In still another embodiment, substitutions resulting in alterations in the pI of the constant region domain will not significantly diminish its effector binding and/or function. It is also contemplated that substitutions resulting in alterations in the pI in an antibody domain may be selected such that both the pI and other pharmacological characteristics of the antibody domain, e.g., the antibody's binding specificity, binding affinity and/or avidity to its target, or the antibody's Fc effector functions are improved. The inventors have found that certain modifications of the hinge region do not change the pI and Tm of the antibody significantly. Thus, in one embodiment, the invention provides a method for engineering an antibody to improve the antibody's biological activity without reducing the antibody's formulation properties.

In one embodiment, the modifications of an antibody domain as described herein may be combined with known modifications of the Fc domain such as those disclosed in Duncan et al, 1988, *Nature* 332:563-564; Lund *et al.*, 1991, *J. Immunol* 147:2657-2662; Lund et al, 1992, *Mol Immunol* 29:53-59; Alegre et al, 1994, *Transplantation* 57:1537-1543; Hutchins *et al.*, 1995, *Proc Natl. Acad Sci U S A* 92:11980-11984; Jefferis et al, 1995, *Immunol Lett.* 44:111-117; Lund *et al.*, 1995, *Faseb J* 9:115-119; Jefferis et al, 1996, *Immunol Lett* 54:101-104; Lund et al, 1996, *Immunol* 157:4963-4969; Armour *et al.*, 1999, *Eur J Immunol* 29:2613-2624; Idusogie et al, 2000, *J Immunol* 164:4178-4184; Reddy et al, 2000, *J Immunol* 164:1925-1933; Xu *et al.*, 2000, *Cell Immunol* 200:16-26; Idusogie et al, 2001, *J Immunol* 166:2571-2575; Shields *et al.*, 2001, *J Biol Chem* 276:6591-6604; Jefferis et al, 2002, *Immunol Lett* 82:57-65; Presta *et al.*, 2002, *Biochem Soc Trans* 30:487-490); U.S. Pat. Nos. 5,624,821; 5,885,573; 6,194,551; U.S. Patent Application Nos. 60/601,634 and 60/608,852; PCT Publication Nos. WO 00/42072 and WO 99/58572; each of which is incorporated herein by reference in its entirety.

[00235] In one embodiment, the antibodies may be engineered to include modifications in the Fc region, typically to alter one or more functional properties of the antibody, such as serum half-life, complement fixation, Fc receptor binding, and/or antigen-dependent cellular cytotoxicity, without reducing the antibodies' pI and Tm. Furthermore, an antibody may be chemically modified (e.g., one or more chemical moieties can be attached to the antibody) or be modified to alter its glycosylation, again to alter one or more functional properties of the antibody.

[00236] In one embodiment, the amino acid sequence of the Fc region is modified by deleting, adding and/or substituting at least amino acid residue to alter one or more of the functional properties of the antibody described above. This approach is described further in Duncan et al. 1988, Nature 332:563-564; Lund et al., 1991, J. Immunol 147:2657-2662; Lund et al, 1992, Mol Immunol 29:53-59; Alegre et al, 1994, Transplantation 57:1537-1543; Hutchins et al., 1995, Proc Natl. Acad Sci USA 92:11980-11984; Jefferis et al, 1995, Immunol Lett. 44:111-117; Lund et al., 1995, Faseb J 9:115-119; Jefferis et al, 1996, Immunol Lett 54:101-104; Lund et al, 1996, J Immunol 157:4963-4969; Armour et al., 1999, Eur J Immunol 29:2613-2624; Idusogie et al, 2000, J Immunol 164:4178-4184; Reddy et al, 2000, J Immunol 164:1925-1933; Xu et al., 2000, Cell Immunol 200:16-26; Idusogie et al, 2001, J Immunol 166:2571-2575; Shields et al., 2001, J Biol Chem 276:6591-6604; Jefferis et al, 2002, Immunol Lett 82:57-65; Presta et al., 2002, Biochem Soc Trans 30:487-490); U.S. Patent Nos. 5,624,821; 5,885,573; 5.677.425; 6.165.745; 6.277.375; 5.869.046; 6.121.022; 5.624.821; 5.648.260; 6.194.551; 6,737,056 U.S. Patent Application Nos. 10/370,749 and PCT Publications WO 94/2935; WO 99/58572; WO 00/42072; WO 04/029207, each of which is incorporated herein by reference in its entirety.

[00237] In still another embodiment, the glycosylation of antibodies is modified. For example, an aglycoslated antibody can be made (*i.e.*, the antibody lacks glycosylation). Glycosylation can be altered to, for example, increase the affinity of the antibody for a target antigen. Such carbohydrate modifications can be accomplished by, for example, altering one or more sites of glycosylation within the antibody sequence. For example, one or more amino acid substitutions can be made that result in elimination of one or more variable region framework glycosylation sites to thereby eliminate glycosylation at that site. Such aglycosylation may increase the affinity of the antibody for antigen. Such an approach is described in further detail in U.S. Patent Nos. 5,714,350 and 6,350,861, each of which is incorporated herein by reference in its entirety.

[00238] Additionally or alternatively, an antibody can be made that has an altered type of glycosylation, such as a hypofucosylated antibody having reduced amounts of fucosyl residues or an antibody having increased bisecting GlcNAc structures. Such altered glycosylation patterns have been demonstrated to increase the ADCC ability of antibodies. Such carbohydrate modifications can be accomplished by, for example, expressing the antibody in a host cell with altered glycosylation machinery. Cells with altered glycosylation machinery have been described in the art and can be used as host cells in which to express recombinant antibodies of the invention to thereby produce an antibody with altered glycosylation. See, for example, Shields, R.L. et al. (2002) J. Biol. Chem. 277:26733-26740; Umana et al. (1999) Nat. Biotech. 17:176-1, as well as, European Patent No: EP 1,176,195; PCT Publications WO 03/035835; WO 99/54342, each of which is incorporated herein by reference in its entirety.

In another embodiment, the antibodies may be engineered to include [00239] modifications in the antigen binding domain to alter the formulation characteristics of the antibody, without reducing the binding characteristics. One skilled in the art will understand that amino acid substitutions and other modifications of an antibody may alter its antigen binding characteristics (examples of binding characteristics include but are not limited to, binding specificity, equilibrium dissociation constant (K_D), dissociation and association rates (K_{off} and K_{on} respectively), binding affinity and/or avidity) and that certain alterations are more or less desirable. For example a modification that preserves or enhances antigen binding would be more preferable then one that diminished or altered antigen binding. The binding characteristics of an antibody for a target antigen, may be determined by a variety of methods including but not limited it, equilibrium methods (e.g., enzyme-linked immunoabsorbent assay (ELISA) or radioimmunoassay (RIA)), or kinetics (e.g., BIACORE® analysis; see Example 2), for example. Other commonly used methods to examine the binding characteristics of antibodies are described in Using Antibodies: A Laboratory Manual, Cold Spring Harbor Laboratory Press, NY, Harrow et al., 1999 and Antibodies: A Laboratory Manual, Cold Spring Harbor Laboratory Press, NY; Harlow et al., 1989.

[00240] It is well known in the art that the equilibrium dissociation constant (K_D) is defined as k_{off}/k_{on} . It is generally understood that an antibody with a low K_D is preferable to an antibody with a high K_D . However, in some instances the value of the k_{on} or k_{off} may be more relevant than the value of the K_D . One skilled in the art can determine which kinetic parameter is most important for a given antigen binding domain and application. In a preferred embodiment, the method of the invention will result in antigen binding domains with improved

formulation characteristics and one or more antigen binding characteristics (*e.g.*, binding specificity, K_D, K_{off}, K_{on}, binding affinity and/or avidity) that are improved by at least 2%, or by at least 5%, or by at least 10%, or by at least 20 %, or by at least 30%, or by at least 40%, or by at least 50%, or by at least 60%, or by at least 70%, or by at least 80% when compared to kinetic parameters of the antigen binding domain without said modification.

In another embodiment, the method of the invention will result in modified antigen binding domains that have improved formulation characteristics, but do not have substantially diminished antigen binding. For example, the method of the invention will generate antigen binding domains that exhibit improved formulation characteristics, but preferably have no reduction in any antigen binding characteristic (*e.g.*, binding specificity, K_D, K_{off}, K_{on}, binding affinity and/or avidity), or have one or more antigen binding characteristics that are reduced by less than 1%, or by less than 5%, or by less than 10%, or by less than 20 %, or by less than 30%, or by less than 40%, or by less than 50%, or by less than 60%, or by less than 60%, or by less than 50% when compared to antigen binding of the antibody without said substitution.

[00242] In one embodiment, selected or engineered antigen binding and antibody constant domains are then used to construct a full anti-RSV antibody using methods known in the art. Such antibodies can then be submitted to formulation development to determine the optimal formulations.

5.3.4 Antibodies that Immunospecifically Bind to Human Metapneumovirus (hMPV)

[00243] The formulations of the present invention comprise an isolated antibody that specifically binds to an antigen of human metapheumovirus (hMPV) and compositions comprising this antibody. The term "anti-hMPV-antigen antibody" refers to an antibody or antibody fragment thereof that binds immunospecifically to a hMPV antigen. A hMPV antigen refers to a hMPV polypeptide or fragment thereof such as of hMPV nucleoprotein, hMPV phosphoprotein, hMPV matrix protein, hMPV small hydrophobic protein, hMPV RNA-dependent hMPV polymerase, hMPV F protein, and hMPV G protein. A hMPV antigen also refers to a polypeptide that has a similar amino acid sequence compared to a hMPV polypeptide or fragment thereof such as of hMPV nucleoprotein, hMPV phosphoprotein, hMPV matrix protein, hMPV small hydrophobic protein, hMPV RNA-dependent hMPV polymerase, hMPV F protein, and hMPV G protein.

[00244] The anti-hMPV-antigen antibodies used in this invention can be monoclonal antibodies, human antibodies, humanized antibodies or chimeric antibodies. In some preferred embodiments, the anti-hMPV antibody of the invention is the antibody disclosed in U.S. Patent Application No. 10/628,088, filed July 25, 2003 and published May 20, 2004, as U.S. Pat. Pub. No. US 2004/0096451 A1.

[00245] The anti-hMPV-antigen antibodies of this section can be made, formulated, administered, used therapeutically or used prophylactically as described in U.S. Patent Application No. 10/628,088, filed July 25, 2003 and published May 20, 2004, as U.S. Pat. Pub. No. US 2004/0096451 A1, the contents of which are hereby incorporated by reference in their entirety.

5.3.5 Antibodies that Immunospecifically Bind to Integrin $\alpha_v \beta_3$

[00246] The formulations of the present invention also comprise an isolated antibody that specifically binds to integrin $\alpha_v \beta_3$ and compositions comprising this antibody. The antibodies can be monoclonal antibodies, human antibodies, humanized antibodies or chimeric antibodies. In some preferred embodiments, the anti- integrin $\alpha_v \beta_3$ antibody of the invention is MEDI-522 (Vitaxin®). Vitaxin® and compositions or formulations comprising Vitaxin® are disclosed, *e.g.*, in International Publication Nos. WO 98/33919, WO 00/78815, and WO 02/070007; U.S. application Serial No. 09/339,222; U.S. Patent Application No. 10/091,236, filed March 4, 2002 and published November 12, 2002, as U.S. Pat. Pub. No. US 2002/0168360, each of which is incorporated herein by reference in its entirety.

[00247] In further embodiments, the antibody that immunospecifically binds to integrin $\alpha_{\nu}\beta_{3}$ is not Vitaxin® or an antigen-binding fragment of Vitaxin®. Examples of known antibodies that immunospecifically bind to integrin $\alpha_{\nu}\beta_{3}$ include, but are not limited to, 11D2 (Searle), the murine monoclonal LM609 (Scripps, International Publication Nos. WO 89/05155 and U.S. Patent No. 5,753,230, which is incorporated herein by reference in its entirety), International Publication Nos WO 98/33919 and WO 00/78815, each of which is incorporated herein by reference in its entirety), 17661-37E and 17661-37E 1-5 (USBiological), MON 2032 and 2033 (CalTag), ab7166 (BV3) and ab 7167 (BV4) (Abcam), and WOW-1 (Kiosses et al., Nature Cell Biology, 3:316-320).

[00248] $\alpha_v \beta_{3}$, an integrin has been found on new blood vessels as well as surface of many solid tumors, activated macrophages, monocytes, and osteoclasts. As the such, the anti-integrin

 $\alpha_{\nu}\beta_{3}$ antibodies of this section can be used, for example, as an investigational antibody, or in the prevention or treatment of several destructive diseases.

[00249] The anti- integrin $\alpha_v \beta_3$ antibodies of this section can be made, formulated, administered, used therapeutically or used prophylactically as described in U.S. Patent Application No. 10/091,236, filed March 4, 2002 and published November 12, 2002, as U.S. Pat. Pub. No. US 2002/0168360; U.S. Patent Application No. 10/769,712, filed January 30, 2004; U.S. Patent Application No. 10/769,720, filed January 30, 2004 and published September 9, 2004, as U.S. Pat. Pub. No. US 2004/0176272; U.S. Patent Application No. 10/379,145, filed March 4, 2003; U.S. Patent Application No. 10/379,189, filed March 4, 2003 and published as U.S. Pat. Pub. No. US 2004/0001835; PCT Application No. PCT/US04/02701, filed January 30, 2004; International Application Publication No.: WO 00/78815 A1, entitled "Anti- $\alpha_v \beta_3$ recombinant human antibodies, nucleic acids encoding same and methods", by Huse et al.; and International Application Publication No.: WO 98/33919 A1, entitled "Anti-alpha-V-veta-3 recombinant humanized antibodies, nucleic acids encoding same and methods of use", by Huse et al.; International Publication No. WO 89/05155, the contents of which are hereby incorporated by reference in their entirety.

5.3.6 Antibodies that Immunospecifically Bind to CD2

The formulations of the present invention comprise an isolated antibody that [00250] immunospecifically binds to CD2 and compositions comprising this antibody. The antibodies can be monoclonal antibodies, human antibodies, humanized antibodies or chimeric antibodies. In some preferred embodiments, the anti-CD2 antibody of the invention is siplizumab (MEDI-507). Siplizumab can selectively binds to cells expressing the CD2 antigen (specifically T cells, natural killer cells and thymocytes) and can be used, for example, in the prophylaxis and treatment of T cell lymphoma or other related conditions. MEDI-507 is disclosed, e.g., in International Publication No. WO 99/03502, International Application Nos. PCT/US02/22273 and PCT/US02/06761, and U.S. Application Serial Nos. 09/462,140, 10/091,268, and 10/091,313, each of which is incorporated herein by reference in its entirety. MEDI-507 is a humanized IgG1k class monoclonal antibody that immunospecifically binds to human CD2 polypeptide. MEDI-507 was constructed using molecular techniques to insert the CDRs from the rat monoclonal antibody LO-CD2a/BTI-322 into a human IgG1 framework. LO-CD2a/BTI-322 has the amino acid sequence disclosed, e.g., in U.S. Patent Nos. 5,730,979, 5,817,311, and 5.951.983; and U.S. application Serial Nos. 09/056,072 and 09/462,140 (each of which is incorporated herein by reference in its entirety), or the amino acid sequence of the monoclonal antibody produced by the cell line deposited with the American Type Culture Collection

(ATCC®), 10801 University Boulevard, Manassas, Virginia 20110-2209 on July 28, 1993 as Accession Number HB 11423.

[00251] The anti- CD2 antibodies of this section can be made, formulated, administered, used therapeutically or prophylactically, or in other context as described in U.S. Patent Application No. 10/091,268, filed March 4, 2002, and published April 15, 2003, as U.S. Pat. Pub. No. US 2003/0068320; U.S. Patent Application No. 10/091,313, filed March 4, 2002, and published March 6, 2003, as U.S. Pat. Pub. No. US 2003/0044406; and U.S. Patent Application No. 10/657,006, filed September 5, 2003, and published December 30, 2004, as U.S. Pat. Pub. No. US 2004/0265315, the contents of which are hereby incorporated by reference in their entirety.

5.3.7 Antibodies that Immunopecifically Bind to CD19

[00252] The formulations of the present invention comprise an isolated antibody that immunospecifically binds to CD19 and a composition comprising this antibody. The antibodies can be monoclonal antibodies, human antibodies, humanized antibodies or chimeric antibodies. In some preferred embodiments, the anti-CD19 antibody of the invention is MT-103. MT-103 is the most-advanced clinical representative of a novel class of antibody derivatives called Bi-Specific T Cell Engagers (BiTE™). The BiTE compound MT-103 directs and activates the patient's own immune system against the cancer cells, stimulating T cells (a very potent type of white blood cell) to destroy B tumor cells (cancerous white blood cells). MT-103 specifically targets a particular protein (the CD19 antigen), which is present on cancerous B cells but not on other types of blood cells or healthy tissues, therefore avoiding the side effects of traditional chemotherapy

[00253] The anti- CD19 antibodies of this section can be made, formulated, administered, used therapeutically or prophylactically, or in other context as described in U.S. Pat. No. 6,723,538, and U.S. Pat. Pub. No. 2004/0162411.

[00254] The human CD19 molecule is a structurally distinct cell surface receptor that is expressed on the surface of human B cells. The invention relates to immunotherapeutic compositions and methods for the prophylaxis and treatment of GVHD, humoral rejection, and post-transplantation lymphoproliferative disorder in human subjects; autoimmune diseases and disorders; and cancers, using therapeutic antibodies that bind to the human CD19 antigen.

[00255] Hybridomas producing HB12a and HB12b anti-CD19 antibodies have been deposited under ATCC deposit nos. PTA-6580 and PTA-6581. See, also, U.S. Application No.

to be assigned (Attorney Docket No.: 11605-006-999) and U.S. Application No. 11/355,905, filed 2/15/2006, each of which is incorporated herein by reference in its entirety.

5.3.8 Antibodies that Immunopecifically Bind to EphA2

[00256] The formulations of the present invention comprise an isolated antibody that immunospecifically binds to EphA2 and a compositions comprising this antibody. The antibodies of the invention can be monoclonal antibodies, human antibodies, humanized antibodies or chimeric antibodies. In some embodiments, the anti-EphA2 antibody of the invention is EA2. In some preferred embodiments, the EA2 antibody is human or humanized. In other embodiments, the is EA5. In some preferred embodiments, the EA5 antibody is human or humanized. Hybridomas producing the anti-EphA2 antibodies of the invention have been deposited with the American Type Culture Collection (ATCC, P.O. Box 1549, Manassas, VA 20108) under the provisions of the Budapest Treaty on the International Recognition of the Deposit of Microorganisms for the Purposes of Patent Procedures, and assigned accession numbers, which are incorporated by reference, as shown in TABLE 4.

TABLE 4:

EphA2 Antibodies	Deposit No.	Date of Deposit
EA2.31	PTA-4380	May 22, 2002
EA5.12	PTA-4381	May 22, 2002
Eph099B-102.147	PTA-4572	August 7, 2002
Eph099B-208.261	PTA-4573	August 7, 2002
Eph099B-210.248	PTA-4574	August 7, 2002
Eph099B-233.152	PTA-5194	May 12, 2003
Eph101.530.241	PTA-4724	September 26, 2002

[00257] EphA2 is a 130 kDa receptor tyrosine kinase that is expressed in adult epithelia, where it is found at low levels and is enriched within sites of cell-cell adhesion (Zantek, et al, *Cell Growth & Differentiation* 10:629, 1999; Lindberg, et al., *Molecular & Cellular Biology* 10: 6316, 1990). EphA2 is upregulated on a large number of aggressive carcinoma cells. The anti-

EphA2 antibodies of this invention can be used, for example, in the treatment of a variety of tumors, including breast, colon, prostate, lung and skin cancers, as well as to prevent metastasis.

The anti-EphA2 antibodies of this section can be made, formulated, administered, used therapeutically or used prophylactically as described in U.S. Patent Application No. 10/823,259, filed April 12, 2004; U.S. Patent. Application No. 10/823,254, filed on April 12, 2004; U.S. Patent. Application No. 10/436,782, filed on May 12, 2003 and published February 12, 2004 as U.S. Pat. Pub. No. 2004/0028685; U.S. Patent. Application No. 10/436,783, filed on May 12, 2003 and published May 13, 2004 as U.S. Pat. Pub. No. 2004/0091486; U.S. Patent. Application No. 11/004,794, filed on December 3, 2004; U.S. Patent. Application No. 11/004,795, filed on December 3, 2004; u.S. Patent. Application No. 11/004,795, filed on December 3, 2004; and U.S. Provisional Application Nos. 60/662,517,60/622,711, 60/622,489, filed October 27, 2004, the contents of which are hereby incorporated by reference in their entirety.

5.3.9 Antibodies that Immunopecifically Bind to EphA4

[00259] The formulations of the present invention comprise an isolated antibody that immunospecifically binds to an antigen of EphA4 and a composition comprising this antibody. The antibodies of the invention can be monoclonal antibodies, human antibodies, humanized antibodies or chimeric antibodies. Hybridomas producing the anti-EphA4 antibodies of the invention have been deposited with the American Type Culture Collection (ATCC, P.O. Box 1549, Manassas, VA 20108) on June 4, 2004 under the provisions of the Budapest Treaty on the International Recognition of the Deposit of Microorganisms for the Purposes of Patent Procedures, and assigned accession number PTA-6044 and PTA-4381 and incorporated by reference.

[00260] EphA4 is a receptor tyrosine kinase that is expressed in brain, heart, lung, muscle, kidney, placenta, pancreas (Fox, et al, *Oncogene* 10:897, 1995) and melanocytes (Easty, et al., *Int. J. Cancer* 71:1061, 1997). EphA4 is overexpressed in a number of cancers. The anti-EphA4 antibodies of this section can be used, for example, to decrease the expression of EphA4 in the treatment of pancreatic cancers etc.

[00261] The anti-EphA4 antibodies of this section can be made, formulated, administered, used therapeutically or used prophylactically as described in U.S. Patent Application No. 10/863,729, filed June 7, 2004; U.S. Patent. Application No. 11/004,794, filed on December 3,

2004; U.S. Patent. Application Nos. 11/004,794 and 11/004,795, filed on December 3, 2004, the contents of which are hereby incorporated by reference in their entirety.

5.3.10 Antibodies that Immunopecifically Bind to IL-9

[00262] The formulations of the present invention comprise an antibody that immunospecifically binds to IL-9 and a composition comprising this antibody. The antibodies of the invention can be monoclonal antibodies, human antibodies, humanized antibodies or chimeric antibodies. In some preferred embodiments, the anti-IL-9 antibodies is MEDI-528.

[00263] It has been shown that IL-9 may be a key mediator of asthma and may also contribute to other respiratory disorders including chronic obstructive pulmonary disease (COPD) and cystic fibrosis. The anti-IL-9 antibodies of this section may be used in the prophylaxis or treatment of asthma.

[00264] The anti-IL-9 antibodies of this section can be made, formulated, administered, used therapeutically or used prophylactically as described in U.S. Patent Application No. 10/823,253, filed April 12, 2004 and published January 6, 2005, as U.S. Pat. Pub. No. US 2005/0002934 A1; U.S. Patent. Application No. 10/823, 810, filed on April 12, 2004; U.S. Provisional Application Nos. 60/371,728 and 60,371, 683, filed April 12, 2002; and U.S. Provisional Application No. 60/561,845, filed April 12, 2004, the contents of which are hereby incorporated by reference in their entirety.

5.3.11. Antibodies that Immunospecifically Bind to HMG1

[00265] The formulations of the present invention can comprise an antibody that immunospecifically binds to HMG1 and a composition comprising this antibody. The antibodies of the invention can be monoclonal antibodies, human antibodies, humanized antibodies or chimeric antibodies.

[00266] The early proinflammatory cytokines (e.g., TNF, IL-1, etc.) mediate inflammation, and induce the late release of high mobility group protein 1 (HMG1) (also known as HMG-1, HMG1, and HMGB1), a protein that accumulates in serum and mediates delayed lethality and further induction of early proinflammatory cytokines.

[00267] It has also been shown that HMG1 can be actively secreted by stimulated macrophages or monocytes in a process requiring acetylation of the molecule, which enables

translocation from the nucleus to secretory lysosomes and results in the secretion of an acetylated form of HMG1. *See*, PCT/IB2003/005718. Thus, HMG1 passively released from necrotic cells and HMGB1 actively secreted by inflammatory cells are molecularly different.

Further, HMG1 has been implicated as a cytokine mediator of delayed lethality in endotoxemia. *See, e.g.*, U.S. patents 6,468,533 and 6,448,223. More specifically, it has been demonstrated that bacterial endotoxin (lipopolysaccharide (LPS)) activates monocytes/macrophages to release HMG1 as a late response to activation, resulting in elevated serum HMG1 levels that are toxic. Antibodies against HMG1 have been shown to prevent lethality of endotoxin even when antibody administration is delayed until after the early cytokine response. Like other proinflammatory cytokines, HMG1 is a potent activator of monocytes. Intratracheal application of HMG1 causes acute lung injury, and anti-HMG1 antibodies protect against endotoxin-induced lung edema. In addition, serum HMG1 levels are elevated in critically ill patients with sepsis or hemorrhagic shock, and levels are significantly higher in non-survivors as compared to survivors.

[00269] The anti-HMG1 antibodies of this section can be made, formulated, administered, used therapeutically or used prophylactically as described in U.S. Patent Publication No. 2006-0099207 A1 filed October 21, 2005, which is incorporated herein by reference in its entirety. Three clones, S6, S16 and G4 have been deposited with the American Type Culture Collection (10801 University Boulevard, Manassas, Va. 20110-2209) and assigned ATCC Deposit Nos. PTA-6143 (Deposited August 4, 2004), PTA-6259 (Deposited October 19, 2004) and PTA-6258 (Deposited October 19, 2004) (also referred to herein as "S6", "S16", and "G4", respectively) as described in U.S. Patent Publication No. 2006-0099207 A1 filed October 21, 2005, which is incorporated herein by reference in its entirety.

5.3.12. Antibodies that Immunospecifically Bind to ALK

[00270] The formulations of the present invention can comprise an antibody that immunospecifically binds to ALK and a composition comprising this antibody. The antibodies of the invention can be monoclonal antibodies, human antibodies, humanized antibodies or chimeric antibodies.

[00271] Monoclonal antibodies against ALK as well as hybridoma cell lines producing ALK monoclonal antibodies 8B10, 16G2-3 and 9C10-5 (deposited with the American Type Culture Collection (10801 University Boulevard, Manassas, Va. 20110-2209) and assigned ATCC Deposit Nos. to be assigned, respectively) as described in U.S. Patent Application No.

09/880,097, filed 06-14-2001 and published March 21, 2002, as U.S. Pat. Pub. No. 20020034768, which is incorporated herein by reference in its entirety.

Pleiotrophin (PTN) is a 136-amino acid, secreted, heparin-binding cytokine that has diverse functions including a role in angiogenesis. PTN has been shown to specifically bind to a receptor tyrosine kinase, Anaplastic Lymphoma Kinase (ALK), and such binding leads to auto-phosphorylation of the receptor and subsequent phosphorylation of a number of signal transduction molecules such as IRS-1, PLC-gamma, PI3 kinase, and Shc, and activates a cell survival pathway. See PCT Pat. App. Pub. No. WO 01/96364. Accordingly, agents and therapeutic treatments that regulate ALK-mediated signal transduction pathways can affect one or more ALK-regulated functions, including, for example, angiogenesis. ALK participates in various disease states, including cancers and diseases related to unwanted or excessive angiogenesis. Additionally, ALK participates in a desirable way in certain processes, such as wound healing. ALK and/or PTN are expressed, often at high levels, in a variety of tumors. Therefore, agents that downregulate ALK and/or PTN function may affect tumors by a direct effect on the tumor cells, an indirect effect on the angiogenic processes recruited by the tumor, or a combination of direct and indirect effects.

5.3.13. Antibodies that Immunospecifically Bind to CD20

[00273] The formulations of the present invention can comprise an antibody that immunospecifically binds to CD20 and a composition comprising this antibody. The antibodies of the invention can be monoclonal antibodies, human antibodies, humanized antibodies or chimeric antibodies.

[00274] CD20 is only expressed by B lymphocytes (Stashenko et al. (1980) *J Immunol* 125:1678-1685; Tedder et al., 1988a). CD20 forms a homo- or hetero-tetrameric complex that is functionally important for regulating cell cycle progression and signal transduction in B lymphocytes (Tedder and Engel, 1994). CD20 additionally regulates transmembrane Ca⁺⁺ conductance, possibly as a functional component of a Ca⁺⁺-permeable cation channel (Bubien et al. *J Cell Biol* 121:1121-1132; Kanzaki et al. (1997a) *J Biol Chem* 272:14733-14739; Kanzaki et al. (1997b) *J Biol Chem* 272:4964-4969; Kanzaki et al. (1995) *J Biol Chem* 270:13099-13104). Antibodies against CD20 are effective in treating non-Hodgkin's lymphoma (McLaughlin et al. (1998) *Oncology* 12:1763-1769; Onrust et al. (1989) *J Biol Chem* 264:15323-15327; Weiner (1999) *Semin Oncol* 26:43-51).

See, also, US Patent Application No. 10/433,287, filed September 30, 2003, published as US 20040137566 on July 15, 2004, which is incorporated herein by reference in its entirety.

5.3.14. Antibodies that Immunospecifically Bind to CD22

[00275] The formulations of the present invention can comprise an antibody that immunospecifically binds to CD22 and a composition comprising this antibody. The antibodies of the invention can be monoclonal antibodies, human antibodies, humanized antibodies or chimeric antibodies.

[00276] Anti-CD22 antibodies have been described, for example, in U.S. Pat. Nos. 5,484,892; 6,183,744; 6,187,287; 6,254,868; 6,306,393, and in Tuscano *et al.*, *Blood* 94(4):1382-92 (1999) (each of which is incorporated herein in its entirety by reference). The use of monoclonal antibodies, including anti-CD22 antibodies, in the treatment of non-Hodgkin's lymphoma is reviewed, for example, by Renner *et al.*, *Leukemia* 11(Suppl. 2):S5509 (1997).

[00277] The use of humanized CD22 antibodies has been described for the treatment of autoimmune disorders (*see*, Tedder U.S. Patent Application Publication No. US2003/0202975) and for the treatment of B cell malignancies, such as lymphomas and leukemias (*see*, Tuscano U.S. Patent Application Publication No. U.S. 2004/0001828). Humanized CD22 antibodies that target specific epitopes on CD22 have been described for use in immunoconjugates for therapeutic uses in cancer (*see* U.S. Patent Nos. 5,789,554 and 6,187,287 to Leung).

[00278] Exemplary VH and VK antibody regions of the invention were deposited with the American Type Culture Collection (ATCC). In particular, a plasmid encoding the humanized anti-CD22 VH sequence of the invention designated RHOv2 was deposited under ATCC deposit no. PTA-7372, on February 9, 2006. A plasmid encoding the humanized anti-CD22 VH sequence of the invention designated RHOv2ACD was deposited under ATCC deposit no. PTA-7373, on February 9, 2006. A plasmid encoding the humanized anti-CD22 VK sequence of the invention, RKA was deposited under ATCC deposit no. PTA-7370, on February 9, 2006. A plasmid encoding the humanized anti-CD22 VK sequence of the invention, RKC, was deposited under ATCC deposit no. PTA-7371, on February 9, 2006.

[00279] See, also, U.S. Provisional Application No. TBA, filed March 6, 2006, attorney docket no. BC320P1, which is incorporated herein by reference in its entirety.

5.3.15. Antibodies that Immunospecifically Bind to Chitinase

[00280] The formulations of the present invention can comprise an antibody that immunospecifically binds to Chitinase and a composition comprising this antibody. The antibodies of the invention can be monoclonal antibodies, human antibodies, humanized antibodies or chimeric antibodies.

[00281] It is described that blocking a chitinase/chitinase-like protein, in vivo results in protection of bone and cartilage as well as a reduction in weight loss in a mouse RA model. These results support the role of chitinase/chitinase-like proteins in chronic inflammatory diseases and more specifically the role of chitinase/chitinase-like proteins in OCL-related diseases including bone metabolism and connective tissue disorders and diseases. Furthermore, these results validate human chitinase/chitinase-like proteins as potential therapeutic targets for the prevention and treatment of OCL-related diseases.

[00282] See, also, U.S. Application No. 10/202,436, filed July 23, 2002, published as US 20030049261 on March 13, 2003, which is incorporated herein by reference in its entirety.

5.3.16. Antibodies that Immunospecifically Bind to Interferon alpha

[00283] The formulations of the present invention can comprise an antibody that immunospecifically binds to interferon alpha and a composition comprising this antibody. The antibodies of the invention can be monoclonal antibodies, human antibodies, humanized antibodies or chimeric antibodies.

The invention provides a method of treating an interferon alpha-mediated disease or disorder in a subject, comprising administering to the subject an anti-IFN alpha antibody of the invention, such that the interferon-alpha mediated disease in the subject is treated. Examples of diseases that can be treated include autoimmune diseases (*e.g.*, systemic lupus erythematosus, multiple sclerosis, insulin dependent diabetes mellitus, inflammatory bowel disease, psoriasis, autoimmune thyroiditis, rheumatoid arthritis and glomerulonephritis), transplant rejection and graft versus host disease.

[00285] Anti-interferon alpha monoclonal antibody has also been described in U.S. serial no. 11/009,410 filed December 10, 2004, which is incorporated herein by reference in its entirety.

5.3.17. Antibodies that Immunospecifically Bind to Interferon alpha receptor

[00286] The formulations of the present invention can comprise an antibody that immunospecifically binds to interferon alpha receptor and a composition comprising this antibody. The antibodies of the invention can be monoclonal antibodies, human antibodies, humanized antibodies or chimeric antibodies.

[00287] The invention also provides a method for inhibiting biological activity of a type I interferon on a cell expressing interferon alpha receptor 1 comprising contacting the cell with the antibody of the invention, such that the biological activity of the type I interferon is inhibited. The invention also provides a method of treating a type I interferon-mediated disease or disorder in a subject in need of treatment comprising administering to the subject the antibody, or antigen-binding portion thereof, of the invention, such that the type-I interferon mediated disease in the subject is treated. The type I interferon-mediated disease can be, for example, an interferon alpha-mediated disease.

[00288] Examples of disease or disorders that can be treated using the methods of the invention include systemic lupus erythematosus, insulin dependent diabetes mellitus, inflammatory bowel disease, multiple sclerosis, psoriasis, autoimmune thyroiditis, rheumatoid arthritis, glomerulonephritis, HIV infection, AIDS, transplant rejection and graft versus host disease.

[00289] Anti-interferon receptor monoclonal antibody has been described in U.S. Patent Publication No. 2006-0029601 A1, published 2/9/2006, filed June 20, 2005, which is incorporated herein by reference in its entirety.

5.3.18 Antibodies That Have Therapeutic Utility

[00290] The formulations of the present invention comprise antibodies that have therapeutic utility, including but not limited to antibodies listed in Table 5.

[00291] TABLE 5. THERAPEUTIC ANTIBODIES THAT CAN BE USED IN CONNECTION WITH THE PRESENT INVENTION

Company	Product	Disease	Target
Abgenix AltaRex	ABX-EGF OvaRex BravaRex	Cancer ovarian cancer metastatic cancers	EGF receptor tumor antigen CA125 tumor antigen MUC1
Antisoma	Theragyn (pemtumomabytrrium- 90)	ovarian cancer	PEM antigen

Company	Product	Disease	Target
	Therex	breast cancer	PEM antigen
Boehringer	Blvatuzumab	head & neck	CD44
Ingelheim		cancer	
Centocor/J&J	Panorex	Colorectal	17-1A
		cancer	
	ReoPro	PTCA	Gp IIIb/IIIa
	ReoPro	Acute MI	Gp IIIb/IIIa
	ReoPro	Ischemic stroke	Gp IIIb/IIIa
Corixa	Bexocar	NHL	$\overline{\text{CD20}}$
CRC Technology	MAb, idiotypic 105AD7	colorectal cancer	Gp72
Cite roumoiogy	2.2120, 2020 J pao	vaccine	-1
Crucell	Anti-EpCAM	cancer	Ep-CAM
Cytoclonal	MAb, lung cancer	non-small cell	NA
CytoCionai	THE TO, THIS CHILDS	lung cancer	
Genentech	Herceptin	metastatic breast	HER-2
Generation	Погеории	cancer	
	Herceptin	early stage	HER-2
	погсерии	breast cancer	TILLE L
	Rituxan	Relapsed/refract	CD20
	Kituxan	ory low-grade or	CB20
		follicular NHL	
	Rituxan	intermediate &	CD20
	Kituxan	high-grade NHL	CDZ0
	MAb-VEGF	NSCLC,	VEGF
	MAG-VEGI	metastatic	V DOI
	MAb-VEGF	Colorectal	VEGF
	MAU-VEGI	cancer,	V LOI
		metastatic	
	AMD Fab	age-related	CD18
	AMD rao	macular	CD10
		degeneration	
	E-26 (2 nd gen. IgE)	allergic asthma	IgE
	E-20 (2 gcn. igE)	& rhinitis	1gL
IDEC	Zevalin (Rituxan +	low grade of	CD20
IDEC	yttrium-90)	follicular,	CD20
	yttridin-50)	relapsed or	
		refractory,	
		CD20-positive,	
		B-cell NHL and	
		Rituximab-	
		refractory NHL	
ImClone	Cetuximab + innotecan	refractory NTIL	EGF receptor
micione		colorectal	Lor receptor
		carcinoma	
	Caturrimoh + signilatin &	newly diagnosed	EGF receptor
	Cetuximab + cisplatin &		TOI ICCOPIUI
		חר דברווזיים אל אמרוים	
	radiation	or recurrent head	
	radiation	& neck cancer	EGE recentor
			EGF receptor

Company	Product	Disease	Target
		pancreatic	
	O-4000100101010101010101	carcinoma	DCD .
	Cetuximab + cisplatin + 5FU or Taxol	recurrent or	EGF receptor
	JFU OF Taxor	metastatic head & neck cancer	
	Cetuximab +	newly diagnosed	EGF receptor
	carboplatin + paclitaxel	non-small cell	EGT receptor
	caroopiatiii - paciitaxci	lung carcinoma	
	Cetuximab + cisplatin	head & neck	EGF receptor
		cancer	F
		(extensive	
		incurable local-	
		regional disease	1
		& distant	
		metasteses)	
	Cetuximab + radiation	locally advanced	EGF receptor
		head & neck	
	DECC \ D :II	carcinoma	
	BEC2 + Bacillus	small cell lung	mimics ganglioside
	Calmette Guerin BEC2 + Bacillus	carcinoma	GD3
	Calmette Guerin	melanoma	mimics ganglioside GD3
	IMC-1C11	colorectal cancer	VEGF-receptor
	INIC-1CII	with liver	V LGT-receptor
		metasteses	
ImmonoGen	nuC242-DM1	Colorectal,	nuC242
		gastric, and	
		pancreatic	
		cancer	
ImmunoMedics	LymphoCide	Non-Hodgkins	CD22
		lymphoma	
	LymphoCide Y-90	Non-Hodgkins	CD22
	CEA CL	lymphoma	OF A
	CEA-Cide	metastatic solid	CEA
	CEA-Cide Y-90	tumors metastatic solid	CEA
	CLA-Clue 1-90	tumors	CLA
	CEA-Scan (Tc-99m-	colorectal cancer	CEA
	labeled arcitumomab)	(radioimaging)	
	CEA-Scan (Tc-99m-	Breast cancer	CEA
	labeled arcitumomab)	(radioimaging)	
	CEA-Scan (Tc-99m-	lung cancer	CEA
	labeled arcitumomab)	(radioimaging)	
	CEA-Scan (Tc-99m-	intraoperative	CEA
	labeled arcitumomab)	tumors (radio	
	T 1 0 00	imaging)	
	LeukoScan (Tc-99m-	soft tissue	CEA
	labeled sulesomab)	infection	
	,	(radioimaging)	

Company	Product	Disease	Target
	LymphoScan (Tc-99m-	lymphomas	CD22
	labeled)	(radioimaging)	
	AFP-Scan (Tc-99m-	liver 7 gem-cell	AFP
	labeled)	cancers	
		(radioimaging)	
Intracel	HumaRAD-HN (+	head & neck	NA
	yttrium-90)	cancer	
	HumaSPECT	colorectal	NA
3.5.1	TOTAL OF COURT AND	imaging	COT 1 4
Medarex	MDX-101 (CTLA-4)	Prostate and	CTLA-4
	MDW 010 (1)	other cancers	TIED 0
	MDX-210 (her-2	Prostate cancer	HER-2
	overexpression)	O	HED 0
M. II	MDX-210/MAK	Cancer	HER-2
MedImmune Merck KGaA	Vitaxin	Cancer	ανβ ₃
Merck KGaA	MAb 425	Various cancers	EGF receptor
3 6:11	IS-IL-2	Various cancers	Ep-CAM CD52
Millennium	Campath	chronic	CD32
	(alemtuzumab)	lymphocytic leukemia	
NeoRx	CD20 atmentagidin (1)		CD20
NeoKx	CD20-streptavidin (+	Non-Hodgkins	CD20
	biotin-yttrium 90) Avidicin (albumin +	lymphoma metastatic	NA
	NRLU13)		IVA
	NKLU15)	cancer	
Peregrine	Oncolym (+ iodine-131)	Non-Hodgkins	HLA-DR 10 beta
		lymphoma	
	Cotara (+ iodine-131)	unresectable	DNA-associated
		malignant	proteins
		glioma	
Pharmacia	C215 (+ staphylococcal	pancreatic	NA
Corporation	enterotoxin)	cancer	771
	MAb, lung/kidney	lung & kidney	NA
	cancer	cancer	3.7. A
	nacolomab tafenatox	colon &	NA
	(C242 + staphylococcal	pancreatic	
D. Asia Dania	enterotoxin)	cancer	CD2
Protein Design	Nuvion	T cell	CD3
Labs	CNAADT NAIDE	malignancies	CD22
	SMART M195	AML NHL	CD33
Titom	SMART 1D10		HLA-DR antigen
Titan	CEAVac	colorectal	CEA
		cancer,	
	TriCom	advanced	CD2 genelieside
	TriGem	metastatic	GD2-ganglioside
		melanoma &	
		small cell lung	
	Tri Ah	cancer metastatic broast	MUC-1
	TriAb	metastatic breast	MIOC-1

Company	Product	Disease	Target
		cancer	
Trilex	CEAVac	colorectal	CEA
		cancer,	
		advanced	
	TriGem	metastatic	GD2-ganglioside
		melanoma &	
		small cell lung	
	73 * 4.1	cancer	
	TriAb	metastatic breast	MUC-1
Viventia Biotech	NovoMAb-G2	cancer	
vivenua biotech	radiolabeled	Non-Hodgkins	NA
		lymphoma	CTT 4
	Monopharm C	colorectal &	SK-1 antigen
		pancreatic carcinoma	
	GlioMAb-H (+ gelonin	gliorna,	NA
	toxin)	melanoma &	INV
	· · · · · · · · · · · · · · · · · · ·	neuroblastoma	
Xoma	Rituxan	Relapsed/refract	CD20
		ory low-grade or	
		follicular NHL	
	Rituxan	intermediate &	CD20
		high-grade NHL	
	ING-1	adenomearcino	Ep-CAM
		ma	

5.3.19. <u>Antibodies That Can Be Used For Inflammatory</u> <u>Disorders or Autoimmune Diseases</u>

[00292] The formulations of the present invention further comprises any of the antibodies known in the art for the treatment and/or prevention of autoimmune disease or inflammatory disease. A non-limiting example of the antibodies that are used for the treatment or prevention of inflammatory disorders which can be engineered according to the invention is presented in Table 6A, and a non-limiting example of the antibodies that are used for the treatment or prevention of autoimmune disorder is presented in Table 6B.

[00293] TABLE 6A: ANTIBODIES FOR INFLAMMATORY DISEASES AND AUTOIMMUNE DISEASES THAT CAN USED IN ACCORDANCE WITH THE INVENTION.

Antibody Name	Target Antigen	Product Type	Isotype	Sponsors	Indication
5G1.1	Complement (C5)	Humanized	IgG	Alexion Pharm Inc	Rheumatoid Arthritis
5G1.1	Complement (C5)	Humanized	IgG	Alexion Pharm Inc	SLE

Antibody Name	Target Antigen	Product Type	Isotype	Sponsors	Indication
5G1.1	Complement	Humanized	IgG	Alexion	Nephritis
	(C5)			Pharm Inc	
5G1.1-SC	Complement	Humanized	ScFv	Alexion	Cardiopulmonary
	(C5)			Pharm Inc	Bypass
5G1.1-SC	Complement	Humanized	ScFv	Alexion	Myocardial
	(C5)	<u> </u>		Pharm Inc	Infarction
5G1.1-SC	Complement	Humanized	ScFv	Alexion	Angioplasty
	(C5)			Pharm Inc	
ABX-CBL	CBL	Human		Abgenix Inc	GvHD
ABX-CBL	CD147	Murine	IgG	Abgenix Inc	Allograft rejection
ABX-IL8	IL-8	Human	IgG2	Abgenix Inc	Psoriasis
Antegren	VLA-4	Humanized	IgG	Athena/Elan	Multiple Sclerosis
Anti-	CD11a	Humanized	IgG1	Genentech	Psoriasis
CD11a		1		Inc/Xoma	Soriusis
Anti-CD18	CD18	Humanized	Fab'2	Genentech Inc	Myocardial
	1				infarction
Anti-LFA1	CD18	Murine	Fab'2	Pasteur-	Allograft rejection
				Merieux/	z mogran rejection
	1			Immunotech	ļ
Antova	CD40L	Humanized	IgG	Biogen	Allograft rejection
Antova	CD40L	Humanized	IgG	Biogen	SLE
BTI-322	CD2	Rat	IgG	Medimmune	GvHD, Psoriasis
			180	Inc	GVID, FSonasis
CDP571	TNF-alpha	Humanized	IgG4	Celltech	Crohn's
CDP571	TNF-alpha	Humanized	IgG4	Celltech	Rheumatoid Arthritis
CDP850	E-selectin	Humanized		Celltech	Psoriasis
Corsevin	Fact VII	Chimeric		Centocor	Anticoagulant
M				Contocor	7 Hiticoagulant
D2E7	TNF-alpha	Human		CAT/BASF	Rheumatoid
	1			CITI/BITOI	Arthritis
Hu23F2G	CD11/18	Humanized		ICOS Pharm	Multiple Sclerosis
				Inc	Muniple Scietosis
Hu23F2G	CD11/18	Humanized	IgG	ICOS Pharm	Stroke
			-50	Inc	PHOVE
IC14	CD14		 -	ICOS Pharm	Toxic shock
		.		Inc	TOXIC SHOCK
ICM3	ICAM-3	Humanized		ICOS Pharm	Psoriasis
	1 2 2 2 2 1 1 2	- Aumanized		Inc	r soriasis
IDEC-114	CD80	Primatised		IDEC	Danissis
		1 mauscu		L I	Psoriasis
				Pharm/Mitsub	
IDEC-131	CD40L	Humanized	 -	ishi	CLE
U/LC-131	CDTOL	rumanized		IDEC	SLE
DEC-131	CD40L	LTypes and 1		Pharm/Eisai	16 12 1 5 1
MEC-131	CD40L	Humanized		IDEC	Multiple Sclerosis
DEC-151	CD4	Desire	T. C1	Pharm/Eisai	7.1
DEC-131	CD4	Primatised	Ig G 1	IDEC	Rheumatoid
		,		Pharm/Glaxo	Arthritis
				SmithKline	

Antibody Name	Target Antigen	Product Type	Isotype	Sponsors	Indication
IDEC-152	CD23	Primatised	·	IDEC Pharm	Asthma/Allergy
Infliximab	TNF-alpha	Chimeric	IgG1	Centocor	Rheumatoid Arthritis
Infliximab	TNF-alpha	Chimeric	IgG1	Centocor	Crohn's
LDP-01	beta2- integrin	Humanized	IgG	Millennium Inc (LeukoSite Inc.)	Stroke
LDP-01	beta2- integrin	Humanized	IgG	Millennium Inc (LeukoSite Inc.)	Allograft rejection
LDP-02	alpha4beta7	Humanized		Millennium Inc (LeukoSite Inc.)	Ulcerative Colitis
MAK- 195F	TNF alpha	Murine	Fab'2	Knoll Pharm, BASF	Toxic shock
MDX-33	CD64 (FcR)	Human		Medarex/Cent eon	Autoimmune haematogical disorders
MDX-CD4	CD4	Human	IgG	Medarex/Eisai / Genmab	
MEDI-507	CD2	Humanized		Medimmune Inc	Psoriasis
MEDI-507	CD2	Humanized		Medimmune Inc	GvHD
OKT4A	CD4	Humanized	IgG	Ortho Biotech	Allograft rejection
OrthoClon e OKT4A	CD4	Humanized	IgG	Ortho Biotech	Autoimmune disease
Orthoclone / anti-CD3 OKT3	CD3	Murine	mIgG2a	Ortho Biotech	Allograft rejection
RepPro/ Abciximab	gpIIbIIIa		Fab	Centocor/Lill y	Complications of coronary angioplasty
rhuMab- E25	IgE		IgG1	Genentech/No vartis/Tanox Biosystems	Asthma/Allergy
SB-240563	IL5	Humanized		GlaxoSmithKl ine	Asthma/Allergy
SB-240683	IL-4	Humanized		GlaxoSmithKl ine	Asthma/Allergy
	IL-5	Humanized		Celltech/Sche ring	Asthma/Allergy
Simulect	CD25	Chimeric	IgG1	Novartis Pharm	Allograft rejection

Antibody	Target	Product Type	Isotype	Sponsors	Indication
Name	Antigen				
SMART	CD3	Humanized		Protein	Autoimmune
a-CD3				Design Lab	disease
SMART	CD3	Humanized		Protein	Allograft rejection
a-CD3				Design Lab	
SMART	CD3	Humanized	IgG	Protein	Psoriasis
a-CD3				Design Lab	
Zenapax	CD25	Humanized	IgG1	Protein	Allograft rejection
1				Design	
				Lab/Hoffman-	
				La Roche	

TABLE 6B: ANTIBODIES FOR AUTOIMMUNE DISORDERS THAT CAN BE USED IN ACCORDANCE WITH THE INVENTION

Antibody	Indication	Target Antigen
ABX-RB2		antibody to CBL antigen on T cells, B cells and NK cells fully human antibody from the Xenomouse
5c8 (Anti CD-40 ligand antibody)	Phase II trials were halted in Oct. 99 examine "adverse events"	CD-40
IDEC 131	systemic lupus erythyematous (SLE)	anti CD40 humanized
IDEC 151	rheumatoid arthritis	primatized; anti-CD4
IDEC 152	Asthma	primatized; anti-CD23
IDEC 114	Psoriasis	primatized anti-CD80
MEDI-507	rheumatoid arthritis; multiple sclerosis Crohn's disease Psoriasis	anti-CD2
LDP-02 (anti-b7 mAb)	inflammatory bowel disease Chron's disease ulcerative colitis	a4b7 integrin receptor on white blood cells (leukocytes)
SMART Anti- Gamma Interferon antibody	autoimmune disorders	Anti-Gamma Interferon
Verteportin	rheumatoid arthritis	
MDX-33	blood disorders caused by autoimmune reactions Idiopathic Thrombocytopenia Purpurea (ITP)	monoclonal antibody against FcRI receptors

Antibody	Indication	Target Antigen
	autoimmune hemolytic anemia	
MDX-CD4	treat rheumatoid arthritis and other autoimmunity	monoclonal antibody against CD4 receptor molecule
VX-497	autoimmune disorders multiple sclerosis rheumatoid arthritis inflammatory bowel disease lupus psoriasis	inhibitor of inosine monophosphate dehydrogenase (enzyme needed to make new RNA and DNA used in production of nucleotides needed for lymphocyte proliferation)
VX-740	rheumatoid arthritis	inhibitor of ICE interleukin-1 beta (converting enzyme controls pathways leading to aggressive immune response)
VX-745	specific to inflammation involved in chemical signalling of immune response onset and progression of inflammation	inhibitor of P38MAP kinase mitogen activated protein kinase
Enbrel (etanercept)		targets TNF (tumor necrosis factor)
IL-8		fully human monoclonal antibody against IL-8 (interleukin 8)
Apogen MP4		recombinant antigen selectively destroys disease associated T-cells induces apoptosis T-cells eliminated by programmed cell death no longer attack body's own cells specific apogens target specific T- cells

5.4 Methods of Producing Antibodies

[00294] The antibodies used in the present invention can be produced by any method known in the art for the synthesis of antibodies, in particular, by chemical synthesis or preferably, by recombinant expression techniques.

[00295] Monoclonal antibodies can be prepared using a wide variety of techniques known in the art including the use of hybridoma, recombinant, and phage display technologies, or a

combination thereof. For example, monoclonal antibodies can be produced using hybridoma techniques including those known in the art and taught, for example, in Harlow et al., *Antibodies: A Laboratory Manual*, (Cold Spring Harbor Laboratory Press, 2nd ed. 1988); Hammerling, et al., in: *Monoclonal Antibodies and T-Cell Hybridomas* 563-681 (Elsevier, N.Y., 1981) (said references incorporated by reference in their entireties). The term "monoclonal antibody" as used herein is not limited to antibodies produced through hybridoma technology. The term "monoclonal antibody" refers to an antibody that is derived from a single clone, including any eukaryotic, prokaryotic, or phage clone, and not the method by which it is produced.

[00296] Methods for producing and screening for specific antibodies using hybridoma technology are routine and well known in the art. Briefly, mice can be immunized with an antigen (either the full length protein or a domain thereof, *e.g.*, the extracellular or the ligand binding domain) and once an immune response is detected, *e.g.*, antibodies specific for the particular antigen are detected in the mouse serum, the mouse spleen is harvested and splenocytes isolated. The splenocytes are then fused by well known techniques to any suitable myeloma cells, for example cells from cell line SP20 available from the ATCC. Hybridomas are selected and cloned by limited dilution. Hybridoma clones are then assayed by methods known in the art for cells that secrete antibodies capable of binding a polypeptide of the invention. Ascites fluid, which generally contains high levels of antibodies, can be generated by immunizing mice with positive hybridoma clones.

[00297] Accordingly, monoclonal antibodies can be generated by culturing a hybridoma cell secreting an antibody of the invention wherein, preferably, the hybridoma is generated by fusing splenocytes isolated from a mouse immunized with the antigen with myeloma cells and then screening the hybridomas resulting from the fusion for hybridoma clones that secrete an antibody able to bind the antigen.

[00298] Antibody fragments used in the present invention may be generated by any technique known to those of skill in the art. For example, Fab and F(ab')2 fragments of the invention may be produced by proteolytic cleavage of immunoglobulin molecules, using enzymes such as papain (to produce Fab fragments) or pepsin (to produce F(ab')2 fragments). F(ab')2 fragments contain the variable region, the light chain constant region and the CH1 domain of the heavy chain. Further, the antibodies of the present invention can also be generated using various phage display methods known in the art.

In phage display methods, functional antibody domains are displayed on the [00299] surface of phage particles which carry the polynucleotide sequences encoding them. In particular, DNA sequences encoding VH and VL domains are amplified from animal cDNA libraries (e.g., human or murine cDNA libraries of lymphoid tissues). The DNA encoding the VH and VL domains are recombined together with an scFv linker by PCR and cloned into a phagemid vector (e.g., p CANTAB 6 or pComb 3 HSS). The vector is electroporated in E. coli and the E. coli is infected with helper phage. Phage used in these methods are typically filamentous phage including fd and M13 and the VH and VL domains are usually recombinantly fused to either the phage gene III or gene VIII. Phage expressing an antigen binding domain that binds to an epitope of interest can be selected or identified with antigen, e.g., using labeled antigen or antigen bound or captured to a solid surface or bead. Examples of phage display methods that can be used to make the antibodies of the present invention include those disclosed in Brinkman et al., 1995, J. Immunol. Methods 182:41-50; Ames et al., 1995, J. Immunol. Methods 184:177; Kettleborough et al., 1994, Eur. J. Immunol. 24:952-958; Persic et al., 1997, Gene 187:9; Burton et al., 1994, Advances in Immunology 57:191-280; International Application No. PCT/GB91/01134; International Publication Nos. WO 90/02809, WO 91/10737, WO 92/01047, WO 92/18619, WO 93/1 1236, WO 95/15982, WO 95/20401, and WO97/13844; and U.S. Patent Nos. 5,698,426, 5,223,409, 5,403,484, 5,580,717, 5,427,908, 5,750,753, 5,821,047, 5,571,698, 5,427,908, 5,516,637, 5,780,225, 5,658,727, 5,733,743 and 5,969,108; each of which is incorporated herein by reference in its entirety.

[00300] Phage may be screened for antigen binding activities. As described in the above references, after phage selection, the antibody coding regions from the phage can be isolated and used to generate whole antibodies, including human antibodies, or any other desired antigen binding fragment, and expressed in any desired host, including mammalian cells, insect cells, plant cells, yeast, and bacteria, *e.g.*, as described below. Techniques to recombinantly produce Fab, Fab' and F(ab')2 fragments can also be employed using methods known in the art such as those disclosed in International Publication No. WO 92/22324; Mullinax et al., 1992, *BioTechniques* 12:864; Sawai et al., 1995, *AJRI* 34:26; and Better et al., 1988, *Science* 240:1041 (said references incorporated by reference in their entireties).

[00301] To generate whole antibodies, PCR primers including VH or VL nucleotide sequences, a restriction site, and a flanking sequence to protect the restriction site can be used to amplify the VH or VL sequences in scFv clones. Utilizing cloning techniques known to those of skill in the art, the PCR amplified VH domains can be cloned into vectors expressing a VH

constant region, *e.g.*, the human gamma 4 constant region, and the PCR amplified VL domains can be cloned into vectors expressing a VL constant region, *e.g.*, human kappa or lambda constant regions. Preferably, the vectors for expressing the VH or VL domains comprise an EF-1 α promoter, a secretion signal, a cloning site for the variable domain, constant domains, and a selection marker such as neomycin. The VH and VL domains may also be cloned into one vector expressing the necessary constant regions. The heavy chain conversion vectors and light chain conversion vectors are then co-transfected into cell lines to generate stable or transient cell lines that express full-length antibodies, *e.g.*, IgG, using techniques known to those of skill in the art.

[00302] For some uses, including *in vivo* use of antibodies in humans and *in vitro* detection assays, it may be preferable to use human or chimeric antibodies. Completely human antibodies are particularly desirable for therapeutic treatment of human subjects. Human antibodies can be made by a variety of methods known in the art including phage display methods described above using antibody libraries derived from human immunoglobulin sequences. See also U.S. Patent Nos. 4,444,887 and 4,716,111; and International Publication Nos. WO 98/46645, WO 98/50433, WO 98/24893, WO 98/16654, WO 96/34096, WO 96/33735, and WO 91/10741; each of which is incorporated herein by reference in its entirety.

Human antibodies can also be produced using transgenic mice which are [00303] incapable of expressing functional endogenous immunoglobulins, but which can express human immunoglobulin genes. For example, the human heavy and light chain immunoglobulin gene complexes may be introduced randomly or by homologous recombination into mouse embryonic stem cells. Alternatively, the human variable region, constant region, and diversity region may be introduced into mouse embryonic stem cells in addition to the human heavy and light chain genes. The mouse heavy and light chain immunoglobulin genes may be rendered non-functional separately or simultaneously with the introduction of human immunoglobulin loci by homologous recombination. In particular, homozygous deletion of the J_H region prevents endogenous antibody production. The modified embryonic stem cells are expanded and microinjected into blastocysts to produce chimeric mice. The chimeric mice are then be bred to produce homozygous offspring which express human antibodies. The transgenic mice are immunized in the normal fashion with a selected antigen, e.g., all or a portion of a polypeptide of the invention. Monoclonal antibodies directed against the antigen can be obtained from the immunized, transgenic mice using conventional hybridoma technology. The human immunoglobulin transgenes harbored by the transgenic mice rearrange during B cell

differentiation, and subsequently undergo class switching and somatic mutation. Thus, using such a technique, it is possible to produce therapeutically useful IgG, IgA, IgM and IgE antibodies. For an overview of this technology for producing human antibodies, see Lonberg and Huszar (1995, *Int. Rev. Immunol.* 13:65-93). For a detailed discussion of this technology for producing human antibodies and human monoclonal antibodies and protocols for producing such antibodies, *see*, *e.g.*, International Publication Nos. WO 98/24893, WO 96/34096, and WO 96/33735; and U.S. Patent Nos. 5,413,923, 5,625,126, 5,633,425, 5,569,825, 5,661,016, 5,545,806, 5,814,318, and 5,939,598, which are incorporated by reference herein in their entirety. In addition, companies such as Abgenix, Inc. (Fremont, CA) and Medarex (Princeton, NJ) can be engaged to provide human antibodies directed against a selected antigen using technology similar to that described above.

[00304] A chimeric antibody is a molecule in which different portions of the antibody are derived from different immunoglobulin molecules such as antibodies having a variable region derived from a non-human antibody and a human immunoglobulin constant region. Methods for producing chimeric antibodies are known in the art. See, e.g., Morrison, 1985, Science 229:1202; Oi et al., 1986, BioTechniques 4:214; Gillies et al., 1989, J. Immunol. Methods 125:191-202; and U.S. Patent Nos. 6,311,415, 5,807,715, 4,816,567, and 4,816,397, which are incorporated herein by reference in their entirety. Chimeric antibodies comprising one or more CDRs from a non-human species and framework regions from a human immunoglobulin molecule can be produced using a variety of techniques known in the art including, for example, CDR-grafting (EP 239,400; International Publication No. WO 91/09967; and U.S. Patent Nos. 5,225,539, 5,530,101, and 5,585,089), veneering or resurfacing (EP 592,106; EP 519,596; Padlan, 1991, Molecular Immunology 28(4/5):489-498; Studnicka et al., 1994, Protein Engineering 7:805; and Roguska et al., 1994, PNAS 91:969), and chain shuffling (U.S. Patent No. 5,565,332).

[00305] Often, framework residues in the framework regions will be substituted with the corresponding residue from the CDR donor antibody to alter, preferably improve, antigen binding. These framework substitutions are identified by methods well known in the art, e.g., by modeling of the interactions of the CDR and framework residues to identify framework residues important for antigen binding and sequence comparison to identify unusual framework residues at particular positions. (See, e.g., U.S. Patent No. 5,585,089; and Riechmann et al., 1988, Nature 332:323, which are incorporated herein by reference in their entireties.)

A humanized antibody is an antibody or its variant or fragment thereof which is [00306] capable of binding to a predetermined antigen and which comprises a framework region having substantially the amino acid sequence of a human immunoglobulin and a CDR having substantially the amino acid sequence of a non-human immunoglobulin. A humanized antibody comprises substantially all of at least one, and typically two, variable domains in which all or substantially all of the CDR regions correspond to those of a non-human immunoglobulin (i.e., donor antibody) and all or substantially all of the framework regions are those of a human immunoglobulin consensus sequence. Preferably, a humanized antibody also comprises at least a portion of an immunoglobulin constant region (Fc), typically that of a human immunoglobulin. Ordinarily, the antibody will contain both the light chain as well as at least the variable domain of a heavy chain. The antibody also may include the CH1, hinge, CH2, CH3, and CH4 regions of the heavy chain. The humanized antibody can be selected from any class of immunoglobulins, including IgM, IgG, IgD, IgA and IgE, and any isotype, including IgG1, IgG2, IgG₃ and IgG₄. Usually the constant domain is a complement fixing constant domain where it is desired that the humanized antibody exhibit cytotoxic activity, and the class is typically IgG1. Where such cytotoxic activity is not desirable, the constant domain may be of the IgG2 class. The humanized antibody may comprise sequences from more than one class or isotype, and selecting particular constant domains to optimize desired effector functions is within the ordinary skill in the art. The framework and CDR regions of a humanized antibody need not correspond precisely to the parental sequences, e.g., the donor CDR or the consensus framework may be mutagenized by substitution, insertion or deletion of at least one residue so that the CDR or framework residue at that site does not correspond to either the consensus or the import antibody. Such mutations, however, will not be extensive. Usually, at least 75% of the humanized antibody residues will correspond to those of the parental framework region (FR) and CDR sequences, more often 90%, and most preferably greater than 95%. Humanized antibodies can be produced using variety of techniques known in the art, including but not limited to, CDR-grafting (European Patent No. EP 239,400; International Publication No. WO 91/09967; and U.S. Patent Nos. 5,225,539, 5,530,101, and 5,585,089), veneering or resurfacing (European Patent Nos. EP 592,106 and EP 519,596; Padlan, 1991, Molecular Immunology 28(4/5):489-498; Studnicka et al., 1994, Protein Engineering 7(6):805-814; and Roguska et al., 1994, PNAS 91:969-973), chain shuffling (U.S. Patent No. 5,565,332), and techniques disclosed in, e.g., U.S. Patent Nos. 6,407,213, 5,766,886, 5,585,089, International Publication No. WO 9317105, Tan et al., 2002, J. Immunol. 169:1119-25, Caldas et al., 2000, Protein Eng. 13:353-60, Morea et al., 2000, Methods 20:267-79, Baca et al., 1997, J. Biol. Chem. 272:10678-84, Roguska et al., 1996, Protein Eng. 9:895-904, Couto et al., 1995, Cancer Res. 55

(23 Supp):5973s-5977s, Couto et al., 1995, Cancer Res. 55:1717-22, Sandhu, 1994, Gene 150:409-10, Pedersen et al., 1994, J. Mol. Biol. 235:959-73, Jones et al., 1986, Nature 321:522-525, Riechmann et al., 1988, Nature 332:323, and Presta, 1992, Curr. Op. Struct. Biol. 2:593-596. Often, framework residues in the framework regions will be substituted with the corresponding residue from the CDR donor antibody to alter, preferably improve, antigen binding. These framework substitutions are identified by methods well known in the art, e.g., by modeling of the interactions of the CDR and framework residues to identify framework residues important for antigen binding and sequence comparison to identify unusual framework residues at particular positions. (See, e.g., Queen et al., U.S. Patent No. 5,585,089; and Riechmann et al., 1988, Nature 332:323, which are incorporated herein by reference in their entireties.)

[00307] Further, the antibodies of the invention can, in turn, be utilized to generate antiidiotype antibodies using techniques well known to those skilled in the art. (See, e.g.,
Greenspan & Bona, 1989, FASEB J. 7:437-444; and Nissinoff, 1991, J. Immunol. 147:24292438). The invention provides methods employing the use of polynucleotides comprising a
nucleotide sequence encoding an antibody of the invention or a fragment thereof.

5.4.1 Recombinant Expression Of An Antibody

Recombinant expression of an antibody used in the invention, a derivative, [00308] analog or fragment thereof, (e.g., a heavy or light chain of an antibody of the invention or a portion thereof or a single chain antibody of the invention), requires construction of an expression vector containing a polynucleotide that encodes the antibody. Once a polynucleotide encoding an antibody molecule or a heavy or light chain of an antibody, or portion thereof (preferably, but not necessarily, containing the heavy or light chain variable domain), of the invention has been obtained, the vector for the production of the antibody molecule may be produced by recombinant DNA technology using techniques well known in the art. Thus, methods for preparing a protein by expressing a polynucleotide containing an antibody encoding nucleotide sequence are described herein. Methods which are well known to those skilled in the art can be used to construct expression vectors containing antibody coding sequences and appropriate transcriptional and translational control signals. These methods include, for example, in vitro recombinant DNA techniques, synthetic techniques, and in vivo genetic recombination. The invention, thus, provides replicable vectors comprising a nucleotide sequence encoding an antibody molecule of the invention, a heavy or light chain of an antibody, a heavy or light chain variable domain of an antibody or a portion thereof, or a heavy or light chain CDR, operably linked to a promoter. Such vectors may include the nucleotide sequence encoding the constant region of the antibody molecule (see, e.g., International Publication Nos.

WO 86/05807 and WO 89/01036; and U.S. Patent No. 5,122,464) and the variable domain of the antibody may be cloned into such a vector for expression of the entire heavy, the entire light chain, or both the entire heavy and light chains.

[00309] The expression vector is transferred to a host cell by conventional techniques and the transfected cells are then cultured by conventional techniques to produce an antibody of the invention. Thus, the invention includes host cells containing a polynucleotide encoding an antibody of the invention or fragments thereof, or a heavy or light chain thereof, or portion thereof, or a single chain antibody of the invention, operably linked to a heterologous promoter. In embodiments for the expression of double-chained antibodies, vectors encoding both the heavy and light chains may be co-expressed in the host cell for expression of the entire immunoglobulin molecule, as detailed below.

A variety of host-expression vector systems may be utilized to express the [00310] antibody molecules of the invention (see, e.g., U.S. Patent No. 5,807,715). Such host-expression systems represent vehicles by which the coding sequences of interest may be produced and subsequently purified, but also represent cells which may, when transformed or transfected with the appropriate nucleotide coding sequences, express an antibody molecule of the invention in situ. These include but are not limited to microorganisms such as bacteria (e.g., E. coli and B. subtilis) transformed with recombinant bacteriophage DNA, plasmid DNA or cosmid DNA expression vectors containing antibody coding sequences; yeast (e.g., Saccharomyces Pichia) transformed with recombinant yeast expression vectors containing antibody coding sequences; insect cell systems infected with recombinant virus expression vectors (e.g., baculovirus) containing antibody coding sequences; plant cell systems infected with recombinant virus expression vectors (e.g., cauliflower mosaic virus, CaMV; tobacco mosaic virus, TMV) or transformed with recombinant plasmid expression vectors (e.g., Ti plasmid) containing antibody coding sequences; or mammalian cell systems (e.g., COS, CHO, BHK, 293, NS0, and 3T3 cells) harboring recombinant expression constructs containing promoters derived from the genome of mammalian cells (e.g., metallothionein promoter) or from mammalian viruses (e.g., the adenovirus late promoter; the vaccinia virus 7.5K promoter). Preferably, bacterial cells such as Escherichia coli, and more preferably, eukaryotic cells, especially for the expression of whole recombinant antibody molecule, are used for the expression of a recombinant antibody molecule. For example, mammalian cells such as Chinese hamster ovary cells (CHO), in conjunction with a vector such as the major intermediate early gene promoter element from human cytomegalovirus is an effective expression system for antibodies (Foecking et al., 1986, Gene

45:101; and Cockett et al., 1990, *BioTechnology* 8:2). In a specific embodiment, the expression of nucleotide sequences encoding antibodies or fragments thereof which immunospecifically bind to and agonize is regulated by a constitutive promoter, inducible promoter or tissue specific promoter.

In bacterial systems, a number of expression vectors may be advantageously [00311] selected depending upon the use intended for the antibody molecule being expressed. For example, when a large quantity of such a protein is to be produced, for the generation of pharmaceutical compositions of an antibody molecule, vectors which direct the expression of high levels of fusion protein products that are readily purified may be desirable. Such vectors include, but are not limited to, the E. coli expression vector pUR278 (Ruther et al., 1983, EMBO 12:1791), in which the antibody coding sequence may be ligated individually into the vector in frame with the lac Z coding region so that a fusion protein is produced; pIN vectors (Inouye & Inouye, 1985, Nucleic Acids Res. 13:3101-3109; Van Heeke & Schuster, 1989, J. Biol. Chem. 24:5503-5509); and the like. pGEX vectors may also be used to express foreign polypeptides as fusion proteins with glutathione 5-transferase (GST). In general, such fusion proteins are soluble and can easily be purified from lysed cells by adsorption and binding to matrix glutathione-agarose beads followed by elution in the presence of free glutathione. The pGEX vectors are designed to include thrombin or factor Xa protease cleavage sites so that the cloned target gene product can be releastd from the GST moiety.

[00312] In an insect system, *Autographa californica* nuclear polyhedrosis virus (AcNPV) is used as a vector to express foreign genes. The virus grows in *Spodoptera frugiperda* cells. The antibody coding sequence may be cloned individually into non-essential regions (for example the polyhedrin gene) of the virus and placed under control of an AcNPV promoter (for example the polyhedrin promoter).

[00313] In mammalian host cells, a number of viral-based expression systems may be utilized. In cases where an adenovirus is used as an expression vector, the antibody coding sequence of interest may be ligated to an adenovirus transcription/translation control complex, e.g., the late promoter and tripartite leader sequence. This chimeric gene may then be inserted in the adenovirus genome by *in vitro* or *in vivo* recombination. Insertion in a non-essential region of the viral genome (e.g., region El or E3) will result in a recombinant virus that is viable and capable of expressing the antibody molecule in infected hosts (e.g., see Logan & Shenk, 1984, PNAS 8 1:355-359). Specific initiation signals may also be required for efficient translation of inserted antibody coding sequences. These signals include the ATG initiation codon and

adjacent sequences. Furthermore, the initiation codon must be in phase with the reading frame of the desired coding sequence to ensure translation of the entire insert. These exogenous translational control signals and initiation codons can be of a variety of origins, both natural and synthetic. The efficiency of expression may be enhanced by the inclusion of appropriate transcription enhancer elements, transcription terminators, etc. (see, *e.g.*, Bittner et al., 1987, *Methods in Enzymol.* 153:516-544).

In addition, a host cell strain may be chosen which modulates the expression of the inserted sequences, or modifies and processes the gene product in the specific fashion desired. Such modifications (*e.g.*, glycosylation) and processing (*e.g.*, cleavage) of protein products may be important for the function of the protein. Different host cells have characteristic and specific mechanisms for the post-translational processing and modification of proteins and gene products. Appropriate cell lines or host systems can be chosen to ensure the correct modification and processing of the foreign protein expressed. To this end, eukaryotic host cells which possess the cellular machinery for proper processing of the primary transcript, glycosylation, and phosphorylation of the gene product may be used. Such mammalian host cells include but are not limited to CHO, VERO, BHK, HeLa, COS, MDCK, 293, 3T3, W138, BT483, Hs578T, HTB2, BT2O, NS1, T47D, NS0 (a murine myeloma cell line that does not endogenously produce any immunoglobulin chains), CRL7O3O and HsS78Bst cells.

[00315] The antibodies comprising at least one zero-order thioether can be recombinantly produced by any cell lines for producing antibodies known to those skilled in the art. It has been found that it is advantageous to produce the antibodies of the invention in melanoma cells. In certain embodiments, the antibodies of the invention are recombinantly produced in melanoma cells. In some embodiments, the antibodies of the invention are not recombinantly produced in CHO cell line. In other embodiments, the antibodies of the invention are not recombinantly produced in NS0 cell line.

[00316] For long-term, high-yield production of recombinant proteins, stable expression is preferred. For example, cell lines which stably express the antibody molecule may be engineered. Rather than using expression vectors which contain viral origins of replication, host cells can be transformed with DNA controlled by appropriate expression control elements (e.g., promoter, enhancer, sequences, transcription terminators, polyadenylation sites, etc.), and a selectable marker. Following the introduction of the foreign DNA, engineered cells may be allowed to grow for 1-2 days in an enriched media, and then are switched to a selective media. The selectable marker in the recombinant plasmid confers resistance to the selection and allows

cells to stably integrate the plasmid into their chromosomes and grow to form foci which in turn can be cloned and expanded into cell lines. This method may advantageously be used to engineer cell lines which express the antibody molecule. Such engineered cell lines may be particularly useful in screening and evaluation of compositions that interact directly or indirectly with the antibody molecule.

A number of selection systems may be used, including but not limited to, the [00317] herpes simplex virus thymidine kinase (Wigler et al., 1977, Cell 11:223), glutamine synthase, hypoxanthine guanine phosphoribosyltransferase (Szybalska & Szybalski, 1992, Proc. Natl. Acad. Sci. USA 48:202), and adenine phosphoribosyltransferase (Lowy et al., 1980, Cell 22:8-17) genes can be employed in tk-, gs-, hgprt- or aprt- cells, respectively. Also, antimetabolite resistance can be used as the basis of selection for the following genes: dhfr, which confers resistance to methotrexate (Wigler et al., 1980, PNAS 77:357; O'Hare et al., 1981, PNAS 78:1527); gpt, which confers resistance to mycophenolic acid (Mulligan & Berg, 1981, PNAS 78:2072); neo, which confers resistance to the aminoglycoside G-418 (Wu and Wu, 1991, Biotherapy 3:87; Tolstoshev, 1993, Ann. Rev. Pharmacol. Toxicol. 32:573; Mulligan, 1993, Science 260:926; and Morgan and Anderson, 1993, Ann. Rev. Biochem. 62: 191; May, 1993, TIB TECH 11:155-); and hygro, which confers resistance to hygromycin (Santerre et al., 1984, Gene 30:147). Methods commonly known in the art of recombinant DNA technology may be routinely applied to select the desired recombinant clone, and such methods are described, for example, in Ausubel et al. (eds.), Current Protocols in Molecular Biology, John Wiley & Sons, NY (1993); Kriegler, Gene Transfer and Expression, A Laboratory Manual, Stockton Press, NY (1990); and in Chapters 12 and 13, Dracopoli et al. (eds), Current Protocols in Human Genetics, John Wiley & Sons, NY (1994); Colberre-Garapin et al., 1981, J. Mol. Biol. 150:1, which are incorporated by reference herein in their entireties.

[00318] The expression levels of an antibody molecule can be increased by vector amplification (for a review, see Bebbington and Hentschel, The use of vectors based on gene amplification for the expression of cloned genes in mammalian cells in DNA cloning, Vol.3. (Academic Press, New York, 1987)). When a marker in the vector system expressing antibody is amplifiable, increase in the level of inhibitor present in culture of host cell will increase the number of copies of the marker gene. Since the amplified region is associated with the antibody gene, production of the antibody will also increase (Crouse et al., 1983, *Mol. Cell. Biol.* 3:257).

[00319] The host cell may be co-transfected with two expression vectors of the invention, the first vector encoding a heavy chain derived polypeptide and the second vector encoding a

light chain derived polypeptide. The two vectors may contain identical selectable markers which enable equal expression of heavy and light chain polypeptides. Alternatively, a single vector may be used which encodes, and is capable of expressing, both heavy and light chain polypeptides. In such situations, the light chain should be placed before the heavy chain to avoid an excess of toxic free heavy chain (Proudfoot, 1986, *Nature* 322:52; and Kohler, 1980, *PNAS* 77:2197). The coding sequences for the heavy and light chains may comprise cDNA or genomic DNA.

[00320] Once an antibody molecule of the invention has been produced by recombinant expression, it may be purified by any method known in the art for purification of an immunoglobulin molecule, for example, by chromatography (e.g., ion exchange, affinity, particularly by affinity for the specific antigen after Protein A, and sizing column chromatography), centrifugation, differential solubility, or by any other standard technique for the purification of proteins. Further, the antibodies of the present invention or fragments thereof may be fused to heterologous polypeptide sequences described herein or otherwise known in the art to facilitate purification.

5.5 Use of the antibodies and compositions of the present invention

[00321] The formulations comprising antibodies and compositions thereof can be used in any context that those of skilled in the art recognize. For example, the formulations of the invention can be used directly against a particular antigen. The formulations of the invention comprising antibodies and compositions can also be used in diagnostic assays either in vivo or in vitro for detection/identification of the expression of an antigen in a subject or a biological sample (e.g., cells or tissues). formulations of the invention comprising antibodies and compositions can be used alone or in combination with other prophylactic or therapeutic agents for treating, managing, preventing or ameliorating a disorder or one or more symptoms thereof.

[00322] The present invention provides methods for preventing, managing, treating, or ameliorating a disorder comprising administering to a subject in need thereof one or more antibodies of the invention alone or in combination with one or more therapies (e.g., one or more prophylactic or therapeutic agents) other than an antibody of the invention. The present invention also provides formulations comprising one or more antibodies of the invention and one or more prophylactic or therapeutic agents other than antibodies of the invention and methods of preventing, managing, treating, or ameliorating a disorder or one or more symptoms thereof utilizing said compositions. Therapeutic or prophylactic agents include, but are not limited to, small molecules, synthetic drugs, peptides, polypeptides, proteins, nucleic acids (e.g.,

DNA and RNA nucleotides including, but not limited to, antisense nucleotide sequences, triple helices, RNAi, and nucleotide sequences encoding biologically active proteins, polypeptides or peptides) antibodies, synthetic or natural inorganic molecules, mimetic agents, and synthetic or natural organic molecules.

Any therapy which is known to be useful, or which has been used or is currently [00323] being used for the prevention, management, treatment, or amelioration of a disorder or one or more symptoms thereof can be used in combination with an antibody or a composition of the invention in accordance with the invention described herein. See, e.g., Gilman et al., Goodman and Gilman's: The Pharmacological Basis of Therapeutics, 10th ed., McGraw-Hill, New York, 2001; The Merck Manual of Diagnosis and Therapy, Berkow, M. D. et al. (eds.), 17th Ed., Merck Sharp & Dohme Research Laboratories, Rahway, N.J., 1999; Cecil Textbook of Medicine, 20th Ed., Bennett and Plum (eds.), W. B. Saunders, Philadelphia, 1996 for information regarding therapies (e.g., prophylactic or therapeutic agents) which have been or are currently being used for preventing, treating, managing, or ameliorating a disorder or one or more symptoms thereof. Examples of such agents include, but are not limited to, immunomodulatory agents, anti-inflammatory agents (e.g., adrenocorticoids, corticosteroids (e.g., beclomethasone, budesonide, flunisolide, fluticasone, triamcinolone, methlyprednisolone, prednisolone, prednisone, hydrocortisone), glucocorticoids, steroids, non-steriodal antiinflammatory drugs (e.g., aspirin, ibuprofen, diclofenac, and COX-2 inhibitors), anti-cancer agents, pain relievers, leukotreine antagonists (e.g., montelukast, methyl xanthines, zafirlukast, and zileuton), beta2-agonists (e.g., albuterol, biterol, fenoterol, isoetharie, metaproterenol, pirbuterol, salbutamol, terbutalin formoterol, salmeterol, and salbutamol terbutaline), anticholinergic agents (e.g., ipratropium bromide and oxitropium bromide), sulphasalazine, penicillamine, dapsone, antihistamines, anti-malarial agents (e.g., hydroxychloroquine), antiviral agents, and antibiotics (e.g., dactinomycin (formerly actinomycin), bleomycin, erythomycin, penicillin, mithramycin, and anthramycin (AMC)).

In a specific embodiment, the present invention provides a method comprising administering a formulation comprising one or more humanized anti-IL-9 antibodies to a subject, preferably a human subject, for preventing, treating, managing, or ameliorating a respiratory condition or one or more symptoms thereof. In one embodiment, the invention encompasses a method of preventing, treating, managing, or ameliorating a respiratory disorder or one or more symptoms thereof (e.g., an allergy, wheezing, and asthma), said method comprising administering to a subject in need thereof a dose of a prophylactically or

therapeutically effective amount of a formulation comprising one or more of humanized anti-IL-9 antibodies. In another embodiment, the invention provides a method of preventing, treating, managing, or ameliorating a respiratory infection or one or more symptoms thereof, said method comprising administering a prophylactically or therapeutic effective amount of one or more humanized anti-IL-9 antibodies.

In a specific embodiment, the present invention provides a method comprising administering a formulation of one or more humanized anti-EphA2 antibodies to a subject, preferably a human subject, for preventing, treating, managing, or ameliorating a hyperproliferative cell disease or one or more symptoms thereof. In one embodiment, one or more humanized anti-EphA2 antibodies are administered alone or in combination with other agents to a subject to prevent, treat, manage, or ameliorate cancer or one or more symptoms thereof (see, e.g., U.S. application Ser. No. 10/436,782, which is incorporated herein by reference in its entirety). In another embodiment, one or more humanized anti-EphA2 antibodies are administered alone or in combination with other agents to a subject to prevent, treat, manage, or ameliorate a disorder involving non-neoplastic hyperproliferative cells, in particular hyperproliferative epithlial and endothelial cells, or one or symptoms thereof (see e.g., U.S. Application Ser. No. 60/462,024, which is incorporated herein by reference in its entirety). In yet another embodiment, one or more humanized anti-EphA2 antibodies are used for diagnostic or screening purposes.

[00326] The formulations comprising antibodies and compositions of the invention can be used directly against a particular antigen. In some embodiments, the antibodies and compositions of the invention belong to a subclass or isotype that is capable of mediating the lysis of cells to which the antibody binds. In a specific embodiment, the antibodies of the invention belong to a subclass or isotype that, upon complexing with cell surface proteins, activates serum complement and/or mediates antibody dependent cellular cytotoxicity (ADCC) by activating effector cells such as natural killer cells or macrophages.

[00327] The biological activities of antibodies are known to be determined, to a large extent, by the constant domains or Fc region of the antibody molecule (Uananue and Benacerraf, Textbook of Immunology, 2nd Edition, Williams & Wilkins, p. 218 (1984)). This includes their ability to activate complement and to mediate antibody-dependent cellular cytotoxicity (ADCC) as effected by leukocytes. Antibodies of different classes and subclasses differ in this respect, as do antibodies from the same subclass but different species; according to the present invention, antibodies of those classes having the desired biological activity are prepared. Preparation of

these antibodies involves the selection of antibody constant domains and their incorporation in the humanized antibody by known technique. For example, mouse immunoglobulins of the IgG3 and IgG2a class are capable of activating serum complement upon binding to the target cells which express the cognate antigen, and therefore humanized antibodies which incorporate IgG3 and IgG2a effector functions are desirable for certain therapeutic applications.

[00328] In some embodiments, formulations of the invention comprising antibodies and compositions are useful in passively immunizing patients.

The formulations of the invention comprising antibodies and compositions can also be used in diagnostic assays either in vivo or in vitro for detection/identification of the expression of an antigen in a subject or a biological sample (*e.g.*, cells or tissues). Non-limiting examples of using an antibody, or a composition comprising an antibody in a diagnostic assay are given in U.S. Pat. Nos. 6,392,020; 6,156,498; 6,136,526; 6,048,528; 6,015,555; 5,833,988; 5,811,310; 8 5,652,114; 5,604,126; 5,484,704; 5,346,687; 5,318,892; 5,273,743; 5,182,107; 5,122,447; 5,080,883; 5,057,313; 4,910,133; 4,816,402; 4,742,000; 4,724,213; 4,724,212; 4,624,846; 4,623,627; 4,618,486; 4,176,174 (all of which are incorporated herein by reference). Suitable diagnostic assays for the antigen and its antibodies depend on the particular antibody used. Non-limiting examples are an ELISA, sandwich assay, and steric inhibition assays. For in vivo diagnostic assays using the antibodies of the invention, the antibodies may be conjugated to a label that can be detected by imaging techniques, such as X-ray, computed tomography (CT), ultrasound, or magnetic resonance imaging (MRI). The antibodies of the invention can also be used for the affinity purification of the antigen from recombinant cell culture or natural sources.

5.5.1 <u>Prophylactic and Therapeutic Use of Formulations</u> of Antibodies against RSV infections

[00330] The present invention provides antibody-based therapies which involve administering antibodies of the invention to a subject, preferably a human, for preventing, treating, or ameliorating a RSV infection (*i.e.*, an upper and/or lower respiratory tract RSV infection), otitis media (stemming from, caused by, or associated with a RSV infection), or a symptom or respiratory condition relating thereto (including, but not limited to, asthma, wheezing, RAD, or a combination thereof). Prophylactic and therapeutic agents of the invention include, but are not limited to, antibodies (including analogs and derivatives thereof as described herein) and nucleic acids encoding antibodies (including analogs and derivatives thereof and anti-idiotypic antibodies as described herein). Antibodies may be provided in pharmaceutically acceptable compositions as known in the art or as described herein.

[00331] Formulations of the present invention comprising antibodies that function as antagonists of a RSV infection can be administered to a subject, preferably a human, to treat, prevent or ameliorate an upper and/or lower respiratory tract RSV infection, otitis media (preferably, stemming from, caused by, or associated with a RSV infection), or a symptom or respiratory condition relating thereto (including, but not limited to, asthma, wheezing, RAD, or a combination thereof). For example, antibodies which disrupt or prevent the interaction between a RSV antigen and its host cell receptor may be administered to subject, preferably a human, to treat, prevent or ameliorate an upper and/or lower respiratory tract RSV infection, otitis media (stemming from, caused by, or associated with a RSV infection), or a symptom or respiratory condition relating thereto (including, but not limited to, asthma, wheezing, RAD, or a combination thereof).

In a specific embodiment, an antibody prevents or inhibits RSV from binding to its host cell receptor by at least 99%, at least 95%, at least 90%, at least 85%, at least 80%, at least 75%, at least 70%, at least 60%, at least 50%, at least 45%, at least 40%, at least 45%, at least 35%, at least 30%, at least 25%, at least 20%, or at least 10% relative to RSV binding to its host cell receptor in the absence of said antibodies or in the presence of a negative control in an assay known to one of skill in the art or described herein. In another embodiment, a combination of antibodies prevent or inhibit RSV from binding to its host cell receptor by at least 99%, at least 95%, at least 90%, at least 85%, at least 80%, at least 75%, at least 70%, at least 60%, at least 50%, at least 45%, at least 45%, at least 35%, at least 30%, at least 25%, at least 20%, or at least 10% relative to RSV binding to its host cell receptor in the absence of said antibodies or in the presence of a negative control in an assay known to one of skill in the art or described herein.

[00333] In a specific embodiment, an antibody prevents or inhibits RSV-induced fusion by at least 99%, at least 95%, at least 90%, at least 85%, at least 80%, at least 75%, at least 70%, at least 60%, at least 50%, at least 45%, at least 45%, at least 35%, at least 30%, at least 25%, at least 20%, or at least 10% relative to RSV-induced fusion in the absence of said antibodies or in the presence of a negative control in an assay known to one of skill in the art or described herein. In another embodiment, a combination of antibodies prevent or inhibit RSV-induced fusion by at least 99%, at least 95%, at least 90%, at least 85%, at least 80%, at least 75%, at least 70%, at least 50%, at least 45%, at least 45%, at least 45%, at least 35%, at least 45%, at least 45

the absence of said antibodies or in the presence of a negative control in an assay known to one of skill in the art or described herein.

In a specific embodiment, an antibody prevents or inhibits RSV-induced fusion after viral attachment to cells by at least 99%, at least 95%, at least 90%, at least 85%, at least 80%, at least 75%, at least 70%, at least 60%, at least 50%, at least 45%, at least 40%, at least 45%, at least 35%, at least 30%, at least 25%, at least 20%, or at least 10% relative to RSV-induced fusion after viral attachment to cells in the absence of said antibodies or in the presence of a negative control in an assay known to one of skill in the art or described herein. In another embodiment, a combination of antibodies prevent or inhibit RSV-induced fusion after viral attachment to cells by at least 99%, at least 95%, at least 90%, at least 85%, at least 80%, at least 75%, at least 70%, at least 60%, at least 50%, at least 45%, at leas

[00335] Antibodies which do not prevent RSV from binding its host cell receptor but inhibit or downregulate RSV replication can also be administered to a subject to treat, prevent or ameliorate an upper and/or lower respiratory tract RSV infection, otitis media (stemming from, caused by, or associated with a RSV infection), or a symptom or respiratory condition relating thereto (including, but not limited to, asthma, wheezing, RAD, or a combination thereof). The ability of an antibody to inhibit or downregulate RSV replication may be determined by techniques described herein or otherwise known in the art. For example, the inhibition or downregulation of RSV replication can be determined by detecting the RSV titer in the lungs of a subject, preferably a human. In further embodiments, the inhibition or downregulation of RSV replication can be determined by detecting the amount of RSV in the nasal passages or in the middle ear by methods known in the art (e.g., Northern blot analysis, RT-PCR, Western Blot analysis, etc.).

In some embodiments, a formulations of the present invention comprises an antibody that results in reduction of about 1-fold, about 1.5-fold, about 2-fold, about 3-fold, about 4-fold, about 5-fold, about 8-fold, about 10-fold, about 15-fold, about 20-fold, about 25-fold, about 30-fold, about 35-fold, about 40-fold, about 45-fold, about 50-fold, about 55-fold, about 60-fold, about 65-fold, about 70-fold, about 75-fold, about 80-fold, about 85-fold, about 90-fold, about 95-fold, about 100-fold, about 105 fold, about 110-fold, about 115-fold, about 120 fold, about 125-fold or higher in RSV titer in the lung. The fold-reduction in RSV titer may

be as compared to a negative control (such as placebo), as compared to another treatment (including, but not limited to treatment with palivizumab), or as compared to the titer in the patient prior to antibody administration.

In a specific embodiment, formulation of the present invention comprises an antibody that inhibits or downregulates RSV replication by at least 99%, at least 95%, at least 90%, at least 85%, at least 80%, at least 75%, at least 70%, at least 60%, at least 50%, at least 45%, at least 35%, at least 30%, at least 25%, at least 20%, or at least 10% relative to RSV replication in absence of said antibodies or in the presence of a negative control in an assay known in the art or described herein. In another embodiment, a combination of antibodies inhibit or downregulate RSV replication by at least 99%, at least 95%, at least 90%, at least 85%, at least 80%, at least 75%, at least 70%, at least 60%, at least 50%, at least 45%, at least 40%, at least 45%, at least 35%, at least 30%, at least 25%, at least 20%, or at least 10% relative to RSV replication in absence of said antibodies or in the presence of a negative control in an assay known in the art or described herein.

In some embodiments, formulation of the present invention comprises an antibody that results in reduction of about 1-fold, about 1.5-fold, about 2-fold, about 3-fold, about 4-fold, about 5-fold, about 8-fold, about 10-fold, about 15-fold, about 20-fold, about 25-fold, about 30-fold, about 35-fold, about 40-fold, about 45-fold, about 50-fold, about 55-fold, about 60-fold, about 65-fold, about 70-fold, about 75-fold, about 80-fold, about 85-fold, about 90-fold, about 95-fold, about 100-fold, about 105 fold, about 110-fold, about 115-fold, about 120 fold, about 125-fold or higher in RSV titer in the upper respiratory tract. The fold-reduction in RSV titer may be as compared to a negative control (such as placebo), as compared to another treatment (including, but not limited to treatment with palivizumab), or as compared to the titer in the patient prior to antibody administration.

In other embodiments, formulation of the present invention comprises an antibody that results in reduction of about 1-fold, about 1.5-fold, about 2-fold, about 3-fold, about 4-fold, about 5-fold, about 8-fold, about 10-fold, about 15-fold, about 20-fold, about 25-fold, about 30-fold, about 35-fold, about 40-fold, about 45-fold, about 50-fold, about 55-fold, about 60-fold, about 65-fold, about 70-fold, about 75-fold, about 80-fold, about 85-fold, about 90-fold, about 95-fold, about 100-fold, about 105 fold, about 110-fold, about 115-fold, about 120 fold, about 125-fold or higher in RSV titer in the lower respiratory tract. The fold-reduction in RSV titer may be as compared to a negative control (such as placebo), as compared to another

treatment (including, but not limited to treatment with palivizumab), or as compared to the titer in the patient prior to antibody administration.

[00340] One or more antibodies in connection with the present invention that immunospecifically bind to one or more RSV antigens may be used locally or systemically in the body as a prophylactic or therapeutic agent. The antibodies may also be advantageously utilized in combination with other monoclonal or chimeric antibodies, or with lymphokines or hematopoietic growth factors (such as, e.g., IL-2, IL-3 and IL-7), which, for example, serve to increase the number or activity of effector cells which interact with the antibodies. The antibodies may also be advantageously utilized in combination with other monoclonal or chimeric antibodies, or with lymphokines or hematopoietic growth factors (such as, e.g., IL-2, IL-3 and IL-7), which, for example, serve to increase the immune response. The antibodies may also be advantageously utilized in combination with one or more drugs used to treat RSV infection such as, for example anti-viral agents. Antibodies of the invention may be used in combination with one or more of the following drugs: NIH-351 (Gemini Technologies), recombinant RSV vaccine (Aviron), RSVf-2 (Intracel), F-50042 (Pierre Fabre), T-786 (Trimeris), VP-36676 (ViroPharma), RFI-641 (American Home Products), VP-14637 (ViroPharma), PFP-1 and PFP-2 (American Home Products), RSV vaccine (Avant Immunotherapeutics), and F-50077 (Pierre Fabre). In a specific embodiment, an effective amount of an antibody and an effective amount of another therapy is used.

[00341] The formulations of the invention comprising antibodies may be administered alone or in combination with other types of therapies (e.g., hormonal therapy, immunotherapy, and anti-inflammatory agents). Generally, administration of products of a species origin or species reactivity (in the case of antibodies) that is the same species as that of the patient is preferred. Thus, in a preferred embodiment, human or humanized antibodies, derivatives, analogs, or nucleic acids, are administered to a human patient for therapy or prophylaxis.

[00342] In specific embodiments, an antibody is administered in combination with one or more anti-IL-9 antibodies (such as those disclosed in U.S. Publication No. 2005/0002934) either alone or in combination with one or more antibodies of the invention and/or other types of therapies or other agents (e.g., hormone therapy, immunotherapy, and anti-inflammatory agents, such as those disclosed in U.S. Publication No. 2005/0002934, which is herein incorporated by reference in its entirety).

[00343] It is preferred to use high affinity and/or potent *in vivo* inhibiting antibodies and/or neutralizing antibodies that immunospecifically bind to a RSV antigen, for both immunoassays directed to RSV, and the prevention, management or treatment of an upper and/or lower respiratory tract RSV infection, otitis media (preferably, stemming from, caused by, or associated with a RSV infection), or a symptom or respiratory condition relating thereto (including, but not limited to, asthma, wheezing, RAD, or a combination thereof). It is also preferred to use polynucleotides encoding high affinity and/or potent *in vivo* inhibiting antibodies and/or neutralizing antibodies that immunospecifically bind to a RSV antigen, for both immunoassays directed to RSV and therapy for an upper and/or lower respiratory tract RSV infection, otitis media (stemming from, caused by, or associated with a RSV infection), or a symptom or respiratory condition relating thereto (including, but not limited to, asthma, wheezing, RAD, or a combination thereof). Such antibodies will preferably have an affinity for the RSV F glycoprotein and/or fragments of the F glycoprotein.

The methods of the invention comprise the administration of one or more [00344] antibodies to patients suffering from or expected to suffer from (e.g., patients with a genetic predisposition for or patients that have previously suffered from) an upper and/or lower respiratory tract RSV infection, otitis media (preferably, stemming from, caused by, or associated with a RSV infection), or a symptom or respiratory condition relating thereto (including, but not limited to, asthma, wheezing, RAD, or a combination thereof). Such patients may have been previously treated or are currently being treated for the infection, otitis media, or a respiratory condition, e.g., with a therapy other than an antibody of the invention. In one embodiment, the methods of the invention comprise the administration of one or more antibodies to patients that are immunocompromised or immunosuppressed. In a certain embodiment, an antibody administered to patients that are immunocompromised or immunosuppressed. In another embodiment, an antibody is administered to a human with cystic fibrosis, bronchopulmonary dysplasia, congenital heart disease, congenital immunodeficiency or acquired immunodeficiency, or to a human who has had a bone marrow transplant to treat, prevent or ameliorate one or more symptoms associated with an upper and/or lower respiratory tract RSV infection or otitis media (preferably, stemming from, caused by, or associated with a RSV infection), or a symptom or respiratory condition relating thereto (including, but not limited to, asthma, wheezing, RAD, or a combination thereof). In another embodiment, a formulation of the invention comprising an antibody is administered to a human infant, preferably a human infant born prematurely or a human infant at risk of hospitalization for RSV infection to treat, prevent or ameliorate one or more symptoms associated with an upper and/or

lower respiratory tract RSV infection or otitis media. In yet another embodiment, a formulation of the invention comprising an antibody is administered to the elderly or people in group homes (e.g., nursing homes or rehabilitation centers). In accordance with the invention, a formulation of the invention comprising an antibody may be used as any line of therapy, including, but not limited to, a first, second, third and fourth line of therapy, including, but not limited to, a first, second, third and fourth line of therapy. Further, in accordance with the invention, a formulation of the invention comprising an antibody can be used before any adverse effects or intolerance of the therapies other than an antibody occurs. The invention encompasses methods for administering one or more antibodies to prevent the onset or recurrence of an upper and/or lower respiratory tract RSV infection or otitis media.

[00345] In one embodiment, the invention also provides methods of treatment, management, prevention and/or amelioration of an upper and/or lower respiratory tract RSV infection (preferably stemming from, caused by, or associated with a RSV infection), otitis media or a symptom or respiratory condition related thereto (including, but not limited to, asthma, wheezing, RAD, or a combination thereof) as alternatives to current therapies. In a specific embodiment, the current therapy has proven or may prove too toxic (i.e., results in unacceptable or unbearable side effects) for the patient. In another embodiment, an antibody decreases the side effects as compared to the current therapy. In another embodiment, the patient has proven refractory to a current therapy. In such embodiments, the invention provides for the administration of one or more antibodies of the invention without any other anti-infection therapies. In certain embodiments, one or more antibodies can be administered to a patient in need thereof instead of another therapy to treat an upper and/or lower respiratory tract RSV infection, otitis media or a symptom or respiratory condition related thereto (including, but not limited to, asthma, wheezing, RAD, or a combination thereof). In one embodiment, the invention provides methods of treating, managing, preventing and/or ameliorating an active upper and/or lower respiratory tract RSV infection, otitis media or a symptom or respiratory condition related thereto (including, but not limited to, asthma, wheezing, RAD, or a combination thereof).

[00346] The present invention also encompasses methods for administering one or more antibodies to treat or ameliorate symptoms of an upper and/or lower respiratory tract RSV infection or otitis media in patients that are or have become refractory to therapies other than the antibodies. The determination whether the infection is refractory can be made either *in vivo* or *in vitro* by any method known in the art for assaying the effectiveness of a therapy on affected

cells in the infection, particularly epithelial cells, or in patients that are or have become refractory to therapies other than antibodies of the invention.

[00347] In certain embodiments, an effective amount of one or more antibodies in the formulation of the invention is administered in combination with one or more supportive measures to a subject in need thereof to prevent, manage, treat, and/or ameliorate an upper and/or lower respiratory tract RSV infection, otitis media (preferably, stemming from, caused by, or associated with a RSV infection), or a symptom or respiratory condition relating thereto (including, but not limited to, asthma, wheezing, RAD, or a combination thereof). Non-limiting examples of supportive measures include humidification of the air by an ultrasonic nebulizer, aerolized recemic epinephrine, oral dexamethasone, intravenous fluids, intubation, fever reducers (e.g., ibuprofen, acetometaphin), and antibiotic and/or anti-fungal therapy (i.e., to prevent or treat secondary bacterial and/or fungal infections).

[00348] In a specific embodiment, the invention provides methods for preventing, treating, managing, and/or ameliorating a RSV infection (*i.e.*, an upper and/or lower respiratory tract RSV infection), otitis media (preferably, stemming from, caused by, or associated with a RSV infection), or a symptom or respiratory condition relating thereto (including, but not limited to, asthma, wheezing, RAD, or a combination thereof), said methods comprising administering to a subject in need thereof an effective amount of one or more antibodies of the invention alone or in combination with one or more anti-viral agents such as, but not limited to, amantadine, rimantadine, oseltamivir, znamivir, ribavarin, RSV-IVIG (*i.e.*, intravenous immune globulin infusion) (RESPIGAMTM), EphA2/EphrinA1 Modulators, and/or an anti-IL-9 antibody (see, *e.g.*, U.S. Publication No. 2005/0002934).

[00349] In a specific embodiment, the invention provides methods for preventing, treating, managing, and/or ameliorating one or more secondary responses to a primary viral infection, said methods comprising administering an effective amount of one or more antibodies alone or in combination with an effective amount of other therapies (e.g., other prophylactic or therapeutic agents). Examples of secondary responses to a primary viral infection include, but are not limited to, asthma-like responsiveness to mucosal stimula, elevated total respiratory resistance, increased susceptibility to secondary viral, bacterial, and fungal infections, and development of conditions such as, but not limited to, bronchiolitis, pneumonia, croup, and febrile bronchitis.

[00350] In a specific embodiment, the invention provides methods of preventing, treating, managing, and/or ameliorating a RSV infection (i.e., an upper and/or lower respiratory tract RSV infection), otitis media (preferably, stemming from, caused by or associated with a RSV infection, such as an upper and/or lower respiratory tract RSV infection) or a symptom or respiratory condition relating thereto (including, but not limited to asthma, wheezing, RAD, or a combination thereof), said methods comprising administering to a subject in need thereof an effective amount of one or more antibodies in combination with an effective amount of an EphA2/EphrinA1 Modulator (U.S. Provisional Appn. Serial No. 60/622,489, filed October 27, 2004, entitled "Use of Modulators of EphA2 and EphrinA1 for the Treatment and Prevention of Infections", which is incorporated by reference herein in its entirety). In another specific embodiment, the invention provides methods for preventing, treating, managing, and/or ameliorating a RSV infection (i.e., an upper and/or lower respiratory tract RSV infection), otitis media (preferably, stemming from, caused by or associated with a RSV infection, such as an upper and/or lower respiratory tract RSV infection) or a symptom or respiratory condition relating thereto (including, but not limited to asthma, wheezing, RAD, or a combination thereof), said methods comprising administering to a subject in need thereof an effective amount of one or more antibodies in combination with an effective amount of siplizumab (MedImmune, Inc., International Pub. No. WO 02/069904, which is incorporated herein by reference in its entirety). In another embodiment, the invention provides methods of preventing, treating, managing and/or ameliorating a RSV infection (i.e., an upper and/or lower respiratory tract RSV infection), otitis media (preferably, stemming from, caused by or associated with a RSV infection, such as an upper and/or lower respiratory tract RSV infection) or a symptom or respiratory condition relating thereto (including, but not limited to asthma, wheezing, RAD, or a combination thereof), said methods comprising administering to a subject in need thereof an effective amount of one or more antibodies in combination with an effective amount of one or more anti-IL-9 antibodies, such as those disclosed in U.S. Publication No. 2005/0002934, which is incorporated herein by reference in its entirety. In yet another embodiment, the invention provides methods for preventing, treating, managing, and/or ameliorating a RSV infection (i.e., an upper and/or lower respiratory tract RSV infection), otitis media (preferably, stemming from, caused by or associated with a RSV infection, such as an upper and/or lower respiratory tract RSV infection) or a symptom or respiratory condition relating thereto (including, but not limited to asthma, wheezing, RAD, or a combination thereof), said methods comprising administering to a subject in need thereof an effective amount of one or more antibodies of the invention in combination with an effective amount of two or more of the following: EphA2/EphrinA1 Modulators, an anti-IL-9 antibody and/or siplizumab.

The formulations of the invention, comprising antibodies, compositions, or [00351] combination therapies of the invention may be used as any line of therapy, including but not limited to, the first, second, third, fourth, or fifth line of therapy, to prevent, treat, and/or ameliorate an upper and/or lower respiratory tract RSV infection, otitis media (preferably, stemming from, caused by or associated with a RSV infection, such as an upper and/or lower respiratory tract RSV infection) or a symptom or respiratory condition relating thereto (including, but not limited to asthma, wheezing, RAD, or a combination thereof). The invention also includes methods of preventing, treating, managing, and/or ameliorating a RSV infection (i.e., an upper and/or lower respiratory tract RSV infection), otitis media (preferably, stemming from, caused by or associated with a RSV infection, such as an upper and/or lower respiratory tract RSV infection), or a symptom or respiratory condition relating thereto (including, but not limited to, asthma, wheezing, RAD, or a combination thereof) in a patient undergoing therapies for other diseases or disorders (e.g., non-RSV infections). The invention encompasses methods of preventing, managing, treating, and/or ameliorating a RSV infection (i.e., an upper and/or lower respiratory tract RSV infection), otitis media (preferably, stemming from, caused by or associated with a RSV infection, such as an upper and/or lower respiratory tract RSV infection) or a symptom or respiratory condition relating thereto (including, but not limited to asthma, wheezing, RAD, or a combination thereof) in a patient before any adverse effects or intolerance to therapies other than antibodies of the invention develops.

[00352] The invention also encompasses methods of preventing, treating, managing, and/or ameliorating a RSV infection (i.e., an upper and/or lower respiratory tract RSV infection), otitis media (preferably, stemming from, caused by or associated with a RSV infection, such as an upper and/or lower respiratory tract RSV infection), or a symptom or respiratory condition relating thereto (including, but not limited to, asthma, wheezing, RAD, or a combination thereof) in refractory patients. In certain embodiments, a patient with an upper and/or lower respiratory tract RSV infection, is refractory to a therapy when the infection has not significantly been eradicated and/or the symptoms have not been significantly alleviated. The determination of whether a patient is refractory can be made either in vivo or in vitro by any method known in the art for assaying the effectiveness of a therapy for infections, using artaccepted meanings of "refractory" in such a context. In various embodiments, a patient with an upper and/or lower respiratory tract RSV infection is refractory when viral replication has not decreased or has increased. The invention also encompasses methods of preventing the onset or reoccurrence of an upper and/or lower respiratory tract RSV infection or otitis media (preferably, stemming from, caused by or associated with a RSV infection, such as an upper

and/or lower respiratory tract RSV infection) in patients at risk of developing such infections or otitis media.

[00353] The invention also encompasses methods of preventing, managing, treating, and/or ameliorating a RSV infection (*i.e.*, an upper and/or lower respiratory tract RSV infection), otitis media (preferably, stemming from, caused by or associated with a RSV infection, such as an upper and/or lower respiratory tract RSV infection), or a symptom or respiratory condition relating thereto (including, but not limited to, asthma, wheezing, RAD, or a combination thereof) in patients who are susceptible to adverse reactions to conventional therapies. The invention further encompasses methods for preventing, treating, managing, and/or ameliorating a RSV infection (*i.e.*, an upper and/or lower respiratory tract RSV infection) or otitis media (preferably, stemming from, caused by or associated with a RSV infection, such as an upper and/or lower respiratory tract RSV infection) or a symptom or respiratory condition relating thereto (including, but not limited to, asthma, wheezing, RAD, or a combination thereof) for which no anti-viral therapy is available.

The invention encompasses methods for preventing, treating, managing, and/or [00354] ameliorating a RSV infection (i.e., an upper and/or lower respiratory tract RSV infection), otitis media (preferably, stemming from, caused by or associated with a RSV infection, such as an upper and/or lower respiratory tract RSV infection), or a symptom or respiratory condition relating thereto (including, but not limited to, asthma, wheezing, RAD, or a combination thereof) in a patient who has proven refractory to therapies other than antibodies of the invention but are no longer on these therapies. In certain embodiments, the patients being treated in accordance with the methods of this invention are patients already being treated with antibiotics, anti-virals, anti-fungals, or other biological therapy/immunotherapy. Among these patients are refractory patients, patients who are too young for conventional therapies, and patients with reoccurring upper and/or lower respiratory tract RSV infections or otitis media (preferably, stemming from, caused by or associated with a RSV infection, such as an upper and/or lower respiratory tract RSV infection) or a symptom or respiratory condition relating thereto (including, but not limited to, asthma, wheezing, RAD, or a combination thereof) despite treatment with existing therapies.

[00355] The present invention encompasses methods for preventing, treating and/or ameliorating a RSV infection (*i.e.*, an upper and/or lower respiratory tract RSV infection), otitis media (preferably, stemming from, caused by or associated with a RSV infection, such as an upper and/or lower respiratory tract RSV infection), or a symptom or respiratory condition

relating thereto (including, but not limited to, asthma, wheezing, RAD, or a combination thereof) as an alternative to other conventional therapies. In specific embodiments, the patient being or treated in accordance with the methods of the invention is refractory to other therapies or is susceptible to adverse reactions from such therapies. The patient may be a person with a suppressed immune system (e.g., post-operative patients, chemotherapy patients, and patients with immunodeficiency disease), a person with impaired renal or liver function, the elderly, children, infants, infants born prematurely, persons with neuropsychiatric disorders or those who take psychotropic drugs, persons with histories of seizures, or persons on medication that would negatively interact with conventional agents used to prevent, treat, and/or ameliorate an upper and/or lower respiratory tract RSV infection, otitis media (preferably, stemming from, caused by or associated with a RSV infection, such as an upper and/or lower respiratory tract RSV infection) or a symptom or respiratory condition relating thereto (including, but not limited to, asthma, wheezing, RAD, or a combination thereof).

[00356] The dosage amounts and frequencies of administration provided herein are encompassed by the terms "effective amount", "therapeutically effective" and "prophylactically" effective. The dosage and frequency further will typically vary according to factors specific for each patient depending on the specific therapeutic or prophylactic agents administered, the severity and type of infection, the route of administration, as well as age, body weight, response, and the past medical history of the patient. Suitable regimens can be selected by one skilled in the art by considering such factors and by following, for example, dosages reported in the literature and recommended in the Physician's Desk Reference (58th ed., 2004). See Section 5.3 for specific dosage amounts and frequencies of administration of the prophylactic and therapeutic agents provided by the invention.

5.6 <u>Methods of Administration of Antibodies</u>

[00357] The a specific embodiment, the invention provides methods of treatment, prophylaxis, and amelioration of an upper and/or lower respiratory tract RSV infection, otitis media (preferably, stemming from, caused by or associated with a RSV infection, such as an upper and/or lower respiratory tract RSV infection) or a symptom or respiratory condition relating thereto (including, but not limited to, asthma, wheezing, RAD, or a combination thereof) by administrating to a subject of an effective amount of antibody, or pharmaceutical composition comprising the formulation comprising an antibody of the invention. In a preferred aspect, an antibody is substantially purified (*i.e.*, substantially free from substances that limit its effect or produce undesired side-effects). The subject administered a therapy is preferably a mammal such as non-primate (*e.g.*, cows, pigs, horses, cats, dogs, rats etc.) and a primate (*e.g.*,

monkey such as a cynomolgous monkey and a human). In a preferred embodiment, the subject is a human. In another preferred embodiment, the subject is a human infant or a human infant born prematurely. In another embodiment, the subject is a human with an upper and/or lower respiratory tract RSV infection, otitis media stemming from, caused by or associated with a RSV infection, cystic fibrosis, bronchopulmonary dysplasia, congenital heart disease, congenital immunodeficiency or acquired immunodeficiency, a human who has had a bone marrow transplant, or an elderly human.

[00358] Various delivery systems are known and can be used to administer a prophylactic or therapeutic agent (e.g., an antibody of the invention), including, but not limited to. encapsulation in liposomes, microparticles, microcapsules, recombinant cells capable of expressing the antibody, receptor-mediated endocytosis (see, e.g., Wu and Wu, J. Biol. Chem. 262:4429-4432 (1987)), construction of a nucleic acid as part of a retroviral or other vector, etc. Methods of administering a prophylactic or therapeutic agent (e.g., an antibody of the invention), or pharmaceutical composition include, but are not limited to, parenteral administration (e.g., intradermal, intramuscular, intraperitoneal, intravenous and subcutaneous), epidural, and mucosal (e.g., intranasal and oral routes). In a specific embodiment, a prophylactic or therapeutic agent (e.g., an antibody of the present invention), or a pharmaceutical composition is administered intramuscularly, intravenously, or subcutaneously. The prophylactic or therapeutic agents, or compositions may be administered by any convenient route, for example by infusion or bolus injection, by absorption through epithelial or mucocutaneous linings (e.g., oral mucosa, rectal and intestinal mucosa, etc.) and may be administered together with other biologically active agents. Administration can be systemic or local. In addition, pulmonary administration can also be employed, e.g., by use of an inhaler or nebulizer, and formulation with an aerosolizing agent. See, e.g., U.S. Patent Nos. 6,019,968, 5,985,320, 5,985,309, 5,934,272, 5,874,064, 5,855,913, 5,290,540, and 4,880,078; and PCT Publication Nos. WO 92/19244, WO 97/32572, WO 97/44013, WO 98/31346, and WO 99/66903, each of which is incorporated herein by reference their entirety. In a specific embodiment, an antibody, or formulation of the invention is administered using Alkermes AIRTM pulmonary drug delivery technology (Alkermes, Inc., Cambridge, MA).

[00359] In a specific embodiment, it may be desirable to administer a prophylactic or therapeutic agent, or a pharmaceutical formulation of the invention locally to the area in need of treatment; this may be achieved by, for example, and not by way of limitation, local infusion, by injection, or by means of an implant, said implant being of a porous, non-porous, or gelatinous

material, including membranes, such as sialastic membranes, or fibers. Preferably, when administering an antibody of the invention, care must be taken to use materials to which the antibody does not absorb.

[00360] In another embodiment, a prophylactic or therapeutic agent, or a formulation of the invention can be delivered in a vesicle, in particular a liposome (see Langer, 1990, Science 249:1527-1533; Treat et al., in Liposomes in the Therapy of Infectious Disease and Cancer, Lopez-Berestein and Fidler (eds.), Liss, New York, pp. 353-365 (1989); Lopez-Berestein, ibid., pp. 317-327; see generally ibid.).

[00361] In another embodiment, a prophylactic or therapeutic agent, or a formulation of the invention can be delivered in a controlled release or sustained release system. In one embodiment, a pump may be used to achieve controlled or sustained release (see Langer, supra; Sefton, 1987, CRC Crit. Ref. Biomed. Eng. 14:20; Buchwald et al., 1980, Surgery 88:507; Saudek et al., 1989, N. Engl. J. Med. 321:574). In another embodiment, polymeric materials can be used to achieve controlled or sustained release of a prophylactic or therapeutic agent (e.g., an antibodies of the invention) or a formulation of the invention (see e.g., Medical Applications of Controlled Release, Langer and Wise (eds.), CRC Pres., Boca Raton, Florida (1974); Controlled Drug Bioavailability, Drug Product Design and Performance, Smolen and Ball (eds.), Wiley, New York (1984); Ranger and Peppas, 1983, J., Macromol, Sci. Rev. Macromol, Chem. 23:61: see also Levy et al., 1985, Science 228:190; During et al., 1989, Ann. Neurol. 25:351; Howard et al., 1989, J. Neurosurg. 7 1:105); U.S. Patent No. 5,679,377; U.S. Patent No. 5,916,597; U.S. Patent No. 5,912,015; U.S. Patent No. 5,989,463; U.S. Patent No. 5,128,326; PCT Publication No. WO 99/15154; and PCT Publication No. WO 99/20253. Examples of polymers used in sustained release formulations include, but are not limited to, poly(2-hydroxy ethyl methacrylate), poly(methyl methacrylate), poly(acrylic acid), poly(ethylene-co-vinyl acetate), poly(methacrylic acid), polyglycolides (PLG), polyanhydrides, poly(N-vinyl pyrrolidone), poly(vinyl alcohol), polyacrylamide, poly(ethylene glycol), polylactides (PLA), poly(lactide-coglycolides) (PLGA), and polyorthoesters. In a preferred embodiment, the polymer used in a sustained release formulation is inert, free of leachable impurities, stable on storage, sterile, and biodegradable. In yet another embodiment, a controlled or sustained release system can be placed in proximity of the therapeutic target, i.e., the nasal passages or lungs, thus requiring only a fraction of the systemic dose (see, e.g., Goodson, in Medical Applications of Controlled Release, supra, vol. 2, pp. 115-138 (1984)).

[00362] Controlled release systems are discussed in the review by Langer (1990, Science 249:1527-1533). Any technique known to one of skill in the art can be used to produce sustained release formulations comprising one or more antibodies of the invention. See, e.g., U.S. Patent No. 4,526,938, PCT publication WO 91/05548, PCT publication WO 96/20698, Ning et al., 1996, "Intratumoral Radioimmunotheraphy of a Human Colon Cancer Xenograft Using a Sustained-Release Gel," Radiotherapy & Oncology 39:179-189, Song et al., 1995, "Antibody Mediated Lung Targeting of Long-Circulating Emulsions," PDA Journal of Pharmaceutical Science & Technology 50:372-397, Cleek et al., 1997, "Biodegradable Polymeric Carriers for a bFGF Antibody for Cardiovascular Application," Pro. Int'l. Symp. Control. Rel. Bioact. Mater. 24:853-854, and Lam et al., 1997, "Microencapsulation of Recombinant Humanized Monoclonal Antibody for Local Delivery," Proc. Int'l. Symp. Control Rel. Bioact. Mater. 24:759-760, each of which is incorporated herein by reference in their entirety.

[00363] In a specific embodiment, a formulation of the invention comprises one, two or more antibodies described, infra. In another embodiment, a formulation of the invention comprises one, two or more antibodies described, infra, and a prophylactic or therapeutic agent other than an said antibody. In a specific embodiment, the agents are known to be useful for or have been or are currently used for the prevention, treatment or ameliorating of a RSV infection (preferably, an upper and/or lower respiratory tract RSV infection), otitis media (preferably stemming from, caused by or associated with a RSV infection, such as an upper and/or lower respiratory tract RSV infection) or a symptom or respiratory condition relating thereto (including, but not limited to, asthma, wheezing, RAD, or a combination thereof). In addition to prophylactic or therapeutic agents, the compositions of the invention may also comprise a carrier.

[00364] The formulations of the invention include bulk drug compositions useful in the manufacture of pharmaceutical compositions (e.g., compositions that are suitable for administration to a subject or patient) which can be used in the preparation of unit dosage forms. In a preferred embodiment, a composition of the invention is a pharmaceutical composition. Such compositions comprise a prophylactically or therapeutically effective amount of one or more prophylactic or therapeutic agents (e.g., an antibody of the invention or other prophylactic or therapeutic agent), and a pharmaceutically acceptable carrier. Preferably, the pharmaceutical compositions are formulated to be suitable for the route of administration to a subject.

[00365] In a specific embodiment, the term "pharmaceutically acceptable" means approved by a regulatory agency of the Federal or a state government or listed in the U.S. Pharmacopeia or other generally recognized pharmacopeia for use in animals, and more particularly in humans. The term "carrier" refers to a diluent, adjuvant (e.g., Freund's adjuvant (complete and incomplete)), excipient, or vehicle with which the therapeutic is administered. Such pharmaceutical carriers can be sterile liquids, such as water and oils, including those of petroleum, animal, vegetable or synthetic origin, such as peanut oil, soybean oil, mineral oil, sesame oil and the like. Water is a preferred carrier when the pharmaceutical composition is administered intravenously. Saline solutions and aqueous dextrose and glycerol solutions can also be employed as liquid carriers, particularly for injectable solutions. Suitable pharmaceutical excipients include starch, glucose, lactose, sucrose, gelatin, malt, rice, flour, chalk, silica gel. sodium stearate, glycerol monostearate, talc, sodium chloride, dried skim milk, glycerol, propylene, glycol, water, ethanol and the like. The composition, if desired, can also contain minor amounts of wetting or emulsifying agents, or pH buffering agents. These compositions can take the form of solutions, suspensions, emulsion, tablets, pills, capsules, powders. sustained-release formulations and the like. Oral formulation can include standard carriers such as pharmaceutical grades of mannitol, lactose, starch, magnesium stearate, sodium saccharine, cellulose, magnesium carbonate, etc. Examples of suitable pharmaceutical carriers are described in "Remington's Pharmaceutical Sciences" by E.W. Martin. Such compositions will contain a prophylactically or therapeutically effective amount of the antibody, preferably in purified form, together with a suitable amount of carrier so as to provide the form for proper administration to the patient. The formulation should suit the mode of administration.

[00366] In a preferred embodiment, the formulations are manufactured in accordance with routine procedures as a pharmaceutical composition adapted for intravenous administration to human beings. Typically, compositions for intravenous administration are solutions in sterile isotonic aqueous buffer. Where necessary, the composition may also include a solubilizing agent and a local anesthetic such as lignocamne to ease pain at the site of the injection.

[00367] Generally, the ingredients of compositions of the invention are supplied either separately or mixed together in unit dosage form, for example, as a dry lyophilized powder or water free concentrate in a hermetically sealed container such as an ampoule or sachette indicating the quantity of active agent. Where the composition is to be administered by infusion, it can be dispensed with an infusion bottle containing sterile pharmaceutical grade water or

saline. Where the composition is administered by injection, an ampoule of sterile water for injection or saline can be provided so that the ingredients may be mixed prior to administration.

[00368] The invention also provides that the formulaiton is packaged in a hermetically sealed container such as an ampoule or sachette indicating the quantity of antibody. In one embodiment, the formulation of the invention comprising an antibody is supplied as a dry sterilized lyophilized powder or water free concentrate in a hermetically sealed container and can be reconstituted, e.g., with water or saline to the appropriate concentration for administration to a subject. In one embodiment, the formulation of the invention comprising an antibody is supplied as a dry sterile lyophilized powder in a hermetically sealed container at a unit dosage of at least 3 mg, more preferably at least 5 mg, at least 10 mg, at least 15 mg, at least 25 mg, at least 30 mg, at least 35 mg, at least 45 mg, at least 50 mg, at least 60 mg, or at least 75 mg. The lyophilized formulation of the invention comprising an antibody should be stored at between 2 and 8° C in its original container and the antibody should be administered within 12 hours, preferably within 6 hours, within 5 hours, within 3 hours, or within 1 hour after being reconstituted. In an alternative embodiment, a formulation of the invention comprising an antibody is supplied in liquid form in a hermetically sealed container indicating the quantity and concentration of the antibody. Preferably, the liquid form of the formulation of the invention comprising an antibody is supplied in a hermetically sealed container at least 1 mg/ml, more preferably at least 2.5 mg/ml, at least 3 mg/ml, at least 5 mg/ml, at least 8 mg/ml, at least 10 mg/ml, at least 15 mg/ml, at least 25 mg/ml, at least 30 mg/ml, or at least 60 mg/ml.

[00369] The formulation of the invention comprising antibodies can be formulated as neutral or salt forms. Pharmaceutically acceptable salts include those formed with anions such as those derived from hydrochloric, phosphoric, acetic, oxalic, tartaric acids, etc., and those formed with cations such as those derived from sodium, potassium, ammonium, calcium, ferric hydroxides, isopropylamine, triethylamine, 2-ethylamino ethanol, histidine, procaine, etc.

[00370] The amount of a prophylactic or therapeutic agent (e.g., an antibody of the invention), or a composition of the invention which will be effective in the treatment, prevention or amelioration of an upper and/or lower respiratory tract RSV infection, otitis media (preferably, stemming from, caused by or associated with a RSV infection, such as an upper and/or lower respiratory tract RSV infection) or a symptom or respiratory condition relating thereto (including, but not limited to, asthma, wheezing, RAD, or a combination thereof) can be determined by standard clinical techniques. For example, the dosage of a prophylactic or therapeutic agent, or a composition which will be effective in the treatment, prevention or

amenoration of an upper and/or lower respiratory tract RSV infection or one or more symptoms thereof can be determined by administering the composition to a cotton rat, measuring the RSV titer after challenging the cotton rat with 10⁵ pfu of RSV and comparing the RSV titer to that obtain for a cotton rat not administered the prophylactic or therapeutic agent, or the composition. Accordingly, a dosage that results in a 2 log decrease or a 99% reduction in RSV titer in the cotton rat challenged with 10⁵ pfu of RSV relative to the cotton rat challenged with 10⁵ pfu of RSV but not administered the prophylactic or therapeutic agent, or the composition is the dosage of the composition that can be administered to a human for the treatment, prevention or amelioration of an upper and/or lower respiratory tract RSV infection, otitis media (preferably, stemming from, caused by or associated with a RSV infection, such as an upper and/or lower respiratory tract RSV infection) or one or more symptoms thereof.

[00371] The dosage of a composition which will be effective in the treatment, prevention or amelioration of an upper and/or lower respiratory tract RSV infection, otitis media (preferably, stemming from, caused by or associated with a RSV infection, such as an upper and/or lower respiratory tract RSV infection) or one or more symptoms thereof can be determined by administering the composition to an animal model (e.g., a cotton rat or monkey) and measuring the serum titer, lung concentration or nasal turbinate and/or nasal secretion concentration of an antibody that immunospecifically bind to a RSV antigen. Accordingly, a dosage of an antibody or a composition that results in a serum titer of at least 1 µg/ml, preferably $2~\mu g/ml,\,5~\mu g/ml,\,10~\mu g/ml,\,15~\mu g/ml,\,20~\mu g/ml,\,25~\mu g/ml,$ at least 30 $\mu g/ml,$ at least 35 $\mu g/ml,$ at least 40 $\mu g/ml$, at least 50 $\mu g/ml$, at least 75 $\mu g/ml$, at least 100 $\mu g/ml$, at least 125 $\mu g/ml$, at least 150 μg/ml, at least 200 μg/ml, at least 250 μg/ml, at least 300 μg/ml, at least 350 μg/ml, at least 400 μ g/ml, or at least 450 μ g/ml can be administered to a human for the treatment, prevention or amelioration of an upper and/or lower respiratory tract RSV infection, otitis media (preferably, stemming from, caused by or associated with a RSV infection, such as an upper and/or lower respiratory tract RSV infection) or one or more symptoms thereof. In addition, in vitro assays may optionally be employed to help identify optimal dosage ranges.

[00372] The precise dose to be employed in the formulation will also depend on the route of administration, and the seriousness of the upper and/or lower respiratory tract RSV infection or otitis media, and should be decided according to the judgment of the practitioner and each patient's circumstances. Effective doses may be extrapolated from dose-response curves derived from *in vitro* or animal model (e.g., the cotton rat or Cynomolgous monkey) test systems.

[00373] For antibodies, the dosage administered to a patient is typically 0.1 mg/kg to 100 mg/kg of the patient's body weight. In some embodiments, the dosage administered to the patient is about 3 mg/kg to about 60 mg/kg of the patient's body weight. Preferably, the dosage administered to a patient is between 0.1 mg/kg and 20 mg/kg of the patient's body weight, more preferably 1 mg/kg to 15 mg/kg of the patient's body weight. Generally, human antibodies have a longer half-life within the human body than antibodies from other species due to the immune response to the foreign polypeptides. Thus, lower dosages of human antibodies and less frequent administration is often possible. Further, the dosage and frequency of administration of antibodies of the invention may be reduced by enhancing uptake and tissue penetration (e.g., into the nasal passages and/or lung) of the antibodies by modifications such as, for example, lipidation. In a preferred embodiment, the dosage of A4B4L1FR-S28R (motavizumab) or antigen-binding fragment thereof to be administered to is 60 mg/kg, 50 mg/kg, 40 mg/kg, 30 mg/kg, 15 mg/kg, 10 mg/kg, 5 mg/kg, 3 mg/kg, or 2 mg/kg of the patient's body weight.

In a specific embodiment, formulations of the invention comprising antibodies or compositions comprising antibodies are administered once a month just prior to or during the RSV season. In another embodiment, formulation of the invention comprising an antibody, or compositions comprising antibodies produced in accordance with the methods of the invention are administered every two months just prior to or during the RSV season. In yet another embodiment, antibodies, or compositions comprising antibodies are administered once just prior to or during the RSV season. The term "RSV season" refers to the season when RSV infection is most likely to occur. Typically, the RSV season in the northern hemisphere commences in November and lasts through April. Preferably, the antibody comprises the VH and VL domain of A4B4L1FR-S28R (motavizumab) (Figure 13) or an antigen-binding fragment thereof.

In one embodiment, approximately 60 mg/kg or less, approximately 45 mg/kg or less, approximately 30 mg/kg or less, approximately 15 mg/kg or less, approximately 10 mg/kg or less, approximately 5 mg/kg or less, approximately 3 mg/kg or less, approximately 2 mg/kg or less, or approximately 1.5 mg/kg or less of an antibody the invention is administered 5 times, 4 times, 3 times, 2 times or 1 time during a RSV season to a subject, preferably a human. In some embodiments, the antibody is administered about 1-12 times during the RSV season to a subject, wherein the doses may be administered as necessary, *e.g.*, weekly, biweekly, monthly, bimonthly, trimonthly, *etc.*, as determined by a physician. In some embodiments, a lower dose (*e.g.*, 5-15 mg/kg) can be administered more frequently (*e.g.*, 3-6 times) during a RSV season. In other embodiments, a higher dose (*e.g.*, 30-60 mg/kg) can be administered less frequently

(e.g., 1-3 times) during a RSV season. However, as will be apparent to those in the art, other dosing amounts and schedules are easily determinable and within the scope of the invention.

[00376] In one embodiment, approximately 60 mg/kg or less, approximately 45 mg/kg or less, approximately 30 mg/kg or less, approximately 15 mg/kg or less, approximately 10 mg/kg or less, approximately 5 mg/kg or less, approximately 3 mg/kg or less, approximately 2 mg/kg or less, or approximately 1.5 mg/kg or less of an antibody is administered to monthly five times during a RSV season to a subject, preferably a human, intramuscularly. In another embodiment, approximately 60 mg/kg, approximately 45 mg/kg or less, approximately 30 mg/kg or less, approximately 15 mg/kg or less, approximately 10 mg/kg or less, approximately 5 mg/kg or less. approximately 3 mg/kg or less, approximately 2 mg/kg or less, or approximately 1.5 mg/kg or less of an antibody the invention is administered monthly three times during a RSV season to a subject, preferably a human, intramuscularly. In yet another embodiment, approximately 60 mg/kg, approximately 45 mg/kg or less, approximately 30 mg/kg or less, approximately 15 mg/kg or less, approximately 10 mg/kg or less, approximately 5 mg/kg or less, approximately 3 mg/kg or less, approximately 2 mg/kg or less, or approximately 1.5 mg/kg or less of an antibody is administered monthly one to two times during a RSV season to a subject, preferably a human, intramuscularly. Preferably, the antibody comprises the VH and VL domain of A4B4L1FR-S28R (motavizumab) (Figure 13) or an antigen-binding fragment thereof.

In a specific embodiment, approximately 60 mg/kg, approximately 45 mg/kg or [00377] less, approximately 30 mg/kg or less, approximately 15 mg/kg or less, approximately 10 mg/kg or less, approximately 5 mg/kg or less, approximately 3 mg/kg or less, approximately 2 mg/kg or less, or approximately 1.5 mg/kg or less of an antibody in a sustained release formulation is administered to a subject, preferably a human, to prevent, treat or ameliorate an upper and/or lower respiratory tract RSV infection, otitis media (preferably, stemming from, caused by or associated with a RSV infection, such as an upper and/or lower respiratory tract RSV infection) or one or more symptoms thereof. In another specific embodiment, an approximately 60 mg/kg, approximately 45 mg/kg or less, approximately 30 mg/kg or less, approximately 15 mg/kg or less, approximately 10 mg/kg or less, approximately 5 mg/kg or less, approximately 3 mg/kg or less, approximately 2 mg/kg or less, or approximately 1.5 mg/kg or less bolus of an antibody the invention not in a sustained release formulation is administered to a subject, preferably a human, to prevent, treat or ameliorate an upper and/or lower respiratory tract RSV infection, otitis media (preferably, stemming from, caused by or associated with a RSV infection, such as an upper and/or lower respiratory tract RSV infection) or one or more symptoms thereof and after a

certain period of time approximately 60 mg/kg, approximately 45 mg/kg or less, approximately 30 mg/kg or less, approximately 15 mg/kg or less, approximately 10 mg/kg or less, approximately 5 mg/kg or less, approximately 3 mg/kg or less, approximately 2 mg/kg or less, or approximately 1.5 mg/kg or less of the invention in a sustained release is administered to said subject intramuscularly two, three or four times during a RSV season. In accordance with this embodiment, a certain period of time can be 1 to 5 days, a week, two weeks, or a month. In another embodiment, approximately 60 mg/kg, approximately 45 mg/kg or less, approximately 30 mg/kg or less, approximately 15 mg/kg or less, approximately 10 mg/kg or less, approximately 5 mg/kg or less, approximately 3 mg/kg or less, approximately 2 mg/kg or less, or approximately 1.5 mg/kg or less of an antibody in a sustained release formulation is administered to a subject, preferably a human, intramuscularly two, three or four times during a RSV season to prevent, treat or ameliorate an upper and/or lower respiratory tract RSV infection, otitis media (preferably, stemming from, caused by or associated with a RSV infection, such as an upper and/or lower respiratory tract RSV infection) or one or more symptoms thereof. Preferably, the antibody is A4B4L1FR-S28 or an antigen-binding fragment thereof.

[00378] In another embodiment, approximately 60 mg/kg, approximately 45 mg/kg or less, approximately 30 mg/kg or less, approximately 15 mg/kg or less, approximately 10 mg/kg or less, approximately 5 mg/kg or less, approximately 3 mg/kg or less, approximately 2 mg/kg or less, or approximately 1.5 mg/kg or less of one or more antibodies of the invention is administered intranasally to a subject to prevent, treat or ameliorate an upper and/or lower respiratory tract RSV infection, otitis media (preferably, stemming from, caused by or associated with a RSV infection, such as an upper and/or lower respiratory tract RSV infection) or one or more symptoms thereof. Preferably, the antibody is A4B4L1FR-S28 or an antigen-binding fragment thereof.

[00379] In one embodiment, a single dose of the formulation of the invention comprising an antibody (preferably motavizumab) is administered to a patient (preferably a human), wherein the dose is selected from the group consisting of about 1 mg/kg, about 3 mg/kg, about 5 mg/kg, about 10 mg/kg, about 15 mg/kg, about 20 mg/kg, about 25 mg/kg, about 30 mg/kg, about 35 mg/kg, about 40 mg/kg, about 45 mg/kg, about 50 mg/kg, about 55 mg/kg, about 60 mg/kg, about 65 mg/kg, about 70 mg/kg, or about 75 mg/kg.

In some embodiments, a single dose of a formulation of the invention comprising an antibody (preferably motavizumab) is administered to a patient (preferably a human) two, three, four, five, six, seven, eight, nine, ten, eleven, twelve times, thirteen, fourteen, fifteen, sixteen, seventeen, eighteen, nineteen, twenty, twenty-one, twenty-two, twenty-three, twenty-four, twenty five, or twenty six at bi-weekly (e.g., about 14 day) intervals over the course of a year (or alternatively over the course of a RSV season), wherein the dose is selected from the group consisting of about 1 mg/kg, about 3 mg/kg, about 5 mg/kg, about 10 mg/kg, about 15 mg/kg, about 20 mg/kg, about 25 mg/kg, about 30 mg/kg, about 35 mg/kg, about 40 mg/kg, about 45 mg/kg, about 50 mg/kg, about 55 mg/kg, about 60 mg/kg, about 65 mg/kg, about 70 mg/kg, about 75 mg/kg, or a combination thereof (i.e., each dose monthly dose may or may not be identical).

[00381] In another embodiment, a single dose of a formulation of the invention comprising an antibody (preferably motavizumab) is administered to patient (preferably a human) two, three, four, five, six, seven. eight, nine, ten, eleven, or twelve times at about monthly (e.g., about 30 day) intervals over the course of a year (or alternatively over the course of a RSV season), wherein the dose is selected from the group consisting of about 1 mg/kg, about 3 mg/kg, about 5 mg/kg, about 10 mg/kg, about 15 mg/kg, about 20 mg/kg, about 25 mg/kg, about 30 mg/kg, about 35 mg/kg, about 40 mg/kg, about 45 mg/kg, about 50 mg/kg, about 55 mg/kg, about 60 mg/kg, about 65 mg/kg, about 70 mg/kg, about 75 mg/kg, or a combination thereof (i.e., each dose monthly dose may or may not be identical).

In one embodiment, a single dose of a formulation of the invention comprising an antibody (preferably motavizumab) is administered to a patient (preferably a human) two, three, four, five, or six times at about bi-monthly (e.g., about 60 day) intervals over the course of a year (or alternatively over the course of a RSV season), wherein the dose is selected from the group consisting of about 1 mg/kg, about 3 mg/kg, about 5 mg/kg, about 10 mg/kg, about 15 mg/kg, about 20 mg/kg, about 25 mg/kg, about 30 mg/kg, about 35 mg/kg, about 40 mg/kg, about 45 mg/kg, about 50 mg/kg, about 55 mg/kg, about 60 mg/kg, about 65 mg/kg, about 70 mg/kg, about 75 mg/kg, or a combination thereof (i.e., each bi-monthly dose may or may not be identical).

[00383] In some embodiments, a single dose of a formulation of the invention comprising an antibody (preferably motavizumab) is administered to a patient (preferably a human) two, three, or four times at about tri-monthly (e.g., about 120 day) intervals over the course of a year (or alternatively over the course of a RSV season), wherein the dose is selected from the group

consisting of about 1 mg/kg, about 3 mg/kg, about 5 mg/kg, about 10 mg/kg, about 15 mg/kg, about 20 mg/kg, about 25 mg/kg, about 30 mg/kg, about 35 mg/kg, about 40 mg/kg, about 45 mg/kg, about 50 mg/kg, about 55 mg/kg, about 60 mg/kg, about 65 mg/kg, about 70 mg/kg, about 75 mg/kg, or a combination thereof (*i.e.*, each tri-monthly dose may or may not be identical).

[00384] In certain embodiments, the route of administration for a dose of a formulation of the invention comprising an antibody to a patient is intramuscular, intravenous, or a combination thereof (i.e., each dose may or may not be administered by an identical route of administration). In some embodiments, an antibody of the invention may be administered via multiple routes of administration simultaneously or subsequently to other doses of the same or a different antibody of the invention.

5.7 Biological Activity

[00385] Formulations of the invention comprising antibodies may be characterized in a variety of ways. In particular, antibodies may be assayed for the ability to immunospecifically bind to a RSV antigen. Such an assay may be performed in solution (*e.g.*, Houghten, 1992, Bio/Techniques 13:412-421), on beads (Lam, 1991, Nature 354:82-84), on chips (Fodor, 1993, Nature 364:555-556), on bacteria (U.S. Patent No. 5,223,409), on spores (U.S. Patent Nos. 5,571,698; 5,403,484; and 5,223,409), on plasmids (Cull et al., 1992, Proc. Natl. Acad. Sci. USA 89:1865-1869) or on phage (Scott and Smith, 1990, Science 249:386-390; Devlin, 1990, Science 249:404-406; Cwirla et al., 1990, Proc. Natl. Acad. Sci. USA 87:6378-6382; and Felici, 1991, J. Mol. Biol. 222:301-310) (each of these references is incorporated herein in its entirety by reference). Antibodies that have been identified to immunospecifically bind to a RSV antigen (*e.g.*, a RSV F antigen) can then be assayed for their specificity and affinity for a RSV antigen.

[00386] Formulations of the invention comprising antibodies may be assayed for immunospecific binding to a RSV antigen and cross-reactivity with other antigens by any method known in the art. Immunoassays which can be used to analyze immunospecific binding and cross-reactivity include, but are not limited to, competitive and non-competitive assay systems using techniques such as western blots, radioimmunoassays, ELISA (enzyme linked immunosorbent assay), "sandwich" immunoassays, immunoprecipitation assays, precipitin reactions, gel diffusion precipitin reactions, immunodiffusion assays, agglutination assays, complement-fixation assays, immunoradiometric assays, fluorescent immunoassays, protein A immunoassays, to name but a few. Such assays are routine and well known in the art (see, e.g.,

Ausubel et al, eds, 1994, Current Protocols in Molecular Biology, Vol. 1, John Wiley & Sons, Inc., New York, which is incorporated by reference herein in its entirety). Exemplary immunoassays are described briefly below (but are not intended by way of limitation).

Immunoprecipitation protocols generally comprise lysing a population of cells in a lysis buffer such as RIPA buffer (1% NP-40 or Triton X-100, 1% sodium deoxycholate, 0.1% SDS, 0.15 M NaCl, 0.01 M sodium phosphate at pH 7.2, 1% Trasylol) supplemented with protein phosphatase and/or protease inhibitors (e.g., EDTA, PMSF, aprotinin, sodium vanadate), adding the antibody of interest to the cell lysate, incubating for a period of time (e.g., 1 to 4 hours) at 40° C, adding protein A and/or protein G sepharose beads to the cell lysate, incubating for about an hour or more at 40° C, washing the beads in lysis buffer and resuspending the beads in SDS/sample buffer. The ability of the antibody of interest to immunoprecipitate a particular antigen can be assessed by, e.g., western blot analysis. One of skill in the art would be knowledgeable as to the parameters that can be modified to increase the binding of the antibody to an antigen and decrease the background (e.g., pre-clearing the cell lysate with sepharose beads). For further discussion regarding immunoprecipitation protocols see, e.g., Ausubel et al, eds, 1994, Current Protocols in Molecular Biology, Vol. 1, John Wiley & Sons, Inc., New York at 10.16.1.

[00388] Western blot analysis generally comprises preparing protein samples, electrophoresis of the protein samples in a polyacrylamide gel (e.g., 8%-20% SDS-PAGE depending on the molecular weight of the antigen), transferring the protein sample from the polyacrylamide gel to a membrane such as nitrocellulose, PVDF or nylon, incubating the membrane in blocking solution (e.g., PBS with 3% BSA or non-fat milk), washing the membrane in washing buffer (e.g., PBS-Tween 20), incubating the membrane with primary antibody (the antibody of interest) diluted in blocking buffer, washing the membrane in washing buffer, incubating the membrane with a secondary antibody (which recognizes the primary antibody, e.g., an anti-human antibody) conjugated to an enzymatic substrate (e.g., horseradish peroxidase or alkaline phosphatase) or radioactive molecule (e.g., ³²P or ¹²⁵I) diluted in blocking buffer, washing the membrane in wash buffer, and detecting the presence of the antigen. One of skill in the art would be knowledgeable as to the parameters that can be modified to increase the signal detected and to reduce the background noise. For further discussion regarding western blot protocols see, e.g., Ausubel et al, eds, 1994, Current Protocols in Molecular Biology, Vol. 1, John Wiley & Sons, Inc., New York at 10.8.1.

ELISAs comprise preparing antigen, coating the well of a 96 well microtiter plate with the antigen, adding the antibody of interest conjugated to a detectable compound such as an enzymatic substrate (e.g., horseradish peroxidase or alkaline phosphatase) to the well and incubating for a period of time, and detecting the presence of the antigen. In ELISAs the antibody of interest does not have to be conjugated to a detectable compound; instead, a second antibody (which recognizes the antibody of interest) conjugated to a detectable compound may be added to the well. Further, instead of coating the well with the antigen, the antibody may be coated to the well. In this case, a second antibody conjugated to a detectable compound may be added following the addition of the antigen of interest to the coated well. One of skill in the art would be knowledgeable as to the parameters that can be modified to increase the signal detected as well as other variations of ELISAs known in the art. For further discussion regarding ELISAs see, e.g., Ausubel et al, eds, 1994, Current Protocols in Molecular Biology, Vol. 1, John Wiley & Sons, Inc., New York at 11.2.1.

[00390] The binding affinity of an antibody to an antigen and the off-rate of an antibody-antigen interaction can be determined by competitive binding assays. One example of a competitive binding assay is a radioimmunoassay comprising the incubation of labeled antigen (e.g., ³H or ¹²⁵I) with the antibody of interest in the presence of increasing amounts of unlabeled antigen, and the detection of the antibody bound to the labeled antigen. The affinity of the antibody of the present invention for a RSV antigen and the binding off-rates can be determined from the data by scatchard plot analysis. Competition with a second antibody can also be determined using radioimmunoassays. In this case, a RSV antigen is incubated with an antibody of the present invention conjugated to a labeled compound (e.g., ³H or ¹²⁵I) in the presence of increasing amounts of an unlabeled second antibody.

[00391] In a preferred embodiment, BIAcore kinetic analysis is used to determine the binding on and off rates of antibodies to a RSV antigen. BIAcore kinetic analysis comprises analyzing the binding and dissociation of a RSV antigen from chips with immobilized antibodies on their surface.

[00392] Formulations of the invention comprising antibodies can also be assayed for their ability to inhibit the binding of RSV to its host cell receptor using techniques known to those of skill in the art. For example, cells expressing the receptor for RSV can be contacted with RSV in the presence or absence of an antibody and the ability of the antibody to inhibit RSV's binding can measured by, for example, flow cytometry or a scintillation assay. RSV (e.g., a RSV antigen such as F glycoprotein or G glycoprotein) or the antibody can be labeled with a

detectable compound such as a radioactive label (e.g., 32P, 35S, and 125I) or a fluorescent label (e.g., fluorescein isothiocyanate, rhodamine, phycocrythrin, phycocyanin, allophycocyanin, ophthaldehyde and fluorescamine) to enable detection of an interaction between RSV and its host cell receptor. Alternatively, the ability of antibodies to inhibit RSV from binding to its receptor can be determined in cell-free assays. For example, RSV or a RSV antigen such as G glycoprotein can be contacted with an antibody and the ability of the antibody to inhibit RSV or the RSV antigen from binding to its host cell receptor can be determined. Preferably, the antibody is immobilized on a solid support and RSV or a RSV antigen is labeled with a detectable compound. Alternatively, RSV or a RSV antigen is immobilized on a solid support and the antibody is labeled with a detectable compound. RSV or a RSV antigen may be partially or completely purified (e.g., partially or completely free of other polypeptides) or part of a cell lysate. Further, a RSV antigen may be a fusion protein comprising the RSV antigen and a domain such as glutathionine S transferase. Alternatively, a RSV antigen can be biotinylated using techniques well known to those of skill in the art (e.g., biotinylation kit, Pierce Chemicals; Rockford, IL).

[00393] Formulations of the invention comprising antibodies can also be assayed for their ability to inhibit or downregulate RSV replication using techniques known to those of skill in the art. For example, RSV replication can be assayed by a plaque assay such as described, *e.g.*, by Johnson et al., 1997, Journal of Infectious Diseases 176:1215-1224. The antibodies of the invention can also be assayed for their ability to inhibit or downregulate the expression of RSV polypeptides. Techniques known to those of skill in the art, including, but not limited to, Western blot analysis, Northern blot analysis, and RT-PCR can be used to measure the expression of RSV polypeptides. Further, the antibodies of the invention can be assayed for their ability to prevent the formation of syncytia.

[00394] Formulations of the invention comprising antibodies are preferably tested *in vitro*, and then *in vivo* for the desired therapeutic or prophylactic activity, prior to use in humans. For example, *in vitro* assays which can be used to determine whether administration of a specific antibody or composition of the present invention is indicated, include *in vitro* cell culture assays in which a subject tissue sample is grown in culture, and exposed to or otherwise administered an antibody or composition of the present invention, and the effect of such an antibody or composition of the present invention upon the tissue sample is observed. In various specific embodiments, *in vitro* assays can be carried out with representative cells of cell types involved in a RSV infection (*e.g.*, respiratory epithelial cells), to determine if an antibody or composition

of the present invention has a desired effect upon such cell types. Preferably, the antibodies or compositions of the invention are also tested in *in vitro* assays and animal model systems prior to administration to humans. In a specific embodiment, cotton rats are administered an antibody the invention, or a composition of the invention, challenged with 10⁵ pfu of RSV, and four or more days later the rats are sacrificed and RSV titer and anti-RSV antibody serum titer is determined. Further, in accordance with this embodiment, the tissues (*e.g.*, the lung tissues) from the sacrificed rats can be examined for histological changes.

[00395] In accordance with the invention, clinical trials with human subjects need not be performed in order to demonstrate the prophylactic and/or therapeutic efficacy of antibodies of the invention. *In vitro* and animal model studies using the antibodies can be extrapolated to humans and are sufficient for demonstrating the prophylactic and/or therapeutic utility of said antibodies.

[00396] Formulations of the invention comprising antibodies or compositions of the present invention for use in therapy can be tested for their toxicity in suitable animal model systems, including but not limited to rats, mice, cows, monkeys, and rabbits. For *in vivo* testing of an antibody or composition's toxicity any animal model system known in the art may be used.

Efficacy in treating or preventing an upper and/or lower respiratory tract RSV [00397]infection may be demonstrated by determining the ability of an antibody or composition of the invention to inhibit the replication of the virus, to inhibit transmission or prevent the virus from establishing itself in its host, to reduce the incidence of an upper and/or lower respiratory tract RSV infection, to prevent or reduce the progression of an upper respiratory tract RSV infection to a lower respiratory tract RSV infection, or to prevent, ameliorate or alleviate one or more symptoms associated with an upper and/or lower respiratory tract RSV infection. Efficacy in treating or preventing otitis media may be demonstrated by determining the ability of an antibody or composition of the invention to reduce the incidence or otitis media, to reduce the duration of otitis media, to prevent or reduce the progression of an upper and/or lower respiratory tract RSV infection to otitis media, or to ameliorate one or more symptoms of otitis media. A therapy is considered therapeutic if there is, for example, a reduction is viral load, amelioration of one or more symptoms of an upper and/or lower respiratory tract RSV infection or otitis media, or a respiratory condition relating thereto (including, but not limited to asthma, wheezing, RAD or a combination thereof), a reduction in the duration of an upper and/or lower respiratory tract RSV infection or otitis media, a reduction in lower respiratory tract RSV infections, or a decrease in mortality and/or morbidity following administration of an antibody

or composition of the invention. Further, the treatment is considered therapeutic if there is an increase in the immune response following the administration of one or more antibodies which immunospecifically bind to one or more RSV antigens.

[00398] Formulations of the invention comprising antibodies or compositions of the invention can be tested *in vitro* and *in vivo* for the ability to induce the expression of cytokines such as IFN-α, IFN-γ, IL-², IL-3, IL-4, IL-5, IL-6, IL-7, IL-8, IL-9, IL-10, IL-12 and IL-15. Techniques known to those of skill in the art can be used to measure the level of expression of cytokines. For example, the level of expression of cytokines can be measured by analyzing the level of RNA of cytokines by, for example, RT-PCR and Northern blot analysis, and by analyzing the level of cytokines by, for example, immunoprecipitation followed by western blot analysis and ELISA. In a preferred embodiment, an antibody or composition of the invention is tested for its ability to induce the expression of IFN-γ.

Formulations of the invention comprising antibodies or compositions of the [00399] invention can be tested in vitro and in vivo for their ability to modulate the biological activity of immune cells, preferably human immune cells (e.g., T-cells, B-cells, and Natural Killer cells). The ability of an antibody or composition of the invention to modulate the biological activity of immune cells can be assessed by detecting the expression of antigens, detecting the proliferation of immune cells, detecting the activation of signaling molecules, detecting the effector function of immune cells, or detecting the differentiation of immune cells. Techniques known to those of skill in the art can be used for measuring these activities. For example, cellular proliferation can be assayed by ³H thymidine incorporation assays and trypan blue cell counts. Antigen expression can be assayed, for example, by immunoassays including, but are not limited to, competitive and non-competitive assay systems using techniques such as western blots, immunohistochemistry radioimmunoassays, ELISA (enzyme linked immunosorbent assay), "sandwich" immunoassays, immunoprecipitation assays, precipitin reactions, gel diffusion precipitin reactions, immunodiffusion assays, agglutination assays, complement-fixation assays, immunoradiometric assays, fluorescent immunoassays, protein A immunoassays and FACS analysis. The activation of signaling molecules can be assayed, for example, by kinase assays and electrophoretic shift assays (EMSAs).

[00400] Formulations of the invention comprising antibodies or compositions of the invention can also be tested for their ability to inhibit viral replication or reduce viral load in *in vitro*, *ex vivo* and *in vivo* assays. Antibodies or compositions of the invention can also be tested for their ability to decrease the time course of a RSV infection (*i.e.*, an upper and/or lower

respiratory tract RSV infection), otitis media (preferably stemming from, caused by or associated with a RSV infection, such as an upper and/or lower respiratory tract infection), or a symptom or respiratory condition relating thereto (including, but not limited to, asthma, wheezing, RAD, or a combination thereof). Antibodies or compositions of the invention can also be tested for their ability to increase the survival period of humans suffering from a RSV infection (preferably, an upper and/or lower respiratory tract RSV infection) by at least 25%, preferably at least 50%, at least 60%, at least 75%, at least 85%, at least 95%, or at least 99%. Further, antibodies or compositions of the invention can be tested for their ability reduce the hospitalization period of humans suffering from a RSV infection (preferably, an upper and/or lower respiratory tract RSV infection) by at least 60%, preferably at least 75%, at least 85%, at least 95%, or at least 99%. Techniques known to those of skill in the art can be used to analyze the function of the antibodies or compositions of the invention *in vivo*.

5.8 Diagnostic Uses of Antibodies for Detecting RSV Infections

Labeled antibodies and derivatives and analogs thereof, which [00401]immunospecifically bind to a RSV antigen can be used for diagnostic purposes to detect, diagnose, or monitor an upper and/or lower respiratory tract RSV infection or otitis media (preferably, stemming from, caused by or associated with a RSV infection, such as an upper and/or lower respiratory tract RSV infection). The invention provides for the detection of a RSV infection (i.e., an upper and/or lower respiratory tract RSV infection), otitis media (preferably stemming from, caused by or associated with a RSV infection, such as an upper and/or lower respiratory tract infection), or a symptom or respiratory condition relating thereto (including, but not limited to, asthma, wheezing, RAD, or a combination thereof) comprising: (a) assaying the expression of a RSV antigen in cells or a tissue sample of a subject using one or more antibodies that immunospecifically bind to the RSV antigen; and (b) comparing the level of the RSV antigen with a control level, e.g., levels in normal tissue samples not infected with RSV, whereby an increase in the assayed level of RSV antigen compared to the control level of the RSV antigen is indicative of a RSV infection (i.e., an upper and/or lower respiratory tract RSV infection), otitis media (preferably stemming from, caused by or associated with a RSV infection, such as an upper and/or lower respiratory tract infection), or a symptom or respiratory condition relating thereto (including, but not limited to, asthma, wheezing, RAD, or a combination thereof).

[00402] The invention provides a diagnostic assay for diagnosing a RSV infection (i.e., an upper and/or lower respiratory tract RSV infection), otitis media (preferably stemming from, caused by or associated with a RSV infection, such as an upper and/or lower respiratory tract

infection), or a symptom or respiratory condition relating thereto (including, but not limited to, asthma, wheezing, RAD, or a combination thereof) comprising: (a) assaying for the level of a RSV antigen in cells or a tissue sample of an individual using one or more antibodies that immunospecifically bind to a RSV antigen; and (b) comparing the level of the RSV antigen with a control level, e.g., levels in normal tissue samples not infected with RSV, whereby an increase in the assayed RSV antigen level compared to the control level of the RSV antigen is indicative of a RSV infection (i.e., an upper and/or lower respiratory tract RSV infection), otitis media (preferably stemming from, caused by or associated with a RSV infection, such as an upper and/or lower respiratory tract infection), or a symptom or respiratory condition relating thereto (including, but not limited to, asthma, wheezing, RAD, or a combination thereof). A more definitive diagnosis of a RSV infection (i.e., an upper and/or lower respiratory tract RSV infection), otitis media (preferably stemming from, caused by or associated with a RSV infection, such as an upper and/or lower respiratory tract infection), or a symptom or respiratory condition relating thereto (including, but not limited to, asthma, wheezing, RAD, or a combination thereof) may allow health professionals to employ preventative measures or aggressive treatment earlier thereby preventing the development or further progression of the RSV infection or otitis media.

[00403] Antibodies of the invention can be used to assay RSV antigen levels in a biological sample using classical immunohistological methods as described herein or as known to those of skill in the art (*e.g.*, see Jalkanen et al., 1985, J. Cell. Biol. 101:976-985; and Jalkanen et al., 1987, J. Cell . Biol. 105:3087-3096). Other antibody-based methods useful for detecting protein gene expression include immunoassays, such as the enzyme linked immunosorbent assay (ELISA) and the radioimmunoassay (RIA). Suitable antibody assay labels are known in the art and include enzyme labels, such as, glucose oxidase; radioisotopes, such as iodine (¹²⁵I, ¹²¹I), carbon (¹⁴C), sulfur (³⁵S), tritium (³H), indium (¹²¹In), and technetium (⁹⁹Tc); luminescent labels, such as luminol; and fluorescent labels, such as fluorescein and rhodamine, and biotin.

[00404] One aspect of the invention is the detection and diagnosis of a RSV infection (*i.e.*, an upper and/or lower respiratory tract RSV infection), otitis media (preferably stemming from, caused by or associated with a RSV infection, such as an upper and/or lower respiratory tract infection), or a symptom or respiratory condition relating thereto (including, but not limited to, asthma, wheezing, RAD, or a combination thereof) in a human. In one embodiment, diagnosis comprises: a) administering (for example, parenterally, subcutaneously, or

intraperitoneally) to a subject an effective amount of a labeled antibody that immunospecifically binds to a RSV antigen; b) waiting for a time interval following the administering for permitting the labeled antibody to preferentially concentrate at sites in the subject (e.g., the nasal passages, lungs, mouth and ears) where the RSV antigen is expressed (and for unbound labeled molecule to be cleared to background level); c) determining background level; and d) detecting the labeled antibody in the subject, such that detection of labeled antibody above the background level indicates that the subject has a RSV infection (i.e., an upper and/or lower respiratory tract RSV infection), otitis media (preferably stemming from, caused by or associated with a RSV infection, such as an upper and/or lower respiratory tract infection), or a symptom or respiratory condition relating thereto (including, but not limited to, asthma, wheezing, RAD, or a combination thereof). Background level can be determined by various methods including, comparing the amount of labeled molecule detected to a standard value previously determined for a particular system.

[00405] It will be understood in the art that the size of the subject and the imaging system used will determine the quantity of imaging moiety needed to produce diagnostic images. In the case of a radioisotope moiety, for a human subject, the quantity of radioactivity injected will normally range from about 5 to 20 millicuries of ⁹⁹Tc. The labeled antibody will then preferentially accumulate at the location of cells which contain the specific protein. *In vivo* tumor imaging is described in S.W. Burchiel et al., "Immunopharmacokinetics of Radiolabeled Antibodies and Their Fragments." (Chapter 13 in Tumor Imaging: The Radiochemical Detection of Cancer, S.W. Burchiel and B.A. Rhodes, eds., Masson Publishing Inc. (1982).

[00406] Depending on several variables, including the type of label used and the mode of administration, the time interval following the administration for permitting the labeled antibody to preferentially concentrate at sites in the subject and for unbound labeled antibody to be cleared to background level is 6 to 48 hours or 6 to 24 hours or 6 to 12 hours. In another embodiment the time interval following administration is 5 to 20 days or 5 to 10 days.

[00407] In one embodiment, monitoring of an upper and/or lower respiratory tract RSV infection is carried out by repeating the method for diagnosing the upper and/or lower respiratory tract RSV infection, for example, one month after initial diagnosis, six months after initial diagnosis, one year after initial diagnosis, etc.

[00408] Presence of the labeled molecule can be detected in the subject using methods known in the art for in vivo scanning. These methods depend upon the type of label used.

Skilled artisans will be able to determine the appropriate method for detecting a particular label. Methods and devices that may be used in the diagnostic methods of the invention include, but are not limited to, computed tomography (CT), whole body scan such as position emission tomography (PET), magnetic resonance imaging (MRI), and sonography.

[00409] In a specific embodiment, the molecule is labeled with a radioisotope and is detected in the patient using a radiation responsive surgical instrument (Thurston et al., U.S. Patent No. 5,441,050). In another embodiment, the molecule is labeled with a fluorescent compound and is detected in the patient using a fluorescence responsive scanning instrument. In another embodiment, the molecule is labeled with a positron emitting metal and is detected in the patient using positron emission-tomography. In yet another embodiment, the molecule is labeled with a paramagnetic label and is detected in a patient using magnetic resonance imaging (MRI).

5.9 <u>Kits</u>

[00410] The invention also provides a pharmaceutical pack or kit comprising one or more containers filled with one or more of the ingredients of the pharmaceutical formulation of the invention. Optionally associated with such container(s) can be a notice in the form prescribed by a governmental agency regulating the manufacture, use or sale of pharmaceuticals or biological products, which notice reflects approval by the agency of manufacture, use or sale for human administration.

In one embodiment, a kit comprises an antibody of the invention, preferably a purified antibody, in one or more containers. In a specific embodiment, the kits of the present invention contain a substantially isolated RSV antigen as a control. Preferably, the kits of the present invention further comprise a control antibody which does not react with the RSV antigen. In another specific embodiment, the kits of the present invention contain a means for detecting the binding of an antibody to a RSV antigen (e.g., the antibody may be conjugated to a detectable substrate such as a fluorescent compound, an enzymatic substrate, a radioactive compound or a luminescent compound, or a second antibody which recognizes the first antibody may be conjugated to a detectable substrate). In specific embodiments, the kit may include a recombinantly produced or chemically synthesized RSV antigen. The RSV antigen provided in the kit may also be attached to a solid support. In a more specific embodiment the detecting means of the above described kit includes a solid support to which RSV antigen is attached. Such a kit may also include a non-attached reporter-labeled anti-human antibody. In this

embodiment, binding of the antibody to the RSV antigen can be detected by binding of the said reporter-labeled antibody.

[00412]

6. EXAMPLES

[00413] The following examples are offered to illustrate this invention and not to be construed in any way as limiting the scope of this invention.

6.1 EXAMPLE: CHARACTERIZATION OF ANTIBODY FORMULATION FOR ANTIBODY FRAGMENTATION AND AGGREGATION

[00414] This example illustrates the characterization of an antibody formulation for antibody fragmentation and aggregation. Antibody A4B4L1FR-S28R is used in this example. As discussed in the sections above antibody A4B4L1FR-S28R is an IgG1 monoclonal antibody produced by recombinant DNA technology that specifically binds to an epitope in the A antigenic site of the fusion (F) protein of RSV. A4B4L1FR-S28R is a humanized antibody and consists of the CDR regions specific for the targeted antigen and the constant regions of a human $\gamma 1$ heavy chain and κ light chain. The monoclonal antibody has two inter-chain disulfide bonds to link heavy and light chains, and another two inter-chain disulfide bonds at the hinge region. Unless otherwise indicated, all antibody samples in this example were formulated at a concentration of 100 mg/ml in 25 mM histidine-HCl, pH 6.0. Further storage conditions are reported in the section describing experimental results.

Materials and Methods

Size Exclusion Chromotography (SEC)

[00415] Size exclusion chromatography was performed to analyze the antibody formulation for the presence of antibody aggregates and fragments. The test samples were injected onto a size exclusion G3000 SW_{XL} column (5 μ m, 300 Å, 7.8 x 300 mm, TosoHaas). The mobile phase was 0.1 M di-sodium phosphate, 0.1 M sodium sulphate and 0.05 % sodium azide (pH 6.7), running isocratically at a flow rate of 0.25 - 1.0 mL/min. Eluted protein was detected by UV absorbance at 280 nm and collected for further characterization. The relative amount of any protein species detected was reported as the area percent of the product peak as compared to the total area of all other detected peaks excluding the initial included volume peak. Peaks eluting earlier than the antibody monomer peak were recorded in the aggregate percentile, while peaks eluting later than the antibody monomer peak, but earlier than the buffer peak, were

recorded in the fragment percentile. The hydrodynamic radius and molecular weight of the individual peaks were obtained with a coupled multiangle light scattering detector.

Analytical Ultracentrifugation (AUC)

[00416] Analytical ultracentrifugation (AUC) was also used to characterize the antibody formulation. AUC is an orthogonal technique which determines the sedimentation coefficients (reported in Svedberg, S) of macromolecules in a liquid sample. Like SEC, AUC is capable of separating and detecting antibody fragments/aggregates from monomers and is further able to provide information on molecular mass. Compared to SEC, AUC eliminates the possibility of aggregate loss due to solid-phase interaction and is better able to resolve differing species of a given macromolecule.

[00417] Sedimentation velocity experiments were performed using a Beckman Optima XL-A analytical ultracentrifuge. Test samples were diluted to an antibody concentration of 0.5 mg/ml with reference buffer (20 mM citric acid, 100 mM NaCl, 1.5% mannitol, 50 μ M diethylenetriamine-pentaacetic acid, 0.02% Polysorbate 80, pH 6.0). 415 μ l of the diluted antibody sample and 412 μ l or the reference buffer were loaded into a 12 mm centrifuge cell in the sample and reference channels, respectively. Loaded cells were placed into an AN-50Ti analytical rotor and equilibrated to 25 °C. Samples were scanned at 280 nm with a rotor speed of 42000 rpm at full vacuum. A total of 80 scans for each cell were collected for analysis. The first scan for each sample was excluded to avoid artifacts caused by meniscus.

The data were analyzed using the c(s) method developed by Peter Shuck at N.I.H. and the SEDFIT (version 8.8) program with implemented c(s). Using the c(s) method, raw data scans are directly fit to a Lamm function of S in order to derive a distribution of sedimentation coefficients. The parameters used for the fitting procedure were resolution, 400; confidence interval, 0.75; grid size, 1000; partial specific volume, 0.7245; buffer density, 1.000; and buffer viscosity, 0.1002. Frictional ratio, meniscus and bottom positions were set as fitted parameters. Time independent noise was also fitted. The detected peaks were integrated and classified as follows: from 0 to 6 S, fragments; from 6 to 9 S, monomer; and from 9 to 20 S, aggregates.

Turbidity Measurement

[00419] Protein aggregation in the antibody formulation was also characterized by turbidity measurement. Turbidity is a measure of the amount by which the particles in a solution scatter light and, thus, may be used as a general indicator of protein aggregation or denaturation.

[00420] Approximately 3 to 4 ml of formulation sample was transferred into a glass test tube and degassed for 2 minutes using an in-line vacuum system. The degassed sample was then placed into a turbidimeter (2100AN or 2100N, Hatch) sample compartment at room temperature for analysis. The turbidimeter was calibrated with STABLCAL® Stabilized Formazin Turbidity standard (Hatch) at 40, 200, 1000 and 4000 NTU (nephelometric turbidity unit) and verified by analyzing control suspensions of formazin at 3, 6, 18, 30 and 60 NTU.

Results

[00421] SEC was used to monitor antibody aggregate and fragment formation in formulations of A4B4L1FR-S28R stored at three temperature ranges over the course of 9 months. Temperature ranges above the proposed storage temperature, 2-8 °C, were used to stress the formulation and were hoped to simulate the effects of prolonged storage. FIGS. 6 A, B and C present the relative percentage of monomer (purity), aggregates and fragments, respectively, for a single formulation of motavizumab stored at 2-8 °C, 20-24 °C and 38-42 °C. The relative percentage of fragmentation and aggregation increased with both time and temperature. For a single temperature range, however, both the fragmentation and aggregation rate were constant. This finding proved that a higher storage temperature would accurately simulate an accelerated time scale.

[00422] The logarithm of the estimated rates of fragmentation/aggregation also showed a linear dependence to the reciprocal of the storage temperature (FIG. 7). Once this linearity is established, it is then possible to predict the rate of aggregation/fragmentation of a given formulation at any temperature or, more importantly, the formulation characteristics at any time at such temperature.

[00423] FIG. 8 presents a representative SEC profile of the antibody formulation after storage at 38-42 °C with 70-80 % relative humidity for 1 month. Under these conditions, SEC was able to clearly separate antibody aggregates and fragments from monomers. However, at low relative levels of aggregates/fragments, the peaks identified as aggregates and fragment I in FIG. 8 begin to become less distinct and merge into the shoulders of the monomer peak. Such shoulders cannot be accurately analyzed.

[00424] As an alternative, AUC was investigated as a method to characterize low relative levels of aggregation and fragmentation in antibody formulations. FIG. 9 and Table 7 compare the AUC and SEC analysis of formulation samples at initial, 9 month and 14 month time points (the 9 and 14 month samples had been stored at 38-42 °C with 70-80 % relative humidity).

AUC identified two major fragmentation peaks at about 50 KDa and about 90 KDa. AUC was also able to better resolve the fragmentation and aggregation peaks. For the 9 month sample, SEC did not resolve the large fragment peak, while AUC was clearly capable of resolving it. For the 14 month sample, the large fragment peak in SEC was observed as a shoulder of the monomer peak and, when integrated, resulted in a higher fragment I percent than that determined by AUC. Aggregate values for AUC and SEC were comparable. AUC estimates of the molecular mass of the aggregate peak indicated that the majority of the aggregates were antibody dimers.

[00425] Compared to SEC, AUC is also able to better resolve differing species of a given macromolecule. It is, however, first necessary to establish the proper sample dilution, as the noise/signal ratio of AUC is dependent on the concentration of antibody in the sample (FIG. 10). For the described formulation of A4B4L1FR-S28R (100 mg/ml in 25 mM histidine-HCl, pH 6.0), a 200 fold dilution was used -- resulting in a sample antibody concentration of 0.5 mg/ml. Under these conditions, AUC was able to resolve the slight changes in formulation composition observed over 5 days of storage at 38-42 °C with 70-80 % relative humidity (FIG. 11).

TABLE 7. COMPARISON OF AUC AND SEC ANALYSIS OF motavizumab
FORMULATIONS AT INITIAL, 9-MONTH AND 14-MONTH TIME
POINTS

Samples		AUC		SEC			
	Fragments%	Monomer%	Aggregates%	Fragments%	Monomer%	Aggregates%	
Initial	0.0	99.2	0.8	0.0	99.5	0.5	
9-month	7.5	89.3	3.2	3.3	93.7	3.0	
14-month	24.5	64.7	10.8	28.8	60.5	9.8	

[00426] As a general indicator of protein aggregation, the antibody formulation may also be monitored for changes in turbidity. Four lots of a formulation containing concentrations of antibody at about 100 mg/ml were measured for turbidity using a HACH turbidimeter after storage at 38-42 °C for one month (Table 8). The results indicate that the turbidity levels of the differing lots of the formulation had comparable turbidity measurements, comparable NTU, but that one lot showed an elevated measurement. Elevated turbidity may indicate a higher level of aggregation or an increased number/ increased size of particles.

TABLE 8. TURBIDITY VALUES OF FOUR LOTS OF A motavizumab FORMULATION

MAb	Lot	Concentration (mg/ml)	Turbidity Value (NTU)	
	A	100	5.8	
	В	100	7.1	
A4B4L1FR-S28R	C	100	6.1	
	D	100	5.6	
	E	100	5.7	

6.2 EXAMPLE: CHARACTERIZATION OF ANTIBODY FRAGMENTS AND FORMULATION PARTICLE SIZE DISTRIBUTION

[00427] This example illustrates the characterization of antibody fragments as identified by AUC or SEC. Antibody A4B4L1FR-S28R is used in this example. Unless otherwise indicated, all antibody samples in this example were formulated at a concentration of 100 mg/ml in 25 mM histidine-HCl, pH 6.0. Further storage conditions are reported in the section desribing experimental results.

Materials and Methods

Liquid Chromatography Mass Spectrometry (LC-MS)

In SEC fragment peaks were collected and digested with N-Glycosidase F, also known as PNGase F, at 37 °C overnight. PNGase F is an amidase that cleaves between the innermost GlcNAc and asparagine residues of high mannose, hybrid and complex oligosaccharides on N-linked glycoproteins. The deglycosylated sample (approximately 7.5 μ L) was mixed with approximately 42.5 μ L of reducing buffer (2.5 mg/mL DTT, 6.0 M guanindine HCl, pH 8.2) and kept at 56 °C in a water bath for 60 minutes. Neat 4-vinylpyridine (Aldrich Chem. Co., WI) (approximately 0.5 μ L) was then added to the sample, and the reaction mixture was held at ambient temperature for 30 minutes. The deglycosylated, reduced and alkylated

sample was immediately loaded onto a reversed phase column to separate the modified samples from the reactants, and to analyzed by LC-ESI-MS.

[00429] Deglycosylated, reduced, and alkylated samples were fractionated using a reversed phase column (Jupiter 5μm C4, 300 Å, 250 x 2.00 mm, Phenomenex) with a binary gradient HPLC system (Agilent 1100). Mobile phase A consisted of 30% acetonitrile in water with 0.1% trifluoroacetic acid and mobile phase B consisted of 50% acetonitrile in water with 0.1% trifluoroacetic acid. The samples were separated using a linear gradient of 30-50% acetonitrile in water, over 16 min. with a flow rate of 200 μL/min. The column effluent was directed to a UV detector and then split 1:1, one half going to a switching valve on the Ion Trap mass spectrometer (LTQ, ThermoElectro, San Jose, CA), and the remaining half to waste. The switching valve diverted the column effluent flow to the mass spectrometer only between the 15 and 30 minutes portion of the chromatographic run.

[00430] A mixture of caffeine, L-methionyl-arginyl-phenylalanyl-alanine acetate H₂O, and Ultramark 1621 was used to calibrate the ion-trap mass spectrometer according to the manufacturer's instruction. The ESI-MS data were acquired in positive ESI full scan mode. The BioWork deconvolution program (ThermoFinnigan) was used to reconstruct the mass spectra and obtain the molecular masses of the peptides/proteins from their original mass spectra.

Disulfide Bond Determination

[00431] Test samples of antibody were denatured in 10 mM phosphate buffer, 250 mM NaCl, 5 mM NEM, 6 M Guanidine, pH 7.0 at 37 °C for 1 to 3 hr. The denatured samples were then diluted 6 fold with 100 mM phosphate buffer, 0.1 mM EDTA, pH 7.0, to which Lys-C was added at a 1:10 enzyme to protein ratio. The reaction mixtures were incubated at 37 °C for 16 to 24 hours. Half of the reaction mixture was reduced by adding 5-10 μ L of 100 mM DTT and incubated at 37 °C for 1 hr.

[00432] Lys-C digests were separated by reverse-phase HPLC (Phenomenex Jupiter 5m C18 column; 250 x 2.1 mm) and analyzed by an UV-detector and an on-line LCQ or LTQ Ion Trap mass spectrometer (ThermoElectron). The RP-HPLC mobile phase A was 0.1% TFA in H2O and the mobile phase B was 0.1% TFA in acetonitrile. The peptides were eluted at a flow rate of 0.2 mL/min with the following gradient:

[**00433**] 0-2 min, 5% Mobile Phase B

[00434] 2-32 min, 5-20% Mobile Phase B

[00435] 32-132 min, 20-40% Mobile Phase B

[00436] 132-152 min, 40-60% Mobile Phase B

[00437] 152-155 min, 60-95% Mobile Phase B

[00438] The column eluant was diverted to waste directly after the UV-detector during the first 15 min to avoid salt contamination of the LCQ source.

Particle Counting

[00439] The number and size of particles in a solution was characterized by a Beckman Coulter Multisizer 3.

Results

[00440] To characterize aggregates and fragments identified by SEC, fragment fractions were collected from the SEC chromatographic system and analyzed by LC-MS (antibody fragment I and antibody fragment II, FIGS. 12 and 13, respectively). The predominant fragments, above the detection limit of LC-MS, were identified for both fragment peaks (antibody type I fragment and antibody type II fragment) (FIG. 14 and Table 9). Antibody Type I and Antibody Type II fragments were generated by cleavage of the heavy chain in one of the hinge regions of the antibody. Observed cleavage sites were between serine 222 and cysteine 223, cysteine 223 and aspartic acid 224, between aspartic acid 224 and lysine 225, between lysine 225 and threonine 226, between threonine 226 and histidine 227, between histidine 227 and threonine 228, and between threonine 228 and cysteine 229.

[00441] A comparison of peptide maps using reduced and non-reduced conditions of LC-MS/MS was also used to detect disulfide bond scrambling or other covalent modification in the monoclonal antibodies. The profile comparison for aggregates, monomer and fragments indicates that only a low level of disulfide bond scrambling existed in the aggregates (FIGS. 15 and 16). The results also suggest that most of the aggregates were non-covalently linked aggregates, as no significant profile change compared to that of monomer was observed.

Table 9. LC MS IDENTIFICATION OF motavizumab FRAGMENTS AFTER STORAGE OF ANTIBODY FORMULATION AT 38-42°C FOR 1 MONTH

	Sequence	Calculated MW	Measured MW	MW Accuracy
Defende	Light Chain	23654.1	23654.9	0.0000
Reference	Heavy Chain	50617.68	50619.6	0.004%
Standard				
	Light Chain	23654.1	23655.2	0.0000
	H1-222+O	24360.69	24364.6	0.016%
	H1-223+O	24568.97	24571.1	0.009%
	H1-224+O	24684.06	24686	0.008%
	H1-226+O	24913.34	24913.5	0.001%
F.,, .,,, ., ., 4 II	H1-227+O	25050.48	25053.2	0.011%
Fragment II		***************************************		
	H1-222	24344.69	24346.2	0.006%
	H1-223	24552.97	24554.9	0.008%
	H1-224	24668.06	24671.5	0.014%
	H1-226	24897.34	24899.9	0.010%
	H1-227	25034.48	25037.9	0.014%
	Light Chain	23654.1	23655.2	0.0000
	H228-449+O	25599.2	25604.6	0.021%
	H227-449+O	25736.34	25742	0.022%
F., 1	H226-449+O	25837.44	25843.3	0.023%
Fragment I	H225-449+O	25965.61	25972.7	0.027%
	H224-449+O	26080.7	26085.7	0.019%
	H1-449+O	50633.68	50640.7	0.014%
	H1-449	50617.68	50624.1	0.013%

[00442] A multisizer was also used to characterize the particle size distribution of the antibody formulation. A test sample of formulation at 100 mg/ml was analyzed in a Beckman Coulter Multisizer 3 (Table 10).

TABLE 10. PARTICLE ANALYSIS OF motavizumab SAMPLE AFTER STORAGE OF ANTIBODY FORMULATION AT 38-42°C FOR 1 MONTH

Size	Dilution 1,run 1	Dilution 1,run 2	Dilution 1,run 3	Dilution 2,run 1	Dilution 2,run 2	Dilution 2,run 3	Average
μm	Particle/mL	Particle/mL	Particle/mL	Particle/mL	Particle/mL	Particle/mL	Particle/mL
2-4	3.08E+05	3.11E+05	3.09E+05	2.81E+05	2.83E+05	2.82E+05	2.96E+05
4-10	3.93E+04	3.79E+04	3.75E+04	3.61E+04	3.54E+04	3.32E+04	3.66E+04
10-20	3.33E+03	3.47E+03	2.69E+03	6.11E+03	3.71E+03	3.74E+03	3.84E+03
20-30	5.97E+02	3.06E+02	2.55E+02	1.01E+03	3.06E+02	2.40E+02	4.52E+02
30-40	1.02E+02	5.10E+01	0.00E+00	1.48E+02	5.10E+01	5.10E+01	6.72E+01
40-60	5.10E+01	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	8.50E+00
2-60 total	3.51E+05	3.52E+05	3.50E+05	3.25E+05	3.22E+05	3.19E+05	3.37E+05

6.3 <u>KINETIC ANALYSIS OF BINDING OF</u> <u>A4B4L1FR-S28R BY BIACORETM</u>

[00443] The kinetics of the interactions of A4B4L1FR-S28R and palivizumab with RSV F-protein were determined by surface plasmon resonance (see, *e.g.*, Jonsson et al., 1991, Biotechniques 11(5):620-627 and Johne, B. (1989). Epitope mapping by surface plasmon

resonance in the BIAcore. Molecular Biotechnology 9(1):65-71) using a BIAcore 3000 instrument (BIAcore, Inc., Piscataway, NJ). A recombinantly produced, C-terminally truncated RSV (A2 strain) F protein (Wathen et al., 1989, J Infect Dis 159(2):255-264) was used as the antigen for these studies. The truncated F protein, lacking the membrane anchor, was produced as a secreted product using a recombinant baculovirus expression system and was purified by successive chromatography steps on concanavalin-A and Q-sepharose columns. Purified F protein was covalently coupled to an N-hydroxysuccinimide-N-ethyl-N'-[3-diethylaminopropyl]-carbodiimide (EDC/NHS) activated CM5 sensor chip at a low protein density according to the manufacturer's protocol; unreacted active ester groups were blocked with 1 M ethanolamine. For reference purposes, a blank surface, containing no antigen, was prepared under identical immobilization conditions.

[00444] For kinetic measurements, a serial 2-fold dilution series of each mAb from 100 nm - 0.2 nm, made in instrument buffer (HBS/Tween-20, BIAcore, Inc.), was injected over the F-protein and reference cell surfaces, which are connected in series. In each analysis, following the dissociation phase, the remaining bound antibody was removed from the sensor chip by passing a brief pulse of 100 mM HCl over the surface. Once an entire data set was collected, the resulting binding curves were globally fitted to a 1:1 Langmuir binding model using BIAevaluation software (BIAcore, Inc., Piscataway, NJ). This algorithm calculates both the association rate (k_{on}) and the dissociation rate (k_{off}), from which the apparent equilibrium binding constant, K_{D} , for each antibody was deduced as the ratio of the two rate constants, k_{off} / k_{on} . A more detailed explanation of how the individual rate constants are derived can be found in the BIAevaluation Software Handbook (BIAcore, Inc., Piscataway, NJ).

[00445] Kinetic analysis of binding by BIAcore evaluation (Table 11) revealed that, under the conditions of a low-density surface that were employed, A4B4L1FR-S28R (motavizumab) had an approximately 70-fold greater affinity for RSV F protein than palivizumab. The increased affinity of motavizumab for the RSV F protein is attributed to a 4-fold increase in the association rate and an approximately 17-fold decrease in the dissociation rate. Since the rate at which motavizumab dissociates from the F protein surface approaches the detection limits of the BIAcore 3000 instrument, the dissociation rate generated for motavizumab is an estimation.

Table 11. Kinetic Analysis of Binding

mAb	$k_{on} (\mathrm{M}^{-1} \mathrm{s}^{-1})$	k_{off} (s ⁻¹)	K _D (pM)	
palivizumab	1.14 E+05	3.95 E-04	3460	
motavizumab	4.73 E+05	2.35 E-05	50	

6.4 EXAMPLE: MICRONEUTRALIZATION ASSAY

Neutralization of the antibodies of the present invention were determined by [00446] microneutralization assay. This microneutralization assay is a modification of the procedures described by Anderson et al. (1985, J. Clin. Microbiol. 22:1050-1052, the disclosure of which is hereby incorporated by reference in its entirety). The procedure used here is described in Johnson et al., 1999, J. Infectious Diseases 180:35-40, the disclosure of which is hereby incorporated by reference in its entirety. Antibody dilutions were made in triplicate using a 96well plate. Ten TCID₅₀ of respiratory syncytial virus (RSV – Long strain) were incubated with serial dilutions of the antibody (or Fabs) to be tested for 2 hours at 37° C in the wells of a 96well plate. RSV susceptible HEp-2 cells (2.5 x 10⁴) were then added to each well and cultured for 5 days at 37° C in 5% CO₂. After 5 days, the medium was aspirated and cells were washed and fixed to the plates with 80% methanol and 20% PBS. RSV replication was then determined by F protein expression. Fixed cells were incubated with a biotin-conjugated anti-F protein monoclonal antibody (pan F protein, C-site-specific mAb 133-1H) washed and horseradish peroxidase conjugated avidin was added to the wells. The wells were washed again and turnover of substrate TMB (3,3',5,5'-tetramethylbenzidine) was measured at 450 nm. The neutralizing titer was expressed as the antibody concentration that caused at least 50% reduction in absorbency at 450 nm (the OD₄₅₀) from virus-only control cells. The results from the assay for the monoclonal antibodies and Fab fragments listed in Table 2 are shown in Table 11, supra, and Table 12, infra.

Table 12. End Point RSV Microneutralization Titer Of High On Rate Mutant IgG and Fab

	Mean		Fold	Mean		Fold	
	IC50	STDEV	Difference	IC50	STDEV	Difference	
	(Curve)	Curve	(Curve	(Control)	Control	(Control	n (assay
Molecule	μg/ml	IC50	IC50)	μ g/ml	IC50	IC50)	repeat)
**palivizumab	0.4527	0.208	-	0.5351	0.238	-	8
**A1e9	0.0625	0.0268	7	0.0645	0.223	8	3
**A17d4(1)	0.0342	0.022	13	0.0354	0.0187	15	4
**P11d4	0.0217	0.0331	21	0.0289	0.0110	19	5
**P12f2	0.0231	0.0141	20	0.0223	0.0083	24	6
**A8c7	0.0337	0.0309	13	0.0383	0.0283	14	5
**A12a6	0.0357	0.0316	13	0.0354	0.0261	15	7
**P12f4	0.0242	0.0163	19	0.0235	0.0076	23	7
**A13c4	0.0376	0.0268	12	0.0375	0.0213	14	6
**A4B4	0.0171	0.0018	27	0.0154	0.00417	35	2
*A1e9	0.157	-	3	0.125	-	4	1
*A17d4(1)	0.0179	-	25	0.0171	-	31	1
*P11d4	>1.00	-	-	>1.00	-	-	1
*P12f2	0.0407	0.0112	11	0.0326	0.00905	16	2
*A8c7	0.177	-	3	0.157	_	34	1
*A12a6	0.0287	0.00417	16	0.0310	0.00982	17	2
*P12f4	0.0464	0.00791	10	0.0351	0.0126	15	2

WO 2007/002543						PCT/US2006/024717		
*A13c4	0.0264	0.00141	17	0.0258	0.00071	21	2	
*A4B4	0.0414	-	11	0.0411	-	13	1	
*A13a11	0.120	0.0222	4	0.1022	0.0260	5	2	
*A1h5	0.194	0.462	2	0.176	0.0625	3	2	
** Monoclo	nal Antibody							

^{*} Fab Fragment

6.5 RSV MICRONEUTRALIZATION ASSAY

[00447] The ability of A4B4L1FR-S28R (motavizumab) and palivizumab to inhibit the in vitro replication of RSV (Long strain) was evaluated using a RSV microneutralization assay. This assay is a modification of the procedure of Anderson et al., (Anderson et al., 1985, J Clin Microbiol 22: 1050-1052) as described by Johnson et al., (Johnson et al., 1997, J Infect Dis 176: 1215-1224). Antibody dilutions were made in duplicate to quadruplicate wells of a 96-well plate. Approximately 100-1000 TCID₅₀ of RSV (Long) were added to each dilution well and incubated for two hours at 37°C. Low passage, RSV susceptible HEp-2 cells (2.5x10⁴) were then added to each well and cultured for five days at 37° C in a humidified 5%CO₂ incubator. After four or five days the cells were washed with PBS - 0.1% Tween 20 and fixed to the plate with 80% acetone with 20% PBS. RSV replication was determined by quantitation of F protein expression using an F protein-specific ELISA. Fixed cells were incubated with the C-site specific, pa RSV F protein mAb 133-1H (Chemicon, Inc.), washed, and then incubated with horseradish peroxidase-conjugated goat anti-mouse IgG and washed again. The peroxidase substrate TMB (3,3',5,5'-tetramethylbenzidine) was added to each well and the reaction was stopped after twenty minutes by the addition of 2 M H₂SO₄. Substrate turnover was measured at 450 nm (OD450) using a microplate reader. The neutralizing titer is expressed as the antibody concentration resulting in at least a 50% reduction in the OD450 value from control wells with virus only (IC₅₀). The results of this assay, shown in FIG. 17, indicate that motavizumab (average $IC_{50} = 18 \text{ ng/ml}$) is approximately 18-fold more potent than palivizumab (average IC_{50} = 315 ng/ml).

6.6 RSV MICRONEUTRALIZATION ASSAY WITH CYNOMOLGUS BAL SAMPLES

[00448] The ability of motavizumab present in the lungs of treated animals to inhibit the *in vitro* replication of RSV was evaluated using the RSV microneutralization assay. Four juvenile female cynomolgus monkeys (average weight 2.0 kg) were sedated with Telazol and dosed intravenously (i.v.) with motavizumab at 30 mg/kg body weight via the saphenous vein using an external infusion pump. Four days later, the animals were anesthetized with Telazol

and a bronchial alveolar lavage (BAL) was performed on one lobe of the right lung with phosphate buffered saline (PBS). Titers of motavizumab in the BAL fluid were determined using a motavizumab-specific ELISA. The BAL samples were tested undiluted and at serial 2-fold dilutions in the RSV microneutralization assay as above with purified motavizumab included as a control. The results of this assay, shown in FIG. 18, show that motavizumab retains full RSV neutralizing activity in the lungs of cynomolgus monkeys four days after infusion.

6.7 RSV FUSION INHIBITION ASSAY

[00449] The ability of the antibodies of the invention to block RSV-induced fusion after viral attachment to the cells is determined in a fusion inhibition assay. This assay is identical to the microneutralization assay, except that the cells are infected with RSV (Long) for four hours prior to addition of antibody (Taylor et al., 1992, J. Gen. Virol. 73:2217-2223).

6.8 PHYSICAL CHARACTERIZATION

[00448] The example illustrates the physical characteristics of motavizumab and palivizumab. A number of parameters were examined including the Tm and pI. In addition, the aggregation rates and viscosity profiles of motavizumab and palivizumab were determined.

Materials and Methods

Generation of Antibody Fragments

[00449] Fab and Fc domains were generated from full length palivizumab antibody using papain. A commercial kit from Pierce (ImmunoPure Fab Preparation Kit Pierce Product # 44885: ImmunoPure IgG Binding Buffer, ImmunoPure IgG Elution Buffer, AffinityPak Immobilized Protein A Column, Immobilized Papain, Cysteine monohydrochloride, Phosphate Buffer, and Serum Separators) was used to digest the intact antibodies. The enzymology was optimized to achieve the best cleavage of the Mab in a reasonable time. Fab and Fc domains were generated from palivizumab using the following steps: a) adding antibody to papain and incubating overnight at 37°C, ~10 mg of IgG per digestion; b) separating crude digest from immobilized enzyme; c) applying digest to Protein A column; d) eluting the Fab fragment in unretained fraction at pH-8.0; e) eluting the Fc fragment at pH-3.0; and f) dialyzing the fragments into a required buffer.

Differential Scanning Calorimetry

[00450] Thermal melting temperatures (T_m) were measured with a VP-DSC (MicroCal, LLC) using a scan rate of 1.0°C/min and a temperature range of 25 –120°C. A filter period of 8 seconds was used along with a 5 minute pre-scan thermostating. Samples were prepared by dialysis into 10 mM Histidine-HCl, pH 6 using Pierce dialysis cups (3.5 kD). Average Mab concentrations were 50 μ g/mL as determined by A_{280} . Melting temperatures were determined following manufacturer procedures using Origin software supplied with the system. Briefly, multiple baselines were run with buffer in both the sample and reference cell to establish thermal equilibrium. After the baseline was subtracted from the sample thermogram, the data were concentration normalized and fitted using the deconvolution function.

Isoelectric Focusing Gel Electrophoresis

[00451] Isoelectric points were determined using a Pharmacia Biotech Multiphor 2 electrophoresis system with a multi temp 3 refrigerated bath recirculation unit and an EPS 3501 XL power supply. Pre-cast ampholine gels (Amersham Biosciences, pI range 2.5-10) were loaded with 5 μg of protein. Broad range pI marker standards (Amersham, pI range 3-10, 8 μL) were used to determine relative pI for the Mabs. Electrophoresis was performed at 1500 V, 50 mA for 105 minutes. The gel was fixed using a Sigma fixing solution (5x) diluted with purified water to 1x. Staining was performed overnight at room temperature using Simply Blue stain (Invitrogen). Destaining was carried out with a solution that consisted of 25% ethanol, 8% acetic acid and 67% purified water. Isoelectric points were determined using a Bio-Rad Densitometer relative to calibration curves of the standards.

Viscosity Profile

[00452] Viscosities of mAB solutions were measured using a ViscoLab 4000 Viscometer System (Cambridge Applied Systems) equipped with a ViscoLab Piston (SN:7497, 0.3055", 1-20 cP) and S6S Reference Standard (Koehler Instrument Company, Inc.). The viscometer was connected to a water bath and equilibrate the system to 20°C. Piston was checked using S6S viscosity reference standard (8.530 cP @ 20.00°C). Piston was also checked using RODI H_2O (1.00 cP @ 20.0°C). The piston was cleaned and rinsed thoroughly with soap and water between measurements of each different solution type. Each Mab was in 10 mM Histidine-HCl, pH 6 at a concentration of 100 mg/mL. The system was then cooled to \leq 2°C. When the system temperature was at or below 2°C, sample was loaded into the chamber and the piston was lowered into the sample. After sample was equilibrated to the temperature of the chamber, measurement was initiated. The temperature was increased at 1°C increments every 7-10

minutes to a final temperature of \geq 25°C. The temperature was adjusted on the water bath but the recorded temperature was what was displayed on the viscometer. The viscosity result was recorded immediately prior to increasing the temperature. The piston remained in motion during measurements to minimize the need for re-equilibration.

Aggregation Rate

[00453] Aggregation profiles over a range of temperatures were determined by HPSEC. Specifically, approximately 250 µg of, for example, the antibody or antibody fragment that immunospecifically binds to a target antigen (approximately 25 µl of a liquid formulation comprising 10 mg/ml said antibody or antibody fragment) was injected onto a TosoH Biosep TSK G3000SWXL column (7.8 mm x 30 cm) fitted with a TSK SW x1 guard column (6.0 mm CX 4.0 cm). The antibody or antibody fragment was eluted isocratically with 0.1 M disodium phosphate containing 0.1 M sodium sulfate and 0.05% sodium azide, at a flow rate of 0.8 to 1.0 ml/min. Eluted protein was detected using UV absorbance at 280 nm. A suitable reference standard was run in the assay as a control, and the results were reported as the area percent of the product monomer peak compared to all other peaks excluding the included volume peak observed at approximately 12 to 14 minutes. Peaks eluting earlier than the monomer peak were recorded as percent aggregate.

Results

[00454] Differential Scanning Calorimetry (DSC) was used to examine the melting curve of the full length palivizumab (FIG. 19, top). Fab and Fc domain fragments were generated from palivizumab and the purified fragments were analyzed individually by DSC (FIG. 19, bottom). The results show that individual Tm peaks in a full antibody may be assigned to individual domains. In particular, the largest peak represents the Tm of the Fab portion of a full length antibody. The Tm of the palivizumab Fab is about 87.6 °C.

[00455] A similar analysis was performed on motavizumab (data not shown). The Tm of the motavizumab Fab was found to be significantly higher, about 93.1. This finding is unexpected as these two molecules differ by only 13 amino acids.

[00456] To further characterize these molecules, the pI for each full length mAb was determined by isoelectric focusing gel electrophoresis. motavizumab has a pI of 9.0 and palivizumab was found to have a pI of 9.1. The Fab-Tm and mAb-pI values for each antibody are plotted in FIG. 20 for comparison.

[00457] The viscosities of 100 mg/ml solutions of motavizumab and palivizumab were respectively examined over a range of temperatures from about 2 to about 25 °C. The viscosity of motavizumab ranged from a high of about 6.0 cP at 2 °C to a low of about 3.0 cP at about 25

°C. The viscosity of palivizumab ranged from a high of about 4.5 cP at 2 °C to a low of about 2.0 cP at about 25 °C (FIG. 21).

[00458] The aggregation rates of palivizumab and motavizumab were plotted against the Fab Tm for each antibody (FIG. 22). A correlation between Fab Tm and reduced aggregation rates is seen. motavizumab, which has a significantly higher Fab Tm, is much less prone to forming aggregates than palivizumab.

7. EQUIVALENTS

[00459] Those skilled in the art will recognize, or be able to ascertain using no more than routine experimentation, many equivalents to the specific embodiments of the invention described herein. Such equivalents are intended to be encompassed by the following claims.

[00460] All publications, patents and patent applications mentioned in this specification are herein incorporated by reference into the specification to the same extent as if each individual publication, patent or patent application was specifically and individually indicated to be incorporated herein by reference.

WHAT IS CLAIMED IS:

1. An antibody formulation comprising a full length IgG₁ antibody, which antibody immunospecifically binds to an RSV antigen and is not palivizumab, wherein (i) within a predetermined period of time after production no more than a predetermined percentage of the total protein fraction of said formulation is antibody type I and antibody type II fragments, wherein said predetermined period of time is at least about 1 week, and wherein said predetermined percentage is about 0.5%; or (ii) within a month after production and under a temperature of 38-42° C. and a pH of 6.0, less than 5 % of the total protein fraction of said formulation comprises antibody aggregates as determined by size exclusion chromatography (SEC) with UV detection.

- 2. The formulation of claim 1, wherein within a predetermined period of time after production no more than a predetermined percentage of the total protein fraction of said formulation is antibody type I and antibody type II fragments, wherein said predetermined period of time is at least about 1 week and wherein said predetermined percentage is about 0.5%.
 - 3. The formulation of claim 1 or 2, wherein said RSV antigen is an F protein epitope.
- 4. The formulation of claim 1 or 2, wherein said RSV antigen comprises the F protein epitope NSELLSLINDMPITNDQKKLMSNN (SEQ ID NO:337).
- 5. The formulation of claim 1 or 2, wherein said RSV antigen consists of the F protein epitope NSELLSLINDMPITNDQKKLMSNN (SEQ ID NO:337).
- 6. The formulation of any one of claims 1 to 5, wherein the antibody competitively inhibits the binding of antibody A4B4L1FR-S28R to said RSV antigen.
- 7. The formulation of any one of claims 1 to 6, wherein the antibody comprises at least one variable heavy (VH) CDR of the antibody A4B4L1FR-S28R, at least two variable heavy (VH) CDRs of the antibody A4B4L1FR-S28R or at least three variable heavy (VH) CDRs of the antibody A4B4L1FR-S28R.

8. The formulation of any one of claims 1 to 6, wherein the antibody comprises at least one variable light (VL) CDR of the antibody A4B4L1FR-S28R, at least two variable light (VL) CDRs of the antibody A4B4L1FR-S28R or at least three variable light (VL) CDR of the antibody A4B4L1FR-S28R.

- 9. The formulation of any one of claims 1 to 7, wherein the antibody comprises a VH domain of the antibody A4B4L1FR-S28R (SEQ ID NO.:48).
- 10. The formulation of any one of claims 1 to 6 and 8, wherein the antibody comprises a VL domain of the antibody A4B4L1FR-S28R (SEQ ID NO.:11).
- 11. An antibody formulation comprising at least 100 mg/ml of a full length IgG₁ antibody comprising a heavy chain having the variable heavy (VH) domain of A4B4L1FR-S28R (SEQ ID NO:48) and a light chain having the variable light (VL) domain of A4B4L1FR-S28R (SEQ ID NO:11), wherein, within 1 month after production and under 38-42° C and pH 6.0, no more than 0.5 % of the total protein fraction of said formulation is antibody type I fragment.
- 12. An antibody formulation comprising at least 100 mg/ml of a full length IgG₁ antibody comprising a heavy chain having the variable heavy (VH) domain of A4B4L1FR-S28R (SEQ ID NO:48) and a light chain having the variable light (VL) domain of A4B4L1FR-S28R (SEQ ID NO:11), wherein, within 1 month after production and under 38-42° C and pH 6.0, no more than 0.5 % of the total protein fraction of said formulation is antibody type II fragment.
- 13. The antibody formulation of any one of claims 1 to 12, wherein each said antibody type I fragment comprises a heavy chain C-terminal portion, said heavy chain C-terminal portion has a molecular weight of about 25.6 kD, about 25.7 kD, about 25.8 kD, about 26.0 kD, or about 26.1 kD as determined by Liquid Chromatography Mass Spectrometry (LC-MS) analysis of samples of said stored antibody that have been deglycosylated, reduced and alkylated.
- 14. The antibody formulation of any one of claims 1 to 12, wherein each said antibody type II fragment comprises a heavy chain N-terminal portion, said heavy chain N-

terminal portion has a molecular weight of about 24.4 kD, about 24.6 kD, about 24.7 kD, about 24.9 kD, or about 25.1 kD as determined by LC-MS analysis of samples of said stored antibody that have been deglycosylated, reduced and alkylated.

- 15. The formulation of any one of claims 1 to 14, wherein within a month after production and under a temperature of 38-42° C. and a pH of 6.0, less than 5 % of the total protein fraction of said formulation comprises antibody aggregates as determined by size exclusion chromatography (SEC) with UV detection.
- 16. The formulation of any one of claims 1 to 15, wherein within a month after production and under a temperature of 38-42° C. and a pH of 6.0, the turbidity value of a degassed sample of said formulation is less than about 6.5 NTU.
- 17. The formulation of any one of claims 1 to 16, wherein within a month after production and under a temperature of 38-42° C. and a pH of 6.0, said formulation comprises a particle profile of less than about 3.4 E +5 particles/ml of diameter 2-4 μ m, less than about 4.0 E +4 particles/ml of diameter 4-10 μ m, less than about 4.2 E +3 particles/ml of diameter 10-20 μ m, less than about 5.0 E +2 particles/ml of diameter 20-30 μ m, less than about 7.5 E +1 particles/ml of diameter 30-40 μ m, and less than about 9.4 particles/ml of diameter 40-60 μ m as determined by a multisizer.
- 18. The formulation of any one of claims 1 to 17, within a month after production and under a temperature of 38-42° C. and a pH of 6.0, said antibody type I fragments comprise one or more C-terminal portions of said heavy chain, which heavy chain C-terminal portion comprises amino acid residues 223-449 of said antibody; amino acid residues 224-449 of said antibody, amino acid residues 225-449 of said antibody, amino acid residues 227-449 of said antibody, amino acid residues 228-449 of said antibody and amino acid residues 229-449 of said antibody.
- 19. The formulation of any one of claims 1 to 18, wherein each said antibody type II fragment comprises a heavy chain N-terminal portion, which heavy chain N-terminal portion comprises amino acid residues 1-222 of said antibody, amino acid residues 1-223 of said antibody, amino acid residues 1-225 of said

antibody, amino acid residues 1-226 of said antibody, animo acid residues 1-227 of said antibody or amino acid residues 1-228 of said antibody.

- 20. The formulation of any one of claims 1 to 19, wherein said formulation further comprises histidine.
- 21. The formulation of claim 20, wherein histidine is at a concentration of about 1 mM to about 10 mM or of about 10 mM to about 50 mM.
- 22. The formulation of claim 20, wherein the histidine is at a concentration of about 20 mM to about 30 mM, and wherein said formulation further comprises glycine at a concentration of less than 2 mM and is substantially free of surfactants, inorganic salts or other excipients.
- 23. The formulation of claim 22, wherein histidine is at a concentration of about 25 mM and glycine is at a concentration of about 1.6mM.
- 24. The formulation of claim 20, wherein said formulation is substantially free of surfactants and inorganic salts.
- 25. The formulation of claim 20, wherein said formulation is substantially free of other excipients.
- 26. The formulation of any one of claims 1 to 19, wherein said formulation further comprises an excipient other than a surfactant.
 - 27. The formulation of claim 26, wherein the excipient is glycine.
- 28. The formulation of claim 27, wherein glycine is at a concentration of less than 150 mM, less than 100 mM, less than 50 mM, less than 3 mM or less than 2 mM.
 - 29. The formulation of claim 36, wherein the formulation has a pH of about 6.0.
 - 30. The formulation of claim 26, wherein the excipient is a saccharide.

- 31. The formulation of claim 30, wherein the saccharide is sucrose.
- 32. The formulation of claim 31, wherein the sucrose is at a concentration of about 1% to about 20%.
- 33. The formulation of claim 26, wherein the excipient is a polyol other than mannitol.
 - 34. The formulation of claim 33, wherein the polyol is polysorbate.
- 35. The formulation of claim 33, wherein the polyol is Tween, which is at a concentration of about 0.001% to about 1%.
- 36. The formulation of any one of claims 1 to 35, wherein said formulation has a pH of between about 5.5 to about 7.0.
- 37. The formulation of claim 36, wherein said formulation has a pH of between about 5.5 to about 6.5.
- 38. The formulation of any one of claims 1 to 40, wherein the antibody is at a concentration of at least 5 mg/ml, at least 10 mg/ml, at least 20 mg/ml, at least 50 mg/ml, at least 100 mg/ml, at least 110 mg/ml, at least 150 mg/ml or at least 160 mg/ml.
- 39. The formulation of any one of claims 1 to 38, wherein said antibody formulation comprises an antibody manufactured and purified by a process comprising (a) transformation of a murine myeloma cell line with a recombinant vector capable of directing transcription of mRNA encoding said antibody; (b) maintenance of transformed cells; (c) collection of conditioned media from cultures of transformed cells; (c) cation/anion exchange chromatography; (e) nanofiltration; and (f) hydroxyapatite chromatography.
- 40. The formulation of claim 1, wherein the antibody comprises at least one CDR of the antibody A4B4L1FR-S28R, at least two CDRs of the antibody A4B4L1FR-S28R, at least three CDRs of the antibody A4B4L1FR-S28R, at least four CDRs of the antibody A4B4L1FR-S28R, at least five CDRs of the antibody A4B4L1FR-S28R or at least six CDRs of the antibody A4B4L1FR-S28R.

41. The formulation of any one of claims 1 to 40, wherein the formulation is in an aqueous carrier,

- 42. The formulation of claim 41, wherein the aqueous carrier is distilled water.
- 43. The formulation of any one of claims 1 to 42, wherein the formulation is sterile.
- 44. The formulation of any one of claims 1 to 43, wherein the formulation is homogenous.
- 45. The formulation of any one of claims 1 to 44, wherein the formulation has been prepared by a method that does not have a drying step.
- 46. The formulation of any one of claims 1 to 45, wherein the formulation has been prepared by a method that does not have a lyophilization step.
- 47. A pharmaceutical dosage form comprising the formulation of any one of claims 1 to 46, which dosage form is suitable for parenteral administration to a human and is in a suitable container.
- 48. The pharmaceutical unit dosage form of claim 47, wherein the formulation is suitable for subcutaneous, intravenous or intramuscular administration.
- 49. The pharmaceutical unit dosage form of any one of claims 1 to 46, which dosage form is suitable for aerosol administration to a human and is in a suitable container.
 - 50. A sealed container comprising the formulation of any one of claims 1 to 46.
- 51. A method of preventing, treating, or ameliorating one or more symptoms associated with a RSV infection in a subject, said method comprising administering a prophylactically or therapeutically effective amount of the formulation of any one of claims 1 to 46.
- 52. The method of claim 51, wherein the formulation is administered parenterally, intraweously, subcutaneously or intranasally.

53. A method of optimizing an antibody formulation for antibody fragmentation, said method comprising comparing by analytical ultracentrifugation (AUC) the fragmentation levels of a first antibody formulation produced according to a first protocol and a second antibody formulations produced according to a second protocol, which first and second antibody formulations comprise at least 100 mg/ml of a full length IgG₁ antibody comprising a heavy chain having the variable heavy (VH) domain of A4B4L1FR-S28R (SEQ ID NO:48) and the light chain comprising the variable light (VL) domain of A4B4L1FR-S28R (SEQ ID NO:11), wherein the first and second antibody formulations have been stored at 38-42° C, pH 6.0 for 1 month, wherein an antibody formulation is optimized for fragmentation levels if the fragmentation levels in the second antibody formulation are reduced relative to fragmentation levels in the first antibody formulation.

- 54. A method of optimizing an antibody formulation for antibody fragmentation, said method comprising comparing a first antibody formulation produced according to a first protocol and a second antibody formulation produced according to a second protocol, which first and second antibody formulations comprise at least 100 mg/ml of a full length IgG₁ antibody comprising a heavy chain having the variable heavy (VH) domain of A4B4L1FR-S28R (SEQ ID NO:48) and the light chain comprising the variable light (VL) domain of A4B4L1FR-S28R (SEQ ID NO:11), for the abundance of antibody type I and antibody type II fragments as determined by Liquid Chromatography Mass Spectrometry (LC-MS) analysis of samples of said stored antibody that have been deglycosylated, reduced and alkylated, wherein said first and second antibody formulations have been stored at 38-42° C, pH 6.0 for 1 month, wherein an antibody formulation is optimized for antibody fragmentation if the levels of antibody type I and antibody type II fragments in the second formulation are reduced relative to the levels of antibody type I and antibody type II fragments in the first antibody formulation.
 - 55. The method of claim 54, wherein said antibody type I fragments comprise one or more heavy chain C-terminal portions, which heavy chain C-terminal portion has a molecular weight of about 25.6 kD, about 25.7 kD, about 25.8 kD, about 26.0 kD, or about 26.1 kD and said antibody type II fragments comprise one or more N-terminal portions of said heavy chain, which heavy chain N-terminal portion has a molecular weight of about 24.4 kD, about 24.6 kD, about 24.7 kD, about 24.9 kD, or about 25.1 kD, wherein the molecular weight of said portion is determined by Liquid Chromatography Mass Spectrometry (LC-MS) analysis

of samples of said antibody formulation that have been deglycosylated, reduced and alkylated.


- 56. The method of any one of claim 53 to 55, wherein said second protocol comprises an additional chromatography purification step.
- 57. The method of claim 56, wherein said chromatography purification step utilizes a hydroxyapatite column.
- 58. The method of any one of claim 53 to 55, wherein said first protocol comprises an additional chromatography purification step.
- 59. The method of claim 58, wherein said chromatography purification step utilizes an rProtein A affinity column.
- 60. The method of any of claims 53 to 55, wherein said second antibody formulation comprises an antibody manufactured and purified by a process comprising (a) transformation of a murine myeloma cell line with a recombinant vector capable of directing transcription of mRNA encoding said antibody; (b) maintenance of transformed cells; (c) collection of conditioned media from cultures of transformed cells; (c) cation/anion exchange chromatography; (e) nanofiltration; and (f) hydroxyapatite chromatography.
- 61. The method of any of claims 53 to 55, wherein said second protocol comprises the alteration of a temperature in one or more steps of said first protocol.
- 62. An antibody comprising a Fab fragment, which immunospecifically binds to an RSV antigen, wherein the Tm of the Fab fragment is at least about 87 °C, and wherein said antibody is not any of palivizumab, AFFF, P12f2, P12f4, P11d4, Ale9, A12a6, A13c4, A17d4, A4B4, A8c7, 1X-493L1FR, H3-3F4, M3H9, Y10H6, DG, AFFF(1), 6H8, L1-7E5, L2-15B10, A13a11, A1h5, A4B4(1), A4B4L1FR-S28R (motavizumab), A4B4-F52S, A17d4(1), A3e2, A14a4, A16b4, A17b5, A17f5, and A17h4.
- 63. The antibody of claim 62, wherein said Fab is different from the Fab of palivizumab.

64. The antibody of claim 62, wherein the antibody comprises a VH domain that is different from the VH domain of palivizumab.


- 65. The antibody of claim 62, wherein the antibody comprises a VL domain that is different from the VL domain of palivizumab.
- 66. The antibody of claim 62, wherein the Tm of the Fab fragment is at least about 90 °C or at least about 93 °C.
- 67. The antibody of claim 62, wherein the pI of the antibody is between about 8.5 to 9.5or between about 9.0 to 9.5.
 - 68. The antibody of claim 62, wherein said RSV antigen is an F protein epitope.
- 69. The antibody of claim 62, wherein said RSV antigen comprises the F protein epitope NSELLSLINDMPITNDQKKLMSNN (SEQ ID NO:337).
- 70. The antibody of claim 62, wherein the antibody competitively inhibits the binding of antibody A4B4L1FR-S28R to said RSV antigen.
- 71. The antibody of claim 62, wherein the antibody comprises a VH domain of the antibody A4B4L1FR-S28R (SEQ ID NO:48).
- 72. The antibody of claim 62, wherein the antibody comprises a VL domain of the antibody A4B4L1FR-S28R (SEQ ID NO:11).
- --- J3. The antibody of claim 62, wherein said Fab is the Fab of antibody A4B4L1FR-S28R.
- 74. An antibody formulation comprising a full length IgG_1 antibody, which immunospecifically binds to an RSV antigen, said formulation having a viscosity of less than about 10.00 cP at any temperature in the range of 1 to 26 °C.
- 75. An antibody formulation comprising a full length IgG₁ antibody, which immunospecifically binds to an RSV antigen, said formulation having an aggregation rate of less than 15% per day at any temperature in the range of 38 to 42 °C.

76. An an antibody formulation comprising an antibody as claimed in any one of claims 62 to 73, wherein said formulation having a viscosity of less than 10.00 cP at any temperature in the range of 1 to 26 °C.

- 77. An an antibody formulation comprising an antibody as claimed in any one of claims 62 to 73, wherein said formulation having an aggregation rate of less than 15% per day at any temperature in the range of 38 to 42 °C.
- 78. A method of preventing, treating, or ameliorating one or more symptoms associated with a RSV infection in a subject, said method comprising administering a prophylactically or therapeutically effective amount of an antibody formulation comprising the antibody of any one of claims 62 to 73 or the antibody formulation of claims 74 to 77.
- 79. The method of claim 51 or 78, wherein RSV infection is an upper respiratory tract infection.
- 80. The method of claim 78, wherein the formulation is administered parenterally, intrawenously, subcutaneously or intranasally.
- 81. The method of claim 51 or 78, wherein said one or more symptoms are one or more of otitis media, asthma, and wheezing.



F19.1



F19. Z

DIOMTOSPET LSASVEDRYT ITCKCOLSVEYMH WYOOKPE 40

Kapkluy Dtsklas Gvpsr Frasgasatef Tltisslapd 80

DFATYYC FOGSGYPFT FGGGTKLEK

108

QVTLRESGPA LVKPTQTLTL TCTFSGFSLS TSGMSVG WIR

OPPGKALEWL A <u>DIWWDDKKDYNPGLKS</u> RLT ISKDT6KNOV 80

VLKVTNIMDPA DTATYYCAR <u>BMITNWYFDV</u> W <u>QAGITYTV68 120</u> CDR H3

<

F14.3

⋖

DIOMTORPST LEASVGDRVT ITCEASSSVGYMH WYGOKPG 40

Kapkiliy disklas gvpgr fgasgsgtef titisslapd bo

DFATYYC FOREGYPFT FGGG TKVEIK
CDR L3

168

m

OVTLRESGPA LYKPTOTLTL TCTF6GF6LS TSGMSVG WIR 40

OPPOKALEWI A DIWWDDKKDYNPSLKS RLT ISKDTSKNOV 80 CDR HZ VLKVTNUAPPA DTATYYCAR EMITNWYFDY WGOGTTVTV66 120

F19.4

: Nucleotide and Translated Amino Acid Sequence of the MEDI-524

1				•									•	1.5	
CM	GTC	: ACA	CTG	AGG	GAG	TCI	GGT	ccı	GCG	CTC	GIN	G AA	A CO	: ACA	
Q	v	T	L	R	E	5	G	P	A	L	V	K	P	T	MEDI-524
															Sypagis ^e
16														30	
CM) ACC	CIC	ACA	CTC) ACC				TCT	GGG	TT	P TC	CN) AGC	
Q	T	L	T	L	Ŧ	C	T	P	8	G	7	8	, Ł	\$	MEDI-524 Synagia
31					,									45	
														ANG	
I	3	<u> </u>	_ж	_8			W	I	R ,	Q	P	P	Œ	K	MEDI-524 Synagia
46										~~				GAC	•
ي ال															
	Ŀ		₩	4	^	W								H	nedr-524 Synagis
61 TAT	327	CCA	200	CTG	110	GAC	caa	CTC	XCA.	AFC	TCC	. 'ARG	GAT.	75 ACC	•
		2													NEDI-524
						\$									Synagis
76		λAC												90	. •
100	-														
3	K	×	Q	V	V	L	K	•	T	Я.		B	P	X	Synagis
91		GCC						~~~	~~	100	1.00		***	105	
					Y		GÇE A		D						VEDT_204
D	T	λ	T	Y	1	C	^		3	Д		Ŧ		.z. V	MEDI-524 Synagie
									•			*		*	PARMETE
106													•	120	
TÁC	TTC	GAT	GTC	TGG	GGC	CAG	GGG	ACC	ACG	GIC	ACC	GTG	AGC	TCA	
<u> </u>	7	D	¥	W	G	Q ·	G	T	T	v	T	V	2	\$	NEDI-524
			-			A									Synagis*


F19.5A

Nucleotide and Translated Amino Acid Sequence of the MEDI-524


													15	
atc I	Q Q	n Vla	ACC T	CAG Q	TCT S	CCT P	TCC S	ACC T	CTG L	TCT S	GCA A	TCT S	ota V	MEDI-524 Synagis
	AGA R	oic C	ACC T	ATC I	act T	TGC C	AGC	gee C	AGC g	TCG	coc 1	GTA V	30 GGT G	MEDI-524 Synagis
DTA W	CAC H	a Log	TAC Y	G CMG	CAG	raa K	CCC	GGG G	XXX K	gcc A	oct P	ang K	45 CTC L	MEDI-524 Synagis
ATC I	TAT Y	GAC D	ACT T	AGT S	XXX	CTG L	ccr	TCT S	ggg G	CIC V	cca P	TCA S	60 AGG R	MEDI-524 Synagis
agc S	GGC G	ngt S	GGA G	TCT S	GGG G	ACA T	E Gyy	TTC F	act T	CTC L	ACC T	atc I	75 AGC 8	MEDI-524 Synagis
CTG L	CAG Q	CCT P	GAC D	gat D	TTT P	GCA.	ACT T	TAT Y	TAC Y	TGC C	777	CAG O	90 90 90	MEDI-524 Synagis
GGG G	TAC Y				TTC F	GCA.	GGG G	996 G	ACC T	aag K	¥	gal E	105 ATA I	MEDI-524
<u>a</u>	<u>Y</u>	P	. F	.I	•	.	•	•	•	•	L	-	~	Synagis*
	GAC D ATG ATC I AGC S CTG GGG	GAC AGA D R ATG CAC M TAT I Y AGC GGC G CAG L Q GGG TAC	GAC AGA GTC ATG CAC TGG M R ATC TAT GAC TAT GAC ACC GGC AGT ACC GAG CCT CTG CAG CCT CTG CAG CCT CTG CAG CCT CTG CAG CCT	I Q M T GAC AGA GTC ACC D R V T ATG CAC TGG TAC M B W Y ATC TAT GAC ACT I Y D T AGC GGC AGT GGA G G G G GAC G G G G G GAC L Q P GAC GGG TAC CCA TTC	GAC AGA GTC ACC ATC D R V T T I ATG CAC TGG TAC CAG M R V Y Q ATC TAT GAC ACT AGT I Y D T S AGC GGC AGT GGA TCT S G S G AGT GAC GAT L Q P GAC GAT D GGG TAC CCA TTC ACG	GAC AGA GTC ACC ATC ACT D R V T T ACC ATC ACT T T ATG CAC TGG TAC CAG CAG M R V T T S ACT T T ATG CAC TGG TAC GAG TCT S GGG AGT GGA TCT C CAG CCT GAC GAT TTT L Q P D C ACG TTC GGG TAC CCA TTC ACG TTC	GAC AGA GTC ACC ATC ACT TGC ATG CAC TGG TAC CAG CAG AAA M H GAC ACT AGT AAA CTG ATC TAT GAC ACT AGT AAA CTG AGC GGC AGT GGA TCT GGG ACA GGG TAC CCA TTC ACG TTC GGA GGG TAC CCA TTC ACG TTC GGA	GAC AGA GTC ACC ATC ACT TGC AGC D R V TC TC ACC ATC ACT TGC AGC E ATG CAC TGG TAC CAG CAG AAA CCC M H Y Q Q Q K PA ATC TAT GAC ACT AGT AAA CTG GCT I Y D T S R L A AGC GGC AGT GGA TCT GGG ACA GAA S G G G ACT ACT TTT GCA ACT L Q P D D GAT TTT GCA ACT T	GAC AGA GTC ACC ATC ACT TGC AGC GCC ATG CAC TGG TAC CAG CAG AAA CCC GGG ATG CAC TGG TAC CAG CAG AAA CCC GGG ATC TAT GAC ACT AGT AAA CTG GCT TCT Y AGC GGC AGT GGA TCT GGG ACA GAA TTC S G G T TTT CTG CAG CCT GAC GAT TTT GCA ACT TAT L GGG TAC CCA TTC ACG TTC GGA GGG GGG	GAC AGA GTC ACC ATC ACT TOC AGC GCC AGC D R V T I T T C R ACC GCC AGC ATG CAC TGG TAC CAG CAG ANA CCC GGG AAA N H W Y Q Q R R P G R ATC TAT GAC ACT AGT AAA CTG GCT TCT GGG AGC GGC AGT G GG ACA GAA TTC ACT T G CAG CCT GAC GAT TTT GCA ACT TAT TAC L Q P D G GAT TTC GCA GGG GGG ACC GGG TAC CCA TTC ACG TTC GGA GGG GGG ACC	GAC AGA GTC ACC ATC ACT TOC AGC GCC AGC TCG B A A G C Q L ATG CAC TGG TAC CAG CAG AAA CCC GGG AAA GCC M H W Y Q Q K P G G K A ATC TAT GAC ACT AGT AAA CTG GCT TCT GGG GTC I Y D T S K L A S G G C AGC GGC AGT GGA TCT GGG ACA GAA TTC ACT CTC B G G G G G G G G ACA GAA TTC ACT CTC CTG CAG CCT GAC GAT TTT GCA ACT TAT TAC TGC GGG TAC CCA TTC ACG TTC GGA GGG GGG ACC AAG	GAC AGA GTC ACC ATC ACT TGC AGC GCC AGC TGG CGC ATG CAC TGG TAC CAG CAG AAA CCC GGG AAA GCC CCT ATC TAT GAC ACT AGT AAA CTG GCT TCT GGG GTC CCA ACC GGC AGT GGA TCT GGG ACA GAA TTC ACT CTC ACC CTG CAG CCT GAC GAT TTT GCA ACT TAT TAC TGC TTT CTG CAG CCT GAC GAT TTT GCA ACT TAT TAC TGC TTT GGG TAC CCA TTC ACG TTC GGA GGG GGG ACC AAG GTC GGG TAC CCA TTC ACG TTC GGA GGG GGG ACC AAG GTC GGG TAC CCA TTC ACG TTC GGA GGG GGG ACC AAG GTC	GAC AGA GTC ACC ATC ACT TGC AGC GCC AGC TGG CGC GTA ATG CAC TGG TAC CAG CAG RAA CCC GGG AAA GCC CCT AMG ATG TAT GAC ACT AGT AAA CTG GCT TCT GGG GTC CCA TCA ATC TAT GAC ACT AGT AAA CTG GCT TCT GGG GTC CCA TCA ACC GGC AGT GGA TCT GGG ACA GAA TTC ACT CTG ACC ATC B CAG CCT GAC GAT TTT GCA ACT TAT TAC TGC TTT CAG CTG CAG CCT GAC GAT TTT GCA ACT TAT TAC TGC TTT CAG GGG TAC CCA TTC ACG TTC GGA GGG GGG ACC AAG GTC GAA GCG TAC CCA TTC ACG TTC GGA GGG GGG ACC AAG GTC GAA GCG TAC CCA TTC ACG TTC GGA GGG GGG ACC AAG GTC GAA GCG TAC CCA TTC ACG TTC GGA GGG GGG ACC AAG GTC GAA GCG TAC CCA TTC ACG TTC GGA GGG GGG ACC AAG GTC GAA GCG TAC CCA TTC ACG TTC GGA GGG GGG ACC AAG GTC GAA GCG TAC CCA TTC ACG TTC GGA GGG GGG ACC AAG GTC GAA	ATC CAG ATG ACC CAG TCT CCT TCC ACC CTG TCT GCA TCT GTA GAC AGA GTC ACC ATC ACT TGC AGC GCC AGC TCG CGC GTA GGT ATG CAC TGG TAC CAG CAG AAA CCC GGG AAA GCC CCT AAG CTC ATC TAT GAC ACT AGT AAA CTG GCT TCT GGG GTC CCA TCA AGG ATC TAT GAC ACT AGT AAA CTG GCT TCT GGG GTC CCA TCA AGG ACC GGC AGT GGA TCT GGG ACA GAA TTC ACT CTC TCT AGC S G G GC AGT GAC GAT TTT GCA ACT TAT TAC TCC TTT CAG GGG CTG CAG CCT GAC GAT TTT GCA ACT TAT TAC TCC TTT CAG GGG GGG TAC CCA TTC ACG TTC GGA GGG GGG ACC AAG GTC GAA ATTA GGG TAC CCA TTC ACG TTC GGA GGG GGG ACC AAG GTC GAA ATTA G Y P F T F G GGA GGG GGG ACC AAG GTC GAA ATTA T K Y E I

x MEDI-524 Synagis


F19.5B




F19.6




F19.7




F19.8







F19.9




F19. 10



F19.11




1 19. 17




F19, 13

Fragmentation Pattern of A4B4L1FR-S28R


3




F14.14




F19.15





F19.16




















F16. 7

PCT/US2006/024717

Sheet 23 of 23

10271-170-228



SEQUENCE LISTING

```
<110> MedImmune, Inc.
<120> Antibody Formulations Having Optimized Aggregation
      and Fragmentation Profiles
<130> 10271-170-228
<140> To be assinged
<141> On even date herewith
<150> 60/693,603
<151> 2005-06-23
<150> 60/699,614
<151> 2005-07-15
<160> 1496
<170> PatentIn version 3.1
<210> 1
<211> 7
<212> PRT
<213> Murine
<400> 1
Thr Ser Gly Met Ser Val Gly
<210> 2
<211> 16
<212> PRT
<213> Murine
<400> 2
Asp Ile Trp Trp Asp Asp Lys Lys Asp Tyr Asn Pro Ser Leu Lys Ser
                           10
<210> 3
<211> 10
<212> PRT
<213> Murine
Ser Met Ile Thr Asn Trp Tyr Phe Asp Val
<210> 4
<211> 10
<212> PRT
<213> Murine
<400> 4
Lys Cys Gln Leu Ser Val Gly Tyr Met His
         5
<210> 5
<211> 7
<212> PRT
<213> Murine
```

```
<400> 5
Asp Thr Ser Lys Leu Ala Ser
<210> 6
<211> 9
<212> PRT
<213> Murine
<400> 6
Phe Gln Gly Ser Gly Tyr Pro Phe Thr
<210> 7
<211> 120
<212> PRT
<213> Artificial Sequence
<220>
<223> Humanized antibody - VH Domain
Gln Val Thr Leu Arg Glu Ser Gly Pro Ala Leu Val Lys Pro Thr Gln
Thr Leu Thr Leu Thr Cys Thr Phe Ser Gly Phe Ser Leu Ser Thr Ser
                               25
Gly Met Ser Val Gly Trp Ile Arg Gln Pro Pro Gly Lys Ala Leu Glu
                           40
Trp Leu Ala Asp Ile Trp Trp Asp Asp Lys Lys Asp Tyr Asn Pro Ser
                       55
Leu Lys Ser Arg Leu Thr Ile Ser Lys Asp Thr Ser Lys Asn Gln Val
                                       75
Val Leu Lys Val Thr Asn Met Asp Pro Ala Asp Thr Ala Thr Tyr Tyr
                                90
Cys Ala Arg Ser Met Ile Thr Asn Trp Tyr Phe Asp Val Trp Gly Ala
           100
Gly Thr Thr Val Thr Val Ser Ser
      115
<210> 8
<211> 106
<212> PRT
<213> Artificial Sequence
<220>
<221> misc_feature
<223> Humanized antibody - VL Domain
<400> 8
Asp Ile Gln Met Thr Gln Ser Pro Ser Thr Leu Ser Ala Ser Val Gly
Asp Arg Val Thr Ile Thr Cys Lys Cys Gln Leu Ser Val Gly Tyr Met
His Trp Tyr Gln Gln Lys Pro Gly Lys Ala Pro Lys Leu Leu Ile Tyr
Asp Thr Ser Lys Leu Ala Ser Gly Val Pro Ser Arg Phe Ser Gly Ser
Gly Ser Gly Thr Glu Phe Thr Leu Thr Ile Ser Ser Leu Gln Pro Asp
Asp Phe Ala Thr Tyr Tyr Cys Phe Gln Gly Ser Gly Tyr Pro Phe Thr
```

```
85
                                                       95
Phe Gly Gly Thr Lys Leu Glu Ile Lys
            100
<210> 9
<211> 120
<212> PRT
<213> Artificial Sequence
<223> Humanized antibody - VH Domain
Gln Val Thr Leu Arg Glu Ser Gly Pro Ala Leu Val Lys Pro Thr Gln
Thr Leu Thr Leu Thr Cys Thr Phe Ser Gly Phe Ser Leu Ser Thr Ala
Gly Met Ser Val Gly Trp Ile Arg Gln Pro Pro Gly Lys Ala Leu Glu
Trp Leu Ala Asp Ile Trp Trp Asp Asp Lys Lys Asp Tyr Asn Pro Ser
                       55
Leu Lys Ser Arg Leu Thr Ile Ser Lys Asp Thr Ser Lys Asn Gln Val
                    70
Val Leu Lys Val Thr Asn Met Asp Pro Ala Asp Thr Ala Thr Tyr Tyr
Cys Ala Arg Ser Met Ile Thr Asn Phe Tyr Phe Asp Val Trp Gly Gln
            100
                            105
Gly Thr Thr Val Thr Val Ser Ser
      115
<210> 10
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 10
Thr Ala Gly Met Ser Val Gly
<210> 11
<211> 106
<212> PRT
<213> Artificial Sequence
<220>
<223> Humanized antibody - VL Domain
Asp Ile Gln Met Thr Gln Ser Pro Ser Thr Leu Ser Ala Ser Val Gly
                                   10
Asp Arg Val Thr Ile Thr Cys Ser Ala Ser Ser Arg Val Gly Tyr Met
                               25
His Trp Tyr Gln Gln Lys Pro Gly Lys Ala Pro Lys Leu Leu Ile Tyr
                          40
Asp Thr Ser Lys Leu Ala Ser Gly Val Pro Ser Arg Phe Ser Gly Ser
```

```
Gly Ser Gly Thr Glu Phe Thr Leu Thr Ile Ser Ser Leu Gln Pro Asp
                   70
                            75
Asp Phe Ala Thr Tyr Tyr Cys Phe Gln Gly Ser Gly Tyr Pro Phe Thr
               8.5
                                  90
Phe Gly Gly Thr Lys Val Glu Ile Lys
<210> 12
<211> 10
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
     antibody and further modified by amino acid
     substitutions
<400> 12
Ser Met Ile Thr Asn Phe Tyr Phe Asp Val
<210> 13
<211> 106
<212> PRT
<213> Artificial Sequence
<220>
<223> Humanized antibody - VL Domain
<400> 13
Asp Ile Gln Met Thr Gln Ser Pro Ser Thr Leu Ser Ala Ser Val Gly
                                  10
Asp Arg Val Thr Ile Thr Cys Ser Ala Ser Ser Ser Val Gly Tyr Met
                               25
His Trp Tyr Gln Gln Lys Pro Gly Lys Ala Pro Lys Leu Leu Ile Tyr
                           40
Asp Thr Phe Lys Leu Ala Ser Gly Val Pro Ser Arg Phe Ser Gly Ser
Gly Ser Gly Thr Glu Phe Thr Leu Thr Ile Ser Ser Leu Gln Pro Asp
                                      75
Asp Phe Ala Thr Tyr Tyr Cys Phe Gln Phe Ser Gly Tyr Pro Phe Thr
                                  90
Phe Gly Gly Thr Lys Val Glu Ile Lys
<210> 14
<211> 10
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
     antibody and further modified by amino acid
     substitutions
<400> 14
Ser Ala Ser Ser Ser Val Gly Tyr Met His
<210> 15
<211> 7
```

```
<212> PRT
<213> Artificial Sequence
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
     substitutions
<400> 15
Asp Thr Phe Lys Leu Ala Ser
<210> 16
<211> 9
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
     substitutions
<400> 16
Phe Gln Phe Ser Gly Tyr Pro Phe Thr
<210> 17
<211> 120
<212> PRT
<213> Artificial Sequence
<220>
<223> Humanized antibody - VH Domain
<400> 17
Gln Val Thr Leu Arg Glu Ser Gly Pro Ala Leu Val Lys Pro Thr Gln
                                   10
Thr Leu Thr Leu Thr Cys Thr Phe Ser Gly Phe Ser Leu Ser Thr Pro
                                25
Gly Met Ser Val Gly Trp Ile Arg Gln Pro Pro Gly Lys Ala Leu Glu
                            40
Trp Leu Ala Asp Ile Trp Trp Asp Asp Lys Lys His Tyr Asn Pro Ser
                       55
Leu Lys Asp Arg Leu Thr Ile Ser Lys Asp Thr Ser Lys Asn Gln Val
                   70
                                       75
Val Leu Lys Val Thr Asn Met Asp Pro Ala Asp Thr Ala Thr Tyr Tyr
               85
                                   90
Cys Ala Arg Asp Met Ile Phe Asn Phe Tyr Phe Asp Val Trp Gly Gln
          100
Gly Thr Thr Val Thr Val Ser Ser
       115
<210> 18
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
     antibody and further modified by amino acid
     substitutions
```

```
<400> 18
Thr Pro Gly Met Ser Val Gly
   5
<210> 19
<211> 16
<212> PRT
<213> Artificial Sequence
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 19
Asp Ile Trp Trp Asp Asp Lys Lys His Tyr Asn Pro Ser Leu Lys Asp
    5
                                   10
<210> 20
<211> 10
<212> PRT
<213> Artificial Sequence
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 20
Asp Met Ile Phe Asn Phe Tyr Phe Asp Val
<210> 21
<211> 106
<212> PRT
<213> Artificial Sequence
<220>
<223> Humanized antibody - VL Domain
Asp Ile Gln Met Thr Gln Ser Pro Ser Thr Leu Ser Ala Ser Val Gly
                                    10
Asp Arg Val Thr Ile Thr Cys Ser Leu Ser Ser Arg Val Gly Tyr Met
                                2.5
His Trp Tyr Gln Gln Lys Pro Gly Lys Ala Pro Lys Leu Leu Ile Tyr
                            40
Asp Thr Phe Tyr Leu Ser Ser Gly Val Pro Ser Arg Phe Ser Gly Ser
                        55
                                            60
Gly Ser Gly Thr Glu Phe Thr Leu Thr Ile Ser Ser Leu Gln Pro Asp
                    70
                                       75
Asp Phe Ala Thr Tyr Tyr Cys Phe Gln Gly Ser Gly Tyr Pro Phe Thr
                85
Phe Gly Gly Gly Thr Lys Val Glu Ile Lys
<210> 22
<211> 10
<212> PRT
```

```
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 22
Ser Leu Ser Ser Arg Val Gly Tyr Met His
                5
<210> 23
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 23
Asp Thr Phe Tyr Leu Ser Ser
<210> 24
<211> 120
<212> PRT
<213> Artificial Sequence
<220>
<223> Humanized antibody - VH Domain
<400> 24
Gln Val Thr Leu Arg Glu Ser Gly Pro Ala Leu Val Lys Pro Thr Gln
                                     10
Thr Leu Thr Leu Thr Cys Thr Phe Ser Gly Phe Ser Leu Ser Thr Pro
                                25
Gly Met Ser Val Gly Trp Ile Arg Gln Pro Pro Gly Lys Ala Leu Glu
                                                45
Trp Leu Ala Asp Ile Trp Trp Asp Gly Lys Lys His Tyr Asn Pro Ser
                        55
Leu Lys Asp Arg Leu Thr Ile Ser Lys Asp Thr Ser Lys Asn Gln Val
                                         75
Val Leu Lys Val Thr Asn Met Asp Pro Ala Asp Thr Ala Thr Tyr Tyr
                                    90
                                                        95
Cys Ala Arg Asp Met Ile Phe Asn Phe Tyr Phe Asp Val Trp Gly Gln
            100
                               105
Gly Thr Thr Val Thr Val Ser Ser
        115
<210> 25
<211> 16
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
```

```
<400> 25
Asp Ile Trp Trp Asp Gly Lys Lys His Tyr Asn Pro Ser Leu Lys Asp
<210> 26
<211> 106
<212> PRT
<213> Artificial Sequence
<220>
<223> Humanized antibody - VL Domain
<400> 26
Asp Ile Gln Met Thr Gln Ser Pro Ser Thr Leu Ser Ala Ser Val Gly
                                   1.0
Asp Arg Val Thr Ile Thr Cys Ser Leu Ser Ser Arg Val Gly Tyr Met
                               25
His Trp Tyr Gln Gln Lys Pro Gly Lys Ala Pro Lys Leu Leu Ile Tyr
                           40
Asp Thr Arg Gly Leu Pro Ser Gly Val Pro Ser Arg Phe Ser Gly Ser
                       55
Gly Ser Gly Thr Glu Phe Thr Leu Thr Ile Ser Ser Leu Gln Pro Asp
                   70
                                      75
Asp Phe Ala Thr Tyr Tyr Cys Phe Gln Gly Ser Gly Tyr Pro Phe Thr
                                   90
Phe Gly Gly Gly Thr Lys Val Glu Ile Lys
<210> 27
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 27
Asp Thr Arg Gly Leu Pro Ser
<210> 28
<211> 120
<212> PRT
<213> Artificial Sequence
<220>
<223> Humanized antibody - VH Domain
<400> 28
Gln Val Thr Leu Arg Glu Ser Gly Pro Ala Leu Val Lys Pro Thr Gln
Thr Leu Thr Leu Thr Cys Thr Phe Ser Gly Phe Ser Leu Ser Thr Pro
                                2.5
Gly Met Ser Val Gly Trp Ile Arg Gln Pro Pro Gly Lys Ala Leu Glu
                            40
Trp Leu Ala Asp Ile Trp Trp Asp Gly Lys Lys His Tyr Asn Pro Ser
                                            60
Leu Lys Asp Arg Leu Thr Ile Ser Lys Asp Thr Ser Lys Asn Gln Val
```

```
65
                                        75
Val Leu Lys Val Thr Asn Met Asp Pro Ala Asp Thr Ala Thr Tyr Tyr
                                    90
Cys Ala Arg Asp Met Ile Phe Asn Trp Tyr Phe Asp Val Trp Gly Gln
            100
                                105
Gly Thr Thr Val Thr Val Ser Ser
        115
<210> 29
<211> 10
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 29
Asp Met Ile Phe Asn Trp Tyr Phe Asp Val
<210> 30
<211> 106
<212> PRT
<213> Artificial Sequence
<220>
<223> Humanized antibody - VL Domain
<400> 30
Asp Ile Gln Met Thr Gln Ser Pro Ser Thr Leu Ser Ala Ser Val Gly
                                    10
Asp Arg Val Thr Ile Thr Cys Ser Pro Ser Ser Arg Val Gly Tyr Met
His Trp Tyr Gln Gln Lys Pro Gly Lys Ala Pro Lys Leu Leu Ile Tyr
Asp Thr Met Arg Leu Ala Ser Gly Val Pro Ser Arg Phe Ser Gly Ser
Gly Ser Gly Thr Glu Phe Thr Leu Thr Ile Ser Ser Leu Gln Pro Asp
                                        75
Asp Phe Ala Thr Tyr Tyr Cys Phe Gln Gly Ser Gly Tyr Pro Phe Thr
Phe Gly Gly Gly Thr Lys Val Glu Ile Lys
            100
<210> 31
<211> 10
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
     substitutions
Ser Pro Ser Ser Arg Val Gly Tyr Met His
<210> 32
```

```
<211> 7
<212> PRT
<213> Artificial Sequence
<223> Amino acid sequence derived from Murine monoclonal
     antibody and further modified by amino acid
     substitutions
<400> 32
Asp Thr Met Arg Leu Ala Ser
1 5
<210> 33
<211> 120
<212> PRT
<213> Artificial Sequence
<220>
<223> Humanized antibody - VH Domain
<400> 33
Gln Val Thr Leu Arg Glu Ser Gly Pro Ala Leu Val Lys Pro Thr Gln
                                   1.0
Thr Leu Thr Leu Thr Cys Thr Phe Ser Gly Phe Ser Leu Ser Thr Ala
                               2.5
Gly Met Ser Val Gly Trp Ile Arg Gln Pro Pro Gly Lys Ala Leu Glu
                           40
Trp Leu Ala Asp Ile Trp Trp Asp Gly Lys Lys His Tyr Asn Pro Ser
                       55
Leu Lys Asp Arg Leu Thr Ile Ser Lys Asp Thr Ser Lys Asn Gln Val
                                       75
                   70
Val Leu Lys Val Thr Asn Met Asp Pro Ala Asp Thr Ala Thr Tyr Tyr
                                  90
               85
Cys Ala Arg Asp Met Ile Phe Asn Trp Tyr Phe Asp Val Trp Gly Gln
           100 105
Gly Thr Thr Val Thr Val Ser Ser
       115
<210> 34
<211> 106
<212> PRT
<213> Artificial Sequence
<220>
<223> Humanized antibody - VL Domain
<400> 34
Asp Ile Gln Met Thr Gln Ser Pro Ser Thr Leu Ser Ala Ser Val Gly
                                   10
Asp Arg Val Thr Ile Thr Cys Ser Leu Ser Ser Arg Val Gly Tyr Met
                                25
His Trp Tyr Gln Gln Lys Pro Gly Lys Ala Pro Lys Leu Leu Ile Tyr
                            40
Asp Thr Phe Lys Leu Ser Ser Gly Val Pro Ser Arg Phe Ser Gly Ser
                        55
Gly Ser Gly Thr Glu Phe Thr Leu Thr Ile Ser Ser Leu Gln Pro Asp
                                       75
Asp Phe Ala Thr Tyr Tyr Cys Phe Gln Gly Ser Gly Tyr Pro Phe Thr
                85
Phe Gly Gly Gly Thr Lys Val Glu Ile Lys
```

105 100 <210> 35 <211> 7 <212> PRT <213> Artificial Sequence <220> <223> Amino acid sequence derived from Murine monoclonal antibody and further modified by amino acid substitutions <400> 35 Asp Thr Phe Lys Leu Ser Ser <210> 36 <211> 120 <212> PRT <213> Artificial Sequence <220> <223> Humanized antibody - VH Domain <400> 36 Gln Val Thr Leu Arg Glu Ser Gly Pro Ala Leu Val Lys Pro Thr Gln Thr Leu Thr Leu Thr Cys Thr Phe Ser Gly Phe Ser Leu Ser Thr Ala 25 Gly Met Ser Val Gly Trp Ile Arg Gln Pro Pro Gly Lys Ala Leu Glu 40 Trp Leu Ala Asp Ile Trp Trp Asp Gly Lys Lys Asp Tyr Asn Pro Ser 55 Leu Lys Asp Arg Leu Thr Ile Ser Lys Asp Thr Ser Lys Asn Gln Val 70 Val Leu Lys Val Thr Asn Met Asp Pro Ala Asp Thr Ala Thr Tyr Tyr 90 Cys Ala Arg Asp Met Ile Phe Asn Phe Tyr Phe Asp Val Trp Gly Gln 100 105 Gly Thr Thr Val Thr Val Ser Ser 115 <210> 37 <211> 16 <212> PRT <213> Artificial Sequence <220> <223> Amino acid sequence derived from Murine monoclonal antibody and further modified by amino acid substitutions Asp Ile Trp Trp Asp Gly Lys Lys Asp Tyr Asn Pro Ser Leu Lys Asp 10 <210> 38 <211> 106 <212> PRT <213> Artificial Sequence

```
<220>
<223> Humanized antibody - VL Domain
<400> 38
Asp Ile Gln Met Thr Gln Ser Pro Ser Thr Leu Ser Ala Ser Val Gly
                               10
Asp Arg Val Thr Ile Thr Cys Ser Ala Ser Ser Arg Val Gly Tyr Met
                               25
His Trp Tyr Gln Gln Lys Pro Gly Lys Ala Pro Lys Leu Leu Ile Tyr
Asp Thr Phe Lys Leu Ser Ser Gly Val Pro Ser Arg Phe Ser Gly Ser
                       55
Gly Ser Gly Thr Glu Phe Thr Leu Thr Ile Ser Ser Leu Gln Pro Asp
                                    75
                   70
Asp Phe Ala Thr Tyr Tyr Cys Phe Gln Gly Ser Gly Tyr Pro Phe Thr
            85
Phe Gly Gly Thr Lys Val Glu Ile Lys
           100
<210> 39
<211> 10
<212> PRT
<213> Artificial Sequence
<223> Amino acid sequence derived from Murine monoclonal
     antibody and further modified by amino acid
      substitutions
<400> 39
Ser Ala Ser Ser Arg Val Gly Tyr Met His
      5
<210> 40
<211> 120
<211> 120
<212> PRT
<213> Artificial Sequence
<223> Humanized antibody - VH Domain
Gln Val Thr Leu Arg Glu Ser Gly Pro Ala Leu Val Lys Pro Thr Gln
                                   10
Thr Leu Thr Leu Thr Cys Thr Phe Ser Gly Phe Ser Leu Ser Thr Ala
                                25
Gly Met Ser Val Gly Trp Ile Arg Gln Pro Pro Gly Lys Ala Leu Glu
                            40
Trp Leu Ala Asp Ile Trp Trp Asp Gly Lys Lys Ser Tyr Asn Pro Ser
                        55
Leu Lys Asp Arg Leu Thr Ile Ser Lys Asp Thr Ser Lys Asn Gln Val
                    70
                                        75
Val Leu Lys Val Thr Asn Met Asp Pro Ala Asp Thr Ala Thr Tyr Tyr
                                   90
Cys Ala Arg Asp Met Ile Phe Asn Phe Tyr Phe Asp Val Trp Gly Gln
           100 105
 Gly Thr Thr Val Thr Val Ser Ser
       115
 <210> 41
 <211> 16
```

```
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 41
Asp Ile Trp Trp Asp Gly Lys Lys Ser Tyr Asn Pro Ser Leu Lys Asp
<210> 42
<211> 106
<212> PRT
<213> Artificial Sequence
<220>
<223> Humanized antibody - VL Domain
<400> 42
Asp Ile Gln Met Thr Gln Ser Pro Ser Thr Leu Ser Ala Ser Val Gly
                                    10
Asp Arg Val Thr Ile Thr Cys Ser Leu Ser Ser Arg Val Gly Tyr Met
                                25
His Trp Tyr Gln Gln Lys Pro Gly Lys Ala Pro Lys Leu Leu Ile Tyr
                            40
Asp Thr Met Tyr Gln Ser Ser Gly Val Pro Ser Arg Phe Ser Gly Ser
                        55
Gly Ser Gly Thr Glu Phe Thr Leu Thr Ile Ser Ser Leu Gln Pro Asp
                    70
                                       75
Asp Phe Ala Thr Tyr Tyr Cys Phe Gln Gly Ser Gly Tyr Pro Phe Thr
               85
Phe Gly Gly Thr Lys Val Glu Ile Lys
            100
<210> 43
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 43
Asp Thr Met Tyr Gln Ser Ser
<210> 44
<211> 120
<212> PRT
<213> Artificial Sequence
<220>
<223> Humanized antibody - VH Domain
Gln Val Thr Leu Arg Glu Ser Gly Pro Ala Leu Val Lys Pro Thr Gln
```

13

```
Thr Leu Thr Leu Thr Cys Thr Phe Ser Gly Phe Ser Leu Ser Thr Ala
                               25
           20
Gly Met Ser Val Gly Trp Ile Arg Gln Pro Pro Gly Lys Ala Leu Glu
                                              45
                           40
Trp Leu Ala Asp Ile Trp Trp Asp Asp Lys Lys Ser Tyr Asn Pro Ser
                       55
Leu Lys Asp Arg Leu Thr Ile Ser Lys Asp Thr Ser Lys Asn Gln Val
                                      75
               70
Val Leu Lys Val Thr Asn Met Asp Pro Ala Asp Thr Ala Thr Tyr Tyr
                                  90
Cys Ala Arg Asp Met Ile Phe Asn Phe Tyr Phe Asp Val Trp Gly Gln
                   105
Gly Thr Thr Val Thr Val Ser Ser
<210> 45
<211> 16
<212> PRT
<213> Artificial Sequence
<223> Amino acid sequence derived from Murine monoclonal
     antibody and further modified by amino acid
      substitutions
<400> 45
Asp Ile Trp Trp Asp Asp Lys Lys Ser Tyr Asn Pro Ser Leu Lys Asp
                           10
<210> 46
<211> 106
<212> PRT
<213> Artificial Sequence
<223> Humanized antibody - VL Domain
<400> 46
Asp Ile Gln Met Thr Gln Ser Pro Ser Thr Leu Ser Ala Ser Val Gly
                                  10
Asp Arg Val Thr Ile Thr Cys Leu Pro Ser Ser Arg Val Gly Tyr Met
His Trp Tyr Gln Gln Lys Pro Gly Lys Ala Pro Lys Leu Leu Ile Tyr
Asp Thr Met Tyr Gln Ser Ser Gly Val Pro Ser Arg Phe Ser Gly Ser
                        55
Gly Ser Gly Thr Glu Phe Thr Leu Thr Ile Ser Ser Leu Gln Pro Asp
                                      75
                    70
Asp Phe Ala Thr Tyr Tyr Cys Phe Gln Gly Ser Gly Tyr Pro Phe Thr
               85
Phe Gly Gly Thr Lys Val Glu Ile Lys
           100
<210> 47
<211> 10
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
```

substitutions

```
<400> 47
Leu Pro Ser Ser Arg Val Gly Tyr Met His
<210> 48
<211> 120
<212> PRT
<213> Artificial Sequence
<220>
<223> Humanized antibody - VH Domain
<400> 48
Gln Val Thr Leu Arg Glu Ser Gly Pro Ala Leu Val Lys Pro Thr Gln
Thr Leu Thr Leu Thr Cys Thr Phe Ser Gly Phe Ser Leu Ser Thr Ala
                              25
Gly Met Ser Val Gly Trp Ile Arg Gln Pro Pro Gly Lys Ala Leu Glu
                          40
Trp Leu Ala Asp Ile Trp Trp Asp Asp Lys Lys His Tyr Asn Pro Ser
                       55
Leu Lys Asp Arg Leu Thr Ile Ser Lys Asp Thr Ser Lys Asn Gln Val
                                       75
Val Leu Lys Val Thr Asn Met Asp Pro Ala Asp Thr Ala Thr Tyr Tyr
                                  90
Cys Ala Arg Asp Met Ile Phe Asn Phe Tyr Phe Asp Val Trp Gly Gln
Gly Thr Thr Val Thr Val Ser Ser
<210> 49
<211> 106
<212> PRT
<213> Artificial Sequence
<220>
<223> Humanized antibody - VL Domain
<400> 49
Asp Ile Gln Met Thr Gln Ser Pro Ser Thr Leu Ser Ala Ser Val Gly
Asp Arg Val Thr Ile Thr Cys Ser Ala Ser Ser Arg Val Gly Tyr Met
His Trp Tyr Gln Gln Lys Pro Gly Lys Ala Pro Lys Leu Leu Ile Tyr
Asp Thr Phe Phe Leu Asp Ser Gly Val Pro Ser Arg Phe Ser Gly Ser
                        55
Gly Ser Gly Thr Glu Phe Thr Leu Thr Ile Ser Ser Leu Gln Pro Asp
Asp Phe Ala Thr Tyr Tyr Cys Phe Gln Gly Ser Gly Tyr Pro Phe Thr
               85
Phe Gly Gly Thr Lys Val Glu Ile Lys
<210> 50
<211> 7
<212> PRT
<213> Artificial Sequence
```

```
<223> Amino acid sequence derived from Murine monoclonal
     antibody and further modified by amino acid
     substitutions
<400> 50
Asp Thr Phe Phe Leu Asp Ser
<210> 51
<211> 120
<212> PRT
<213> Artificial Sequence
<223> Humanized antibody - VH Domain
<400> 51
Gln Val Thr Leu Arg Glu Ser Gly Pro Ala Leu Val Lys Pro Thr Gln
                                   1.0
Thr Leu Thr Leu Thr Cys Thr Phe Ser Gly Phe Ser Leu Ser Thr Ala
Gly Met Ser Val Gly Trp Ile Arg Gln Pro Pro Gly Lys Ala Leu Glu
Trp Leu Ala Asp Ile Trp Trp Asp Asp Lys Lys Ser Tyr Asn Pro Ser
                       55
Leu Lys Asp Arg Leu Thr Ile Ser Lys Asp Thr Ser Lys Asn Gln Val
                                       75
                   70
Val Leu Lys Val Thr Asn Met Asp Pro Ala Asp Thr Ala Thr Tyr Tyr
                                  90
Cys Ala Arg Asp Met Ile Phe Asn Trp Tyr Phe Asp Val Trp Gly Gln
          100 105
Gly Thr Thr Val Thr Val Ser Ser
        115
<210> 52
<211> 106
<212> PRT
<213> Artificial Sequence
<223> Humanized antibody - VL Domain
<400> 52
Asp Ile Gln Met Thr Gln Ser Pro Ser Thr Leu Ser Ala Ser Val Gly
                                   10
Asp Arg Val Thr Ile Thr Cys Ser Pro Ser Ser Arg Val Gly Tyr Met
                                25
His Trp Tyr Gln Gln Lys Pro Gly Lys Ala Pro Lys Leu Leu Ile Tyr
                           40
Asp Thr Arg Tyr Gln Ser Ser Gly Val Pro Ser Arg Phe Ser Gly Ser
Gly Ser Gly Thr Glu Phe Thr Leu Thr Ile Ser Ser Leu Gln Pro Asp
                                       75
Asp Phe Ala Thr Tyr Tyr Cys Phe Gln Gly Ser Gly Tyr Pro Phe Thr
               85
Phe Gly Gly Gly Thr Lys Val Glu Ile Lys
<210> 53
<211> 7
```

```
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 53
Asp Thr Arg Tyr Gln Ser Ser
<210> 54
<211> 106
<212> PRT
<213> Artificial Sequence
<220>
<223> Humanized antibody - VL Domain
<400> 54
Asp Ile Gln Met Thr Gln Ser Pro Ser Thr Leu Ser Ala Ser Val Gly
                                    10
Asp Arg Val Thr Ile Thr Cys Ser Ala Ser Ser Ser Val Gly Tyr Met
                               25
His Trp Tyr Gln Gln Lys Pro Gly Lys Ala Pro Lys Leu Leu Ile Tyr
        35
Asp Thr Ser Lys Leu Ala Ser Gly Val Pro Ser Arg Phe Ser Gly Ser
                       55
Gly Ser Gly Thr Glu Phe Thr Leu Thr Ile Ser Ser Leu Gln Pro Asp
                                      75
Asp Phe Ala Thr Tyr Tyr Cys Phe Gln Gly Ser Gly Tyr Pro Phe Thr
                                   90
                85
Phe Gly Gly Thr Lys Val Glu Ile Lys
            100
<210> 55
<211> 120
<212> PRT
<213> Artificial Sequence
<220>
<223> Humanized antibody - VH Domain
<400> 55
Gln Val Thr Leu Arg Glu Ser Gly Pro Ala Leu Val Lys Pro Thr Gln
                                   1.0
Thr Leu Thr Leu Thr Cys Thr Phe Ser Gly Phe Ser Leu Ser Thr Ala
                               25
Gly Met Ser Val Gly Trp Ile Arg Gln Pro Pro Gly Lys Ala Leu Glu
                           40
Trp Leu Ala Asp Ile Trp Trp Asp Asp Lys Lys Asp Tyr Asn Pro Ser
                       55
Leu Lys Ser Arg Leu Thr Ile Ser Lys Asp Thr Ser Lys Asn Gln Val
                   70
                                      75
Val Leu Lys Val Thr Asn Met Asp Pro Ala Asp Thr Ala Thr Tyr Tyr
                                   90
Cys Ala Arg Asp Met Ile Phe Asn Trp Tyr Phe Asp Val Trp Gly Gln
          100
                               105
Gly Thr Thr Val Thr Val Ser Ser
       115
```

```
<210> 56
<211> 106
<212> PRT
<213> Artificial Sequence
<223> Humanized antibody - VL Domain
<400> 56
Asp Ile Gln Met Thr Gln Ser Pro Ser Thr Leu Ser Ala Ser Val Gly
                                   10
Asp Arg Val Thr Ile Thr Cys Ser Ala Ser Ser Ser Val Gly Tyr Met
                               2.5
His Trp Tyr Gln Gln Lys Pro Gly Lys Ala Pro Lys Leu Leu Ile Tyr
                           40
Asp Thr Phe Lys Leu Ala Ser Gly Val Pro Ser Arg Phe Ser Gly Ser
                        55
Gly Ser Gly Thr Glu Phe Thr Leu Thr Ile Ser Ser Leu Gln Pro Asp
                                       75
                    70
Asp Phe Ala Thr Tyr Tyr Cys Phe Gln Gly Ser Gly Tyr Pro Phe Thr
             85
Phe Gly Gly Gly Thr Lys Val Glu Ile Lys
           100
<210> 57
<211> 7
<212> PRT
<213> Artificial Sequence
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 57
Asp Thr Tyr Lys Gln Thr Ser
<210> 58
<211> 106
<212> PRT
<213> Artificial Sequence
<220>
<223> Humanized antibody - VL Domain
<400> 58
Asp Ile Gln Met Thr Gln Ser Pro Ser Thr Leu Ser Ala Ser Val Gly
                                    10
Asp Arg Val Thr Ile Thr Cys Ser Ala Ser Ser Ser Val Gly Tyr Met
                                25
His Trp Tyr Gln Gln Lys Pro Gly Lys Ala Pro Lys Leu Leu Ile Tyr
                            40
 Asp Thr Arg Tyr Leu Ser Ser Gly Val Pro Ser Arg Phe Ser Gly Ser
 Gly Ser Gly Thr Glu Phe Thr Leu Thr Ile Ser Ser Leu Gln Pro Asp
                                       75
                   70
 Asp Phe Ala Thr Tyr Tyr Cys Phe Gln Gly Ser Gly Tyr Pro Phe Thr
                 85
 Phe Gly Gly Gly Thr Lys Val Glu Ile Lys
```

100 105 <210> 59 <211> 7 <212> PRT <213> Artificial Sequence <220> <223> Amino acid sequence derived from Murine monoclonal antibody and further modified by amino acid substitutions <400> 59 Asp Thr Arg Tyr Leu Ser Ser <210> 60 <211> 106 <212> PRT <213> Artificial Sequence <220> <223> Humanized antibody - VL Domain <400> 60 Asp Ile Gln Met Thr Gln Ser Pro Ser Thr Leu Ser Ala Ser Val Gly 10 Asp Arg Val Thr Ile Thr Cys Ser Ala Ser Ser Ser Val Gly Tyr Met 20 25 His Trp Tyr Gln Gln Lys Pro Gly Lys Ala Pro Lys Leu Leu Ile Tyr 40 Asp Thr Phe Lys Leu Ala Ser Gly Val Pro Ser Arg Phe Ser Gly Ser 55 Gly Ser Gly Thr Glu Phe Thr Leu Thr Ile Ser Ser Leu Gln Pro Asp 70 75 Asp Phe Ala Thr Tyr Tyr Cys Phe Gln Gly Ser Phe Tyr Pro Phe Thr 85 Phe Gly Gly Gly Thr Lys Val Glu Ile Lys 100 <210> 61 <211> 9 <212> PRT <213> Artificial Sequence <220> <223> Amino acid sequence derived from Murine monoclonal antibody and further modified by amino acid substitutions <400> 61 Phe Gln Gly Ser Phe Tyr Pro Phe Thr <210> 62 <211> 106 <212> PRT <213> Artificial Sequence <220> <223> Humanized antibody - VL Domain

```
<400> 62
Asp Ile Gln Met Thr Gln Ser Pro Ser Thr Leu Ser Ala Ser Val Gly
                    10
Asp Arg Val Thr Ile Thr Cys Ser Ala Ser Ser Ser Val Gly Tyr Met
His Trp Tyr Gln Gln Lys Pro Gly Lys Ala Pro Lys Leu Leu Ile Tyr
                          40
Asp Thr Phe Lys Leu Thr Ser Gly Val Pro Ser Arg Phe Ser Gly Ser
                       55
Gly Ser Gly Thr Glu Phe Thr Leu Thr Ile Ser Ser Leu Gln Pro Asp
                  70
Asp Phe Ala Thr Tyr Tyr Cys Phe Gln Gly Ser Gly Tyr Pro Phe Thr
              85
Phe Gly Gly Gly Thr Lys Val Glu Ile Lys
           100
<210> 63
<211> 7
<212> PRT
<213> Artificial Sequence
<223> Amino acid sequence derived from Murine monoclonal
     antibody and further modified by amino acid
      substitutions
<400> 63
Asp Thr Phe Lys Leu Thr Ser
<210> 64
<211> 106
<212> PRT
<213> Artificial Sequence
<220>
<223> Humanized antibody - VL Domain
<400> 64
Asp Ile Gln Met Thr Gln Ser Pro Ser Thr Leu Ser Ala Ser Val Gly
                                  10
Asp Arg Val Thr Ile Thr Cys Ser Ala Ser Ser Arg Val Gly Tyr Met
                               2.5
His Trp Tyr Gln Gln Lys Pro Gly Lys Ala Pro Lys Leu Leu Ile Tyr
                           40
Asp Thr Phe Lys Leu Ala Ser Gly Val Pro Ser Arg Phe Ser Gly Ser
                       55
Gly Ser Gly Thr Glu Phe Thr Leu Thr Ile Ser Ser Leu Gln Pro Asp
                                      75
                  70
Asp Phe Ala Thr Tyr Tyr Cys Phe Gln Gly Ser Gly Tyr Pro Phe Thr
                85
Phe Gly Gly Thr Lys Val Glu Ile Lys
            100
<210> 65
<211> 106
 <212> PRT
 <213> Artificial Sequence
<220>
```

```
<223> Humanized antibody - VL Domain
Asp Ile Gln Met Thr Gln Ser Pro Ser Thr Leu Ser Ala Ser Val Gly
                                    10
Asp Arg Val Thr Ile Thr Cys Ser Ala Ser Ser Ser Val Gly Tyr Met
                               25
His Trp Tyr Gln Gln Lys Pro Gly Lys Ala Pro Lys Leu Leu Ile Tyr
                            40
Asp Thr Phe Arg Leu Ala Ser Gly Val Pro Ser Arg Phe Ser Gly Ser
Gly Ser Gly Thr Glu Phe Thr Leu Thr Ile Ser Ser Leu Gln Pro Asp
                                       75
                   70
Asp Phe Ala Thr Tyr Tyr Cys Phe Gln Gly Ser Gly Tyr Pro Phe Thr
               85
Phe Gly Gly Thr Lys Val Glu Ile Lys
            100
<210> 66
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 66
Asp Thr Phe Arg Leu Ala Ser
<210> 67
<211> 07
<211> 120
<212> PRT
<213> Artificial Sequence
<220>
<223> Humanized antibody - VH Domain
Gln Val Thr Leu Arg Glu Ser Gly Pro Ala Leu Val Lys Pro Thr Gln
                                    10
Thr Leu Thr Leu Thr Cys Thr Phe Ser Gly Phe Ser Leu Ser Thr Ala
Gly Met Ser Val Gly Trp Ile Arg Gln Pro Pro Gly Lys Ala Leu Glu
Trp Leu Ala Asp Ile Trp Trp Asp Asp Lys Lys His Tyr Asn Pro Ser
                        55
Leu Lys Asp Arg Leu Thr Ile Ser Lys Asp Thr Ser Lys Asn Gln Val
                    70
                                        75
Val Leu Lys Val Thr Asn Met Asp Pro Ala Asp Thr Ala Thr Tyr Tyr
                                    90
Cys Ala Arg Asp Met Ile Phe Asn Trp Tyr Phe Asp Val Trp Gly Gln
            100 105
Gly Thr Thr Val Thr Val Ser Ser
<210> 68
<211> 106
<212> PRT
```

```
<213> Artificial Sequence
 <220>
 <223> Humanized antibody - VL Domain
 <400> 68
Asp Ile Gln Met Thr Gln Ser Pro Ser Thr Leu Ser Ala Ser Val Gly
                                     10
Asp Arg Val Thr Ile Thr Cys Ser Pro Ser Ser Arg Val Gly Tyr Met
His Trp Tyr Gln Gln Lys Pro Gly Lys Ala Pro Lys Leu Leu Ile Tyr
        35
                            40
Asp Thr Tyr Arg His Ser Ser Gly Val Pro Ser Arg Phe Ser Gly Ser
Gly Ser Gly Thr Glu Phe Thr Leu Thr Ile Ser Ser Leu Gln Pro Asp
                    70
Asp Phe Ala Thr Tyr Tyr Cys Phe Gln Gly Ser Gly Tyr Pro Phe Thr
                85
Phe Gly Gly Gly Thr Lys Val Glu Ile Lys
            100
<210> 69
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 69
Asp Thr Tyr Arg His Ser Ser
<210> 70
<211> 106
<212> PRT
<213> Artificial Sequence
<220>
<223> Humanized antibody - VL Domain
Asp Ile Gln Met Thr Gln Ser Pro Ser Thr Leu Ser Ala Ser Val Gly
                                    10
Asp Arg Val Thr Ile Thr Cys Ser Ala Ser Ser Ser Val Gly Tyr Met
                                25
His Trp Tyr Gln Gln Lys Pro Gly Lys Ala Pro Lys Leu Leu Ile Tyr
                            40
Asp Thr Tyr Lys Gln Thr Ser Gly Val Pro Ser Arg Phe Ser Gly Ser
                        55
Gly Ser Gly Thr Glu Phe Thr Leu Thr Ile Ser Ser Leu Gln Pro Asp
                    70
                                       75
Asp Phe Ala Thr Tyr Tyr Cys Phe Gln Gly Ser Gly Tyr Pro Phe Thr
Phe Gly Gly Gly Thr Lys Val Glu Ile Lys
<210> 71
```

22

```
<211> 106
<212> PRT
<213> Artificial Sequence
<223> Humanized antibody - VL Domain
<400> 71
Asp Ile Gln Met Thr Gln Ser Pro Ser Thr Leu Ser Ala Ser Val Gly
                                   10
Asp Arg Val Thr Ile Thr Cys Ser Leu Ser Ser Val Gly Tyr Met
                               25
His Trp Tyr Gln Gln Lys Pro Gly Lys Ala Pro Lys Leu Leu Ile Tyr
                           40
Asp Thr Phe Phe His Arg Ser Gly Val Pro Ser Arg Phe Ser Gly Ser
                       55
Gly Ser Gly Thr Glu Phe Thr Leu Thr Ile Ser Ser Leu Gln Pro Asp
                   70
                                       75
Asp Phe Ala Thr Tyr Tyr Cys Phe Gln Gly Ser Gly Tyr Pro Phe Thr
               85
                                   90
Phe Gly Gly Thr Lys Val Glu Ile Lys
<210> 72
<211> 10
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
     antibody and further modified by amino acid
     substitutions
<400> 72
Ser Leu Ser Ser Ser Val Gly Tyr Met His
<210> 73
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
     antibody and further modified by amino acid
     substitutions
<400> 73
Asp Thr Phe Phe His Arg Ser
<210> 74
<211> 106
<212> PRT
<213> Artificial Sequence
<220>
<223> Humanized antibody - VL Domain
<400> 74
```

```
Asp Ile Gln Met Thr Gln Ser Pro Ser Thr Leu Ser Ala Ser Val Gly
                                10
Asp Arg Val Thr Ile Thr Cys Ser Ala Ser Ser Arg Val Gly Tyr Met
                                25
His Trp Tyr Gln Gln Lys Pro Gly Lys Ala Pro Lys Leu Leu Ile Tyr
                            40
Asp Thr Leu Leu Leu Asp Ser Gly Val Pro Ser Arg Phe Ser Gly Ser
                        55
Gly Ser Gly Thr Glu Phe Thr Leu Thr Ile Ser Ser Leu Gln Pro Asp
                                        75
Asp Phe Ala Thr Tyr Tyr Cys Phe Gln Gly Ser Gly Tyr Pro Phe Thr
                                    90
                85
Phe Gly Gly Gly Thr Lys Val Glu Ile Lys
            100
<210> 75
<211> 7
<212> PRT
<213> Artificial Sequence
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 75
Asp Thr Leu Leu Leu Asp Ser
<210> 76
<211> 106
<212> PRT
<213> Artificial Sequence
<223> Humanized antibody - VL Domain
Asp Ile Gln Met Thr Gln Ser Pro Ser Thr Leu Ser Ala Ser Val Gly
                                     10
Asp Arg Val Thr Ile Thr Cys Ser Ala Ser Ser Arg Val Gly Tyr Met
His Trp Tyr Gln Gln Lys Pro Gly Lys Ala Pro Lys Leu Leu Ile Tyr
                             40
Asp Thr Ser Phe Leu Asp Ser Gly Val Pro Ser Arg Phe Ser Gly Ser
Gly Ser Gly Thr Glu Phe Thr Leu Thr Ile Ser Ser Leu Gln Pro Asp
                                         75
Asp Phe Ala Thr Tyr Tyr Cys Phe Gln Gly Ser Gly Tyr Pro Phe Thr
                85
Phe Gly Gly Gly Thr Lys Val Glu Ile Lys
<210> 77
 <211> 7
 <212> PRT
<213> Artificial Sequence
 <220>
 <223> Amino acid sequence derived from Murine monoclonal
```

antibody and further modified by amino acid substitutions

```
<400> 77
Asp Thr Ser Phe Leu Asp Ser
<210> 78
<211> 120
<212> PRT
<213> Artificial Sequence
<220>
<223> Humanized antibody - VH Domain
<400> 78
Gln Val Thr Leu Arg Glu Ser Gly Pro Ala Leu Val Lys Pro Thr Gln
                                  10
Thr Leu Thr Leu Thr Cys Thr Phe Ser Gly Phe Ser Leu Ser Thr Ala
                              25
Gly Met Ser Val Gly Trp Ile Arg Gln Pro Pro Gly Lys Ala Leu Glu
                          40
Trp Leu Ala Asp Ile Trp Trp Asp Asp Lys Lys Asp Tyr Asn Pro Ser
                     55
Leu Lys Ser Arg Leu Thr Ile Ser Lys Asp Thr Ser Lys Asn Gln Val
               70
                                      75
Val Leu Lys Val Thr Asn Met Asp Pro Ala Asp Thr Ala Thr Tyr Tyr
              85
                               90
Cys Ala Arg Asp Met Ile Thr Asn Phe Tyr Phe Asp Val Trp Gly Gln
          100 105
Gly Thr Thr Val Thr Val Ser Ser
       115
<210> 79
<211> 10
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
     antibody and further modified by amino acid
     substitutions
<400> 79
Asp Met Ile Thr Asn Phe Tyr Phe Asp Val
<210> 80
<211> 10
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
     antibody and further modified by amino acid
     substitutions
<400> 80
Lys Cys Gln Ser Ser Val Gly Tyr Met His
```

```
<210> 81
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 81
Asp Thr Ser Tyr Leu Ala Ser
                 5
<210> 82
<211> 16
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
Asp Ile Trp Trp Asp Asp Lys Lys His Tyr Asn Pro Ser Leu Lys Ser
                                  10
<210> 83
<211> 10
<211> 10
<212> PRT
<213> Artificial Sequence
<223> Amino acid sequence derived from Murine monoclonal
       antibody and further modified by amino acid
       substitutions
Asp Met Ile Thr Asn Trp Tyr Phe Asp Val
                 5
<210> 84
<211> 10
<212> PRT
<213> Artificial Sequence
 <223> Amino acid sequence derived from Murine monoclonal
       antibody and further modified by amino acid
       substitutions
 <400> 84
 Lys Cys Gln Ser Arg Val Gly Tyr Met His
 <210> 85
 <211> 7
 <212> PRT
 <213> Artificial Sequence
```

```
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 85
Asp Thr Ser Tyr Leu Ser Ser
               5
<210> 86
<211> 16
<212> PRT
<213> Artificial Sequence
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 86
Asp Ile Trp Trp Asp Asp Lys Lys Asp Tyr Asn Pro Ser Leu Lys Asp
                                    10
<210> 87
<211> 10
<212> PRT
<213> Artificial Sequence
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 87
Lys Cys Gln Leu Arg Val Gly Tyr Met His
<210> 88
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 88
Asp Thr Lys Lys Leu Ser Ser
<210> 89
<211> 10
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
       antibody and further modified by amino acid
       substitutions
```

```
<400> 89
Lys Leu Gln Leu Ser Val Gly Tyr Met His
<210> 90
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 90
Asp Thr Phe Tyr Leu Ser Ser
<210> 91
<211> 16
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 91
Asp Ile Trp Trp Asp Asp Lys Lys His Tyr Asn Pro Ser Leu Lys Ser
                                     10
<210> 92
<211> 10
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
       antibody and further modified by amino acid
       substitutions
 <400> 92
 Lys Leu Gln Ser Ser Val Gly Tyr Met His
 <210> 93
 <211> 16
 <212> PRT
 <213> Artificial Sequence
 <220>
 <223> Amino acid sequence derived from Murine monoclonal
       antibody and further modified by amino acid
       substitutions
 <400> 93
 Asp Ile Trp Trp Asp Asp Lys Lys His Tyr Asn Pro Ser Leu Lys Ser
                                      10
                 5
```

```
<210> 94
<211> 10
 <212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 94
Ser Met Ile Phe Asn Trp Tyr Phe Asp Val
                5
<210> 95
<211> 10
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 95
Lys Leu Gln Ser Arg Val Gly Tyr Met His
                5
<210> 96
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 96
Asp Thr Phe Lys Leu Ser Ser
<210> 97
<211> 10
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 97
Ser Met Ile Phe Asn Phe Tyr Phe Asp Val
               5
                  , 10
<210> 98
<211> 10
<212> PRT
<213> Artificial Sequence
```

```
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 98
Lys Leu Gln Leu Arg Val Gly Tyr Met His
               5
<210> 99
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 99
Asp Thr Phe Tyr Leu Ala Ser
                5
<210> 100
<211> 16
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 100
Asp Ile Trp Trp Asp Gly Lys Lys Asp Tyr Asn Pro Ser Leu Lys Ser
                           10
<210> 101
<211> 10
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
     antibody and further modified by amino acid
     substitutions
<400> 101
Lys Leu Ser Leu Ser Val Gly Tyr Met His
<210> 102
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
     antibody and further modified by amino acid
     substitutions
```

```
<400> 102
Asp Thr Ser Lys Leu Pro Ser
<210> 103
<211> 16
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
       substitutions
Asp Ile Trp Trp Asp Gly Lys Lys Asp Tyr Asn Pro Ser Leu Lys Asp
                                       10
<210> 104
<211> 10
<212> PRT
<213> Artificial Sequence
<223> Amino acid sequence derived from Murine monoclonal
       antibody and further modified by amino acid
       substitutions
Lys Leu Ser Ser Ser Val Gly Tyr Met His
<210> 105
<211> 7
<212> PRT
<213> Artificial Sequence
<223> Amino acid sequence derived from Murine monoclonal
       antibody and further modified by amino acid
       substitutions
<400> 105
Asp Thr Ser Gly Leu Ala Ser
<210> 106
<211> 16
<212> PRT
 <213> Artificial Sequence
 <220>
 <223> Amino acid sequence derived from Murine monoclonal
        antibody and further modified by amino acid
        substitutions
 <400> 106
 Asp Ile Trp Trp Asp Gly Lys Lys His Tyr Asn Pro Ser Leu Lys Ser
                                         10
```

```
<210> 107
<211> 10
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
Lys Leu Ser Ser Arg Val Gly Tyr Met His
                 5
<210> 108
<211> 7
<212> PRT
<213> Artificial Sequence
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 108
Asp Thr Ser Gly Leu Pro Ser
<210> 109
<211> 16
<212> PRT
<213> Artificial Sequence
<223> Amino acid sequence derived from Murine monoclonal
       antibody and further modified by amino acid
       substitutions
<400> 109
Asp Ile Trp Trp Asp Asp Lys Lys Ser Tyr Asn Pro Ser Leu Lys Ser
                                      10
<210> 110
<211> 10
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
       antibody and further modified by amino acid
       substitutions
<400> 110
Lys Leu Ser Leu Arg Val Gly Tyr Met His
                 5
<210> 111
 <211> 16
 <212> PRT
 <213> Artificial Sequence
```

```
<220>
<223> Amino acid sequence derived from Murine monoclonal
     antibody and further modified by amino acid
     substitutions
<400> 111
Asp Ile Trp Trp Asp Asp Lys Lys Ser Tyr Asn Pro Ser Leu Lys Asp
<210> 112
<211> 10
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 112
Lys Cys Ser Leu Ser Val Gly Tyr Met His
<210> 113
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 113
Asp Thr Arg Lys Leu Ala Ser
                5
<210> 114
<211> 16
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
Asp Ile Trp Trp Asp Gly Lys Lys Ser Tyr Asn Pro Ser Leu Lys Ser
                               10
<210> 115
<211> 10
<211> 10
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
```

```
<400> 115
Lys Cys Ser Ser Ser Val Gly Tyr Met His
<210> 116
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
     antibody and further modified by amino acid
     substitutions
<400> 116
Asp Thr Arg Gly Leu Ala Ser
<210> 117
<211> 10
<212> PRT
<213> Artificial Sequence
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 117
Lys Cys Ser Ser Arg Val Gly Tyr Met His
<210> 118
<211> 7
<212> PRT
<213> Artificial Sequence
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
     substitutions
<400> 118
Asp Thr Arg Lys Leu Pro Ser
<210> 119
<211> 10
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
     antibody and further modified by amino acid
     substitutions
<400> 119
Lys Cys Ser Leu Arg Val Gly Tyr Met His
                                   10
```

```
<210> 120
<211> 10
<212> PRT
<213> Artificial Sequence
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 120
Ser Leu Ser Leu Ser Val Gly Tyr Met His
<210> 121
<211> 7
<212> PRT
<213> Artificial Sequence
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 121
Asp Thr Met Lys Leu Ala Ser
<210> 122
<211> 10
<212> PRT
<213> Artificial Sequence
<223> Amino acid sequence derived from Murine monoclonal
       antibody and further modified by amino acid
       substitutions
<400> 122
Ser Leu Ser Ser Ser Val Gly Tyr Met His
<210> 123
<211> 7
<212> PRT
<213> Artificial Sequence
<223> Amino acid sequence derived from Murine monoclonal
       antibody and further modified by amino acid
       substitutions
<400> 123
Asp Thr Ser Arg Leu Ala Ser
<210> 124
<211> 7
<212> PRT
<213> Artificial Sequence
```

```
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 124
Asp Thr Ser Leu Leu Ala Ser
                 5
<210> 125
<211> 10
<212> PRT
<213> Artificial Sequence
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
Ser Leu Ser Leu Arg Val Gly Tyr Met His
                 5
<210> 126
<211> 7
<212> PRT
<213> Artificial Sequence
<223> Amino acid sequence derived from Murine monoclonal
       antibody and further modified by amino acid
       substitutions
<400> 126
Asp Thr Ser Leu Leu Asp Ser
<210> 127
<211> 10
<212> PRT
<213> Artificial Sequence
 <223> Amino acid sequence derived from Murine monoclonal
       antibody and further modified by amino acid
       substitutions
 <400> 127
 Ser Cys Gln Leu Ser Val Gly Tyr Met His
                                       10
 <210> 128
 <211> 7
 <212> PRT
 <213> Artificial Sequence
 <220>
 <223> Amino acid sequence derived from Murine monoclonal
       antibody and further modified by amino acid
       substitutions
```

```
<400> 128
Asp Thr Ser Lys Leu Asp Ser
    5
<210> 129
<211> 10
<212> PRT
<213> Artificial Sequence
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 129
Ser Cys Gln Ser Ser Val Gly Tyr Met His
     5
<210> 130
<211> 10
 <212> PRT
 <213> Artificial Sequence
 <223> Amino acid sequence derived from Murine monoclonal
       antibody and further modified by amino acid
       substitutions
 <400> 130
 Ser Cys Gln Ser Arg Val Gly Tyr Met His
                5
 <210> 131
<211> 7
<212> PRT
<213> Artificial Sequence
 <223> Amino acid sequence derived from Murine monoclonal
       antibody and further modified by amino acid
        substitutions
  Asp Thr Leu Lys Leu Asp Ser
                 5
  <210> 132
<211> 10
<212> PRT
<213> Artificial Sequence
  <223> Amino acid sequence derived from Murine monoclonal
        antibody and further modified by amino acid
        substitutions
  <400> 132
  Ser Cys Gln Leu Arg Val Gly Tyr Met His
  1
```

```
<210> 133
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 133
Asp Thr Leu Leu Leu Ala Ser
                5
<210> 134
<211> 10
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 134
Ser Leu Gln Leu Ser Val Gly Tyr Met His
                5
<210> 135
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 135
Asp Thr Leu Lys Leu Ala Ser
1
                5
<210> 136
<211> 10
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 136
Ser Leu Gln Ser Ser Val Gly Tyr Met His
<210> 137
<211> 7
<212> PRT
<213> Artificial Sequence
```

```
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 137
Asp Thr Ser Lys Leu Ser Ser
<210> 138
<211> 10
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 138
Ser Leu Gln Ser Arg Val Gly Tyr Met His
1
                5
                                     10
<210> 139
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 139
Asp Thr Ser Lys Gln Ala Ser
                5
<210> 140
<211> 10
<212> PRT
<213> Artificial Sequence
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 140
Ser Leu Gln Leu Arg Val Gly Tyr Met His
                5
<210> 141
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
```

substitutions

```
<400> 141
Asp Thr Ser Lys Gln Ser Ser
<210> 142
<211> 10
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
     antibody and further modified by amino acid
     substitutions
<400> 142
Ser Cys Ser Leu Ser Val Gly Tyr Met His
<210> 143
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 143
Asp Thr Ser Tyr Leu Ala Ser
<210> 144
<211> 10
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 144
Ser Cys Ser Ser Ser Val Gly Tyr Met His
<210> 145
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 145
Asp Thr Ser Tyr Leu Ser Ser
               5
```

```
<210> 146
<211> 10
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
     antibody and further modified by amino acid
      substitutions
<400> 146
Ser Cys Ser Ser Arg Val Gly Tyr Met His
<210> 147
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 147
Asp Thr Ser Tyr Gln Ala Ser
<210> 148
<211> 10
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 148
Ser Cys Ser Leu Arg Val Gly Tyr Met His
<210> 149
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 149
Asp Thr Ser Tyr Gln Ser Ser
<210> 150
<211> 10
<212> PRT
<213> Artificial Sequence
```

```
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 150
Lys Pro Ser Ser Arg Val Gly Tyr Met His
<210> 151
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 151
Asp Thr Met Tyr Gln Ala Ser
               5
<210> 152
<211> 10
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 152
Lys Pro Ser Leu Arg Val Gly Tyr Met His
<210> 153
<211> 10
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
Lys Pro Ser Ser Ser Val Gly Tyr Met His
             5
                    1.0
<210> 154
<211> 7
<212> PRT
<213> Artificial Sequence
 <223> Amino acid sequence derived from Murine monoclonal
       antibody and further modified by amino acid
       substitutions
```

```
<400> 154
Asp Thr Met Lys Gln Ala Ser
    5
<210> 155
<211> 10
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 155
Lys Pro Ser Leu Ser Val Gly Tyr Met His
<210> 156
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 156
Asp Thr Met Lys Gln Ser Ser
<210> 157
<211> 10
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
Lys Pro Gln Ser Arg Val Gly Tyr Met His
<210> 158
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 158
Asp Thr Met Tyr Leu Ala Ser
                5
```

```
<210> 159
<211> 10
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 159
Lys Pro Gln Leu Arg Val Gly Tyr Met His
<210> 160
<211> 7
<212> PRT
<213> Artificial Sequence
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 160
Asp Thr Met Tyr Leu Ser Ser
               5
<210> 161
<211> 10
<212> PRT
<213> Artificial Sequence
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 161
Lys Pro Gln Ser Ser Val Gly Tyr Met His
<210> 162
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 162
Asp Thr Met Lys Leu Ala Ser
<210> 163
<211> 10
<212> PRT
<213> Artificial Sequence
```

```
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 163
Lys Pro Gln Leu Ser Val Gly Tyr Met His
               5
<210> 164
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 164
Asp Thr Met Lys Leu Ser Ser
                5
<210> 165
<211> 7
<212> PRT
<213> Artificial Sequence
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 165
Asp Thr Ser Lys Leu Ser Ser
<210> 166
<211> 10
<212> PRT
<213> Artificial Sequence
 <223> Amino acid sequence derived from Murine monoclonal
       antibody and further modified by amino acid
       substitutions
 <400> 166
 Ser Pro Ser Leu Arg Val Gly Tyr Met His
                 5
 <210> 167
 <211> 7
 <212> PRT
 <213> Artificial Sequence
 <220>
 <223> Amino acid sequence derived from Murine monoclonal
       antibody and further modified by amino acid
```

substitutions

```
<400> 167
Asp Thr Arg Tyr Gln Ala Ser
<210> 168
<211> 10
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 168
Ser Pro Ser Ser Ser Val Gly Tyr Met His
            5
<210> 169
<211> 10
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 169
Ser Pro Ser Leu Ser Val Gly Tyr Met His
              5
<210> 170
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 170
Asp Thr Arg Tyr Gln Ala Ser
               5
<210> 171
<211> 10
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 171
Ser Pro Gln Ser Arg Val Gly Tyr Met His
               5
```

```
<210> 172
<211> 7
<212> PRT
<213> Artificial Sequence
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 172
Asp Thr Arg Lys Gln Ser Ser
               5
<210> 173
<211> 10
<212> PRT
<213> Artificial Sequence
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 173
Ser Pro Gln Leu Arg Val Gly Tyr Met His
<210> 174
<211> 7
<212> PRT
<213> Artificial Sequence
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 174
Asp Thr Arg Lys Leu Ala Ser
<210> 175
<211> 7
<212> PRT
<213> Artificial Sequence
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
       substitutions
<400> 175
Asp Thr Arg Lys Leu Ser Ser
                 5
 <210> 176
<211> 10
<212> PRT
```

```
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
     antibody and further modified by amino acid
      substitutions
<400> 176
Ser Pro Gln Ser Ser Val Gly Tyr Met His
<210> 177
<211> 10
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 177
Ser Pro Gln Leu Ser Val Gly Tyr Met His
               5
<210> 178
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 178
Asp Thr Arg Tyr Leu Ala Ser
<210> 179
<211> 10
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 179
Lys Ala Gln Ser Arg Val Gly Tyr Met His
<210> 180
<211> 10
<212> PRT
<213> Artificial Sequence
<220>
```

```
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 180
Lys Ala Gln Leu Arg Val Gly Tyr Met His
<210> 181
<211> 10
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 181
Lys Ala Gln Ser Ser Val Gly Tyr Met His
<210> 182
<211> 10
 <212> PRT
 <213> Artificial Sequence
<220>
 <223> Amino acid sequence derived from Murine monoclonal
       antibody and further modified by amino acid
       substitutions
 <400> 182
 Lys Ala Gln Leu Ser Val Gly Tyr Met His
                5
<210> 183
 <211> 10
 <212> PRT
 <213> Artificial Sequence
 <223> Amino acid sequence derived from Murine monoclonal
       antibody and further modified by amino acid
       substitutions
 <400> 183
 Lys Ala Ser Ser Arg Val Gly Tyr Met His
 <210> 184
 <211> 10
 <212> PRT
 <213> Artificial Sequence
 <220>
 <223> Amino acid sequence derived from Murine monoclonal
       antibody and further modified by amino acid
       substitutions
```

```
<400> 184
Lys Ala Ser Leu Arg Val Gly Tyr Met His
            5
<210> 185
<211> 10
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
     antibody and further modified by amino acid
     substitutions
<400> 185
Lys Ala Ser Ser Ser Val Gly Tyr Met His
               5
<210> 186
<211> 10
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 186
Lys Ala Ser Leu Ser Val Gly Tyr Met His
<210> 187
<211> 10
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 187
Ser Ala Ser Leu Arg Val Gly Tyr Met His
               5
<210> 188
<211> 10
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 188
Ser Ala Ser Leu Ser Val Gly Tyr Met His
                5
```

```
<210> 189
<211> 10
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 189
Ser Ala Gln Ser Arg Val Gly Tyr Met His
<210> 190
<211> 10
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 190
Ser Ala Gln Leu Arg Val Gly Tyr Met His
<210> 191
<211> 10
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 191
Ser Ala Gln Ser Ser Val Gly Tyr Met His
<210> 192
<211> 10
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 192
Leu Pro Ser Leu Ser Val Gly Tyr Met His
                 5
<210> 193
<211> 10
<212> PRT
<213> Artificial Sequence
```

```
<220>
<223> Amino acid sequence derived from Murine monoclonal
     antibody and further modified by amino acid
     substitutions
<400> 193
Leu Pro Ser Ser Ser Val Gly Tyr Met His
               5
<210> 194
<211> 10
<212> PRT
<213> Artificial Sequence
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 194
Leu Pro Ser Leu Arg Val Gly Tyr Met His
               5
<210> 195
<211> 10
<212> PRT
<213> Artificial Sequence
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 195
Leu Cys Ser Ser Arg Val Gly Tyr Met His
<210> 196
<211> 10
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 196
Leu Cys Ser Leu Ser Val Gly Tyr Met His
<210> 197
<211> 10
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
    substitutions
```

```
<400> 197
Leu Cys Ser Ser Ser Val Gly Tyr Met His
<210> 198
<211> 10
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 198
Leu Cys Ser Leu Arg Val Gly Tyr Met His
             5
<210> 199
<211> 10
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 199
Leu Pro Gln Ser Arg Val Gly Tyr Met His
               5
<210> 200
<211> 10
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 200
Leu Pro Gln Leu Ser Val Gly Tyr Met His
               5
<210> 201
<211> 10
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 201
Leu Pro Gln Ser Ser Val Gly Tyr Met His
                5
<210> 202
```

```
<211> 10
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 202
Leu Pro Gln Leu Arg Val Gly Tyr Met His
              5
<210> 203
<211> 10
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 203
Leu Cys Gln Ser Arg Val Gly Tyr Met His
<210> 204
<211> 10
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 204
Leu Cys Gln Leu Ser Val Gly Tyr Met His
<210> 205
<211> 10
<212> PRT
<213> Artificial Sequence
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 205
Leu Cys Gln Ser Ser Val Gly Tyr Met His
<210> 206
<211> 10
 <212> PRT
<213> Artificial Sequence
```

```
<220>
<223> Amino acid sequence derived from Murine monoclonal
     antibody and further modified by amino acid
     substitutions
<400> 206
Leu Cys Gln Leu Arg Val Gly Tyr Met His
<210> 207
<211> 10
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
     antibody and further modified by amino acid
      substitutions
<400> 207
Ser Ala Gln Leu Ser Val Gly Tyr Met His
<210> 208
<211> 450
<212> PRT
<213> Artificial Sequence
<220>
<223> Humanized antibody - VH Chain
<400> 208
Gln Val Thr Leu Arg Glu Ser Gly Pro Ala Leu Val Lys Pro Thr Gln
Thr Leu Thr Leu Thr Cys Thr Phe Ser Gly Phe Ser Leu Ser Thr Ser
                               25
Gly Met Ser Val Gly Trp Ile Arg Gln Pro Pro Gly Lys Ala Leu Glu
Trp Leu Ala Asp Ile Trp Trp Asp Asp Lys Lys Asp Tyr Asn Pro Ser
                        55
Leu Lys Ser Arg Leu Thr Ile Ser Lys Asp Thr Ser Lys Asn Gln Val
                                        75
                   70
Val Leu Lys Val Thr Asn Met Asp Pro Ala Asp Thr Ala Thr Tyr Tyr
Cys Ala Arg Ser Met Ile Thr Asn Trp Tyr Phe Asp Val Trp Gly Ala
                               105
            100
Gly Thr Thr Val Thr Val Ser Ser Ala Ser Thr Lys Gly Pro Ser Val
                           120
Phe Pro Leu Ala Pro Ser Ser Lys Ser Thr Ser Gly Gly Thr Ala Ala
                        1.35
                                           140
Leu Gly Cys Leu Val Lys Asp Tyr Phe Pro Glu Pro Val Thr Val Ser
                                       155
                    150
Trp Asn Ser Gly Ala Leu Thr Ser Gly Val His Thr Phe Pro Ala Val
                                   170
                165
Leu Gln Ser Ser Gly Leu Tyr Ser Leu Ser Ser Val Val Thr Val Pro
                                185
            180
Ser Ser Ser Leu Gly Thr Gln Thr Tyr Ile Cys Asn Val Asn His Lys
                           200
                                               205
Pro Ser Asn Thr Lys Val Asp Lys Arg Val Glu Pro Lys Ser Cys Asp
                        215
```

```
Lys Thr His Thr Cys Pro Pro Cys Pro Ala Pro Glu Leu Leu Gly Gly
225 230 235
Pro Ser Val Phe Leu Phe Pro Pro Lys Pro Lys Asp Thr Leu Met Ile
                           250
      245
Ser Arg Thr Pro Glu Val Thr Cys Val Val Val Asp Val Ser His Glu
 260 265 270
Asp Pro Glu Val Lys Phe Asn Trp Tyr Val Asp Gly Val Glu Val His
275 280 285
Asn Ala Lys Thr Lys Pro Arg Glu Glu Gln Tyr Asn Ser Thr Tyr Arg
  290 295 300
Val Val Ser Val Leu Thr Val Leu His Gln Asp Trp Leu Asn Gly Lys
305 310 315 320
Glu Tyr Lys Cys Lys Val Ser Asn Lys Ala Leu Pro Ala Pro Ile Glu
           325 330 335
Lys Thr Ile Ser Lys Ala Lys Gly Gln Pro Arg Glu Pro Gln Val Tyr
        340 345 350
Thr Leu Pro Pro Ser Arg Glu Glu Met Thr Lys Asn Gln Val Ser Leu
                     360 365
Thr Cys Leu Val Lys Gly Phe Tyr Pro Ser Asp Ile Ala Val Glu Trp
 370 375 380
Glu Ser Asn Gly Gln Pro Glu Asn Asn Tyr Lys Thr Thr Pro Pro Val
               390 395 400
Leu Asp Ser Asp Gly Ser Phe Phe Leu Tyr Ser Lys Leu Thr Val Asp
           405 410 415
Lys Ser Arg Trp Gln Gln Gly Asn Val Phe Ser Cys Ser Val Met His
   420 425 430
Glu Ala Leu His Asn His Tyr Thr Gln Lys Ser Leu Ser Leu Ser Pro
                     440
Gly Lys
 450
<210> 209
<211> 213
<212> PRT
<213> Artificial Sequence
<223> Humanized antibody - VL Chain
<400> 209
Asp Ile Gln Met Thr Gln Ser Pro Ser Thr Leu Ser Ala Ser Val Gly
Asp Arg Val Thr Ile Thr Cys Lys Cys Gln Leu Ser Val Gly Tyr Met
His Trp Tyr Gln Gln Lys Pro Gly Lys Ala Pro Lys Leu Leu Ile Tyr
                     40
Asp Thr Ser Lys Leu Ala Ser Gly Val Pro Ser Arg Phe Ser Gly Ser
                  55
Gly Ser Gly Thr Glu Phe Thr Leu Thr Ile Ser Ser Leu Gln Pro Asp
Asp Phe Ala Thr Tyr Tyr Cys Phe Gln Gly Ser Gly Tyr Pro Phe Thr
Phe Gly Gly Gly Thr Lys Leu Glu Ile Lys Arg Thr Val Ala Ala Pro
                        105
Ser Val Phe Ile Phe Pro Pro Ser Asp Glu Gln Leu Lys Ser Gly Thr
      115 120
Ala Ser Val Val Cys Leu Leu Asn Asn Phe Tyr Pro Arg Glu Ala Lys
  130 135
                                  140
Val Gln Trp Lys Val Asp Asn Ala Leu Gln Ser Gly Asn Ser Gln Glu
                               155
             150
Ser Val Thr Glu Gln Asp Ser Lys Asp Ser Thr Tyr Ser Leu Ser Ser
```

```
170
Thr Leu Thr Leu Ser Lys Ala Asp Tyr Glu Lys His Lys Val Tyr Ala
                 185
Cys Glu Val Thr His Gln Gly Leu Ser Ser Pro Val Thr Lys Ser Phe
 195
                      200
Asn Arg Gly Glu Cys
   210
<210> 210
<211> 450
<212> PRT
<213> Artificial Sequence
<220>
<223> Humanized antibody - VH Chain
<400> 210
Gln Val Thr Leu Arg Glu Ser Gly Pro Ala Leu Val Lys Pro Thr Gln
Thr Leu Thr Leu Thr Cys Thr Phe Ser Gly Phe Ser Leu Ser Thr Ala
                          25
Gly Met Ser Val Gly Trp Ile Arg Gln Pro Pro Gly Lys Ala Leu Glu
                   40
Trp Leu Ala Asp Ile Trp Trp Asp Asp Lys Lys Asp Tyr Asn Pro Ser
                  55
Leu Lys Ser Arg Leu Thr Ile Ser Lys Asp Thr Ser Lys Asn Gln Val
65 70
Val Leu Lys Val Thr Asn Met Asp Pro Ala Asp Thr Ala Thr Tyr Tyr
                            90 95
    85
Cys Ala Arg Ser Met Ile Thr Asn Phe Tyr Phe Asp Val Trp Gly Gln
   100 105
Gly Thr Thr Val Thr Val Ser Ser Ala Ser Thr Lys Gly Pro Ser Val
 115 120 125
Phe Pro Leu Ala Pro Ser Ser Lys Ser Thr Ser Gly Gly Thr Ala Ala
  130 135 140
Leu Gly Cys Leu Val Lys Asp Tyr Phe Pro Glu Pro Val Thr Val Ser
    150 155
Trp Asn Ser Gly Ala Leu Thr Ser Gly Val His Thr Phe Pro Ala Val
            165 170 175
Leu Gln Ser Ser Gly Leu Tyr Ser Leu Ser Ser Val Val Thr Val Pro
         180 185
Ser Ser Ser Leu Gly Thr Gln Thr Tyr Ile Cys Asn Val Asn His Lys
                      200
Pro Ser Asn Thr Lys Val Asp Lys Arg Val Glu Pro Lys Ser Cys Asp
   210 215
                                   220
Lys Thr His Thr Cys Pro Pro Cys Pro Ala Pro Glu Leu Leu Gly Gly
      230 235
Pro Ser Val Phe Leu Phe Pro Pro Lys Pro Lys Asp Thr Leu Met Ile
            245
                            250
Ser Arg Thr Pro Glu Val Thr Cys Val Val Val Asp Val Ser His Glu
                          265 270
         260
Asp Pro Glu Val Lys Phe Asn Trp Tyr Val Asp Gly Val Glu Val His
                      280
Asn Ala Lys Thr Lys Pro Arg Glu Glu Gln Tyr Asn Ser Thr Tyr Arg
                   295
                                   300
Val Val Ser Val Leu Thr Val Leu His Gln Asp Trp Leu Asn Gly Lys
                                315 320
Glu Tyr Lys Cys Lys Val Ser Asn Lys Ala Leu Pro Ala Pro Ile Glu
                            330 335
Lys Thr Ile Ser Lys Ala Lys Gly Gln Pro Arg Glu Pro Gln Val Tyr
                          345
```

```
Thr Leu Pro Pro Ser Arg Glu Glu Met Thr Lys Asn Gln Val Ser Leu
355 360 365
Thr Cys Leu Val Lys Gly Phe Tyr Pro Ser Asp Ile Ala Val Glu Trp
370 375 380
Glu Ser Asn Gly Gln Pro Glu Asn Asn Tyr Lys Thr Thr Pro Pro Val
385 390 395
Leu Asp Ser Asp Gly Ser Phe Phe Leu Tyr Ser Lys Leu Thr Val Asp
      405 410 415
Lys Ser Arg Trp Gln Gln Gly Asn Val Phe Ser Cys Ser Val Met His
 420 425 430
Glu Ala Leu His Asn His Tyr Thr Gln Lys Ser Leu Ser Leu Ser Pro
435 440
Gly Lys
 450
<210> 211
<211> 213
<212> PRT
<213> Artificial Sequence
<220>
<223> Humanized antibody - VL Chain
<400> 211
Asp Ile Gln Met Thr Gln Ser Pro Ser Thr Leu Ser Ala Ser Val Gly
               10 15
Asp Arg Val Thr Ile Thr Cys Ser Ala Ser Ser Ser Val Gly Tyr Met
              25 30
His Trp Tyr Gln Gln Lys Pro Gly Lys Ala Pro Lys Leu Leu Ile Tyr
          40
Asp Thr Phe Lys Leu Ala Ser Gly Val Pro Ser Arg Phe Ser Gly Ser
                 55 60
Gly Ser Gly Thr Glu Phe Thr Leu Thr Ile Ser Ser Leu Gln Pro Asp
         70 75 80
Asp Phe Ala Thr Tyr Tyr Cys Phe Gln Phe Ser Gly Tyr Pro Phe Thr
               90 95
Phe Gly Gly Gly Thr Lys Val Glu Ile Lys Arg Thr Val Ala Ala Pro
       100 105 110
Ser Val Phe Ile Phe Pro Pro Ser Asp Glu Gln Leu Lys Ser Gly Thr
     115 120 125
Ala Ser Val Val Cys Leu Leu Asn Asn Phe Tyr Pro Arg Glu Ala Lys
  130 135 140
Val Gln Trp Lys Val Asp Asn Ala Leu Gln Ser Gly Asn Ser Gln Glu
      150 155 160
Ser Val Thr Glu Gln Asp Ser Lys Asp Ser Thr Tyr Ser Leu Ser Ser
           165 170 175
Thr Leu Thr Leu Ser Lys Ala Asp Tyr Glu Lys His Lys Val Tyr Ala
        180 185 190
Cys Glu Val Thr His Gln Gly Leu Ser Ser Pro Val Thr Lys Ser Phe
Asn Arg Gly Glu Cys
  210
<210> 212
<211> 450
<212> PRT
<213> Artificial Sequence
<223> Humanized antibody - VH Chain
```

```
<400> 212
Gln Val Thr Leu Arg Glu Ser Gly Pro Ala Leu Val Lys Pro Thr Gln
                             10
Thr Leu Thr Leu Thr Cys Thr Phe Ser Gly Phe Ser Leu Ser Thr Pro
                          25
Gly Met Ser Val Gly Trp Ile Arg Gln Pro Pro Gly Lys Ala Leu Glu
                       40
Trp Leu Ala Asp Ile Trp Trp Asp Asp Lys Lys His Tyr Asn Pro Ser
Leu Lys Asp Arg Leu Thr Ile Ser Lys Asp Thr Ser Lys Asn Gln Val
                                 75
Val Leu Lys Val Thr Asn Met Asp Pro Ala Asp Thr Ala Thr Tyr Tyr
                              90
             8.5
Cys Ala Arg Asp Met Ile Phe Asn Phe Tyr Phe Asp Val Trp Gly Gln
                          105
          100
Gly Thr Thr Val Thr Val Ser Ser Ala Ser Thr Lys Gly Pro Ser Val
                                       125
                       120
Phe Pro Leu Ala Pro Ser Ser Lys Ser Thr Ser Gly Gly Thr Ala Ala
                                    140
                    135
Leu Gly Cys Leu Val Lys Asp Tyr Phe Pro Glu Pro Val Thr Val Ser
                                 155 160
                150
Trp Asn Ser Gly Ala Leu Thr Ser Gly Val His Thr Phe Pro Ala Val
                            170 175
             165
Leu Gln Ser Ser Gly Leu Tyr Ser Leu Ser Ser Val Val Thr Val Pro
                           185
          180
Ser Ser Ser Leu Gly Thr Gln Thr Tyr Ile Cys Asn Val Asn His Lys
                            205
                       200
Pro Ser Asn Thr Lys Val Asp Lys Arg Val Glu Pro Lys Ser Cys Asp
                    215
                                    220
Lys Thr His Thr Cys Pro Pro Cys Pro Ala Pro Glu Leu Leu Gly Gly
                                 235
                 230
Pro Ser Val Phe Leu Phe Pro Pro Lys Pro Lys Asp Thr Leu Met Ile
                              250 255
             245
Ser Arg Thr Pro Glu Val Thr Cys Val Val Val Asp Val Ser His Glu
                           265 270
           260
Asp Pro Glu Val Lys Phe Asn Trp Tyr Val Asp Gly Val Glu Val His
                       280
Asn Ala Lys Thr Lys Pro Arg Glu Glu Gln Tyr Asn Ser Thr Tyr Arg
                                     300
                  295
Val Val Ser Val Leu Thr Val Leu His Gln Asp Trp Leu Asn Gly Lys
                                  315
                 310
 Glu Tyr Lys Cys Lys Val Ser Asn Lys Ala Leu Pro Ala Pro Ile Glu
                               330 335
              325
 Lys Thr Ile Ser Lys Ala Lys Gly Gln Pro Arg Glu Pro Gln Val Tyr
                           345
           340
 Thr Leu Pro Pro Ser Arg Glu Glu Met Thr Lys Asn Gln Val Ser Leu
                        360
 Thr Cys Leu Val Lys Gly Phe Tyr Pro Ser Asp Ile Ala Val Glu Trp
                     375
 Glu Ser Asn Gly Gln Pro Glu Asn Asn Tyr Lys Thr Thr Pro Pro Val
                                  395
                 390
 Leu Asp Ser Asp Gly Ser Phe Phe Leu Tyr Ser Lys Leu Thr Val Asp
              405
                               410
 Lys Ser Arg Trp Gln Gln Gly Asn Val Phe Ser Cys Ser Val Met His
                           425 430
 Glu Ala Leu His Asn His Tyr Thr Gln Lys Ser Leu Ser Leu Ser Pro
                        440
 Gly Lys
    450
```

<210> 213

```
<211> 213
<212> PRT
<213> Artificial Sequence
<220>
<223> Humanized antibody - VL Chain
<400> 213
Asp Ile Gln Met Thr Gln Ser Pro Ser Thr Leu Ser Ala Ser Val Glv
                    10 15
Asp Arg Val Thr Ile Thr Cys Ser Leu Ser Ser Arg Val Gly Tyr Met
                     25
His Trp Tyr Gln Gln Lys Pro Gly Lys Ala Pro Lys Leu Leu Ile Tyr
                       40
Asp Thr Phe Tyr Leu Ser Ser Gly Val Pro Ser Arg Phe Ser Gly Ser
Gly Ser Gly Thr Glu Phe Thr Leu Thr Ile Ser Ser Leu Gln Pro Asp
                      75
Asp Phe Ala Thr Tyr Tyr Cys Phe Gln Gly Ser Gly Tyr Pro Phe Thr
                              90
Phe Gly Gly Gly Thr Lys Val Glu Ile Lys Arg Thr Val Ala Ala Pro
                  105
Ser Val Phe Ile Phe Pro Pro Ser Asp Glu Gln Leu Lys Ser Gly Thr
      115 120 125
Ala Ser Val Val Cys Leu Leu Asn Asn Phe Tyr Pro Arg Glu Ala Lys
                    135 140
Val Gln Trp Lys Val Asp Asn Ala Leu Gln Ser Gly Asn Ser Gln Glu
              150 155 160
Ser Val Thr Glu Gln Asp Ser Lys Asp Ser Thr Tyr Ser Leu Ser Ser
             165 170 175
Thr Leu Thr Leu Ser Lys Ala Asp Tyr Glu Lys His Lys Val Tyr Ala
   180 185 190
Cys Glu Val Thr His Gln Gly Leu Ser Ser Pro Val Thr Lys Ser Phe
                       200
Asn Arg Gly Glu Cys
  210
<210> 214
<211> 450
<212> PRT
<213> Artificial Sequence
<223> Humanized antibody - VH Chain
<400> 214
Gln Val Thr Leu Arg Glu Ser Gly Pro Ala Leu Val Lys Pro Thr Gln
Thr Leu Thr Leu Thr Cys Thr Phe Ser Gly Phe Ser Leu Ser Thr Pro
                           25
Gly Met Ser Val Gly Trp Ile Arg Gln Pro Pro Gly Lys Ala Leu Glu
                        40
Trp Leu Ala Asp Ile Trp Trp Asp Gly Lys Lys His Tyr Asn Pro Ser
                     55
Leu Lys Asp Arg Leu Thr Ile Ser Lys Asp Thr Ser Lys Asn Gln Val
                                  75
                 70
Val Leu Lys Val Thr Asn Met Asp Pro Ala Asp Thr Ala Thr Tyr Tyr
              85
                             90
Cys Ala Arg Asp Met Ile Phe Asn Phe Tyr Phe Asp Val Trp Gly Gln
                   105
           100
 Gly Thr Thr Val Thr Val Ser Ser Ala Ser Thr Lys Gly Pro Ser Val
```

```
115
                           120
Phe Pro Leu Ala Pro Ser Ser Lys Ser Thr Ser Gly Gly Thr Ala Ala
                      135
                                          140
Leu Gly Cys Leu Val Lys Asp Tyr Phe Pro Glu Pro Val Thr Val Ser
                  150
                            155
Trp Asn Ser Gly Ala Leu Thr Ser Gly Val His Thr Phe Pro Ala Val
              165
                                 170
Leu Gln Ser Ser Gly Leu Tyr Ser Leu Ser Ser Val Val Thr Val Pro
           180
                               185
Ser Ser Ser Leu Gly Thr Gln Thr Tyr Ile Cys Asn Val Asn His Lys
                           200
Pro Ser Asn Thr Lys Val Asp Lys Arg Val Glu Pro Lys Ser Cys Asp
                       215
                                          220
Lys Thr His Thr Cys Pro Pro Cys Pro Ala Pro Glu Leu Leu Gly Gly
                   230
                                       235
Pro Ser Val Phe Leu Phe Pro Pro Lys Pro Lys Asp Thr Leu Met Ile
                                   250
Ser Arg Thr Pro Glu Val Thr Cys Val Val Val Asp Val Ser His Glu
                               265
Asp Pro Glu Val Lys Phe Asn Trp Tyr Val Asp Gly Val Glu Val His
                          280
Asn Ala Lys Thr Lys Pro Arg Glu Glu Gln Tyr Asn Ser Thr Tyr Arg
                      295
                                          300
Val Val Ser Val Leu Thr Val Leu His Gln Asp Trp Leu Asn Gly Lys
                   310
                                      315
Glu Tyr Lys Cys Lys Val Ser Asn Lys Ala Leu Pro Ala Pro Ile Glu
               325
                                   330
Lys Thr Ile Ser Lys Ala Lys Gly Gln Pro Arg Glu Pro Gln Val Tyr
                               345
Thr Leu Pro Pro Ser Arg Glu Glu Met Thr Lys Asn Gln Val Ser Leu
                           360
Thr Cys Leu Val Lys Gly Phe Tyr Pro Ser Asp Ile Ala Val Glu Trp
                       375
                               380
Glu Ser Asn Gly Gln Pro Glu Asn Asn Tyr Lys Thr Thr Pro Pro Val
                        395
                   390
Leu Asp Ser Asp Gly Ser Phe Phe Leu Tyr Ser Lys Leu Thr Val Asp
               405
                                  410
Lys Ser Arg Trp Gln Gln Gly Asn Val Phe Ser Cys Ser Val Met His
                              425
Glu Ala Leu His Asn His Tyr Thr Gln Lys Ser Leu Ser Leu Ser Pro
                          440
Gly Lys
  450
<210> 215
<211> 213
<212> PRT
<213> Artificial Sequence
<223> Humanized antibody - VL Chain
Asp Ile Gln Met Thr Gln Ser Pro Ser Thr Leu Ser Ala Ser Val Gly
Asp Arg Val Thr Ile Thr Cys Ser Leu Ser Ser Arg Val Gly Tyr Met
His Trp Tyr Gln Gln Lys Pro Gly Lys Ala Pro Lys Leu Leu Ile Tyr
                          40
Asp Thr Arg Gly Leu Pro Ser Gly Val Pro Ser Arg Phe Ser Gly Ser
```

```
Gly Ser Gly Thr Glu Phe Thr Leu Thr Ile Ser Ser Leu Gln Pro Asp
                   70
                                      75
 Asp Phe Ala Thr Tyr Tyr Cys Phe Gln Gly Ser Gly Tyr Pro Phe Thr
               85
                                 90
 Phe Gly Gly Gly Thr Lys Val Glu Ile Lys Arg Thr Val Ala Ala Pro
            100
                             105
 Ser Val Phe Ile Phe Pro Pro Ser Asp Glu Gln Leu Lys Ser Gly Thr
             120
 Ala Ser Val Val Cys Leu Leu Asn Asn Phe Tyr Pro Arg Glu Ala Lys
          135
                                        140
 Val Gln Trp Lys Val Asp Asn Ala Leu Gln Ser Gly Asn Ser Gln Glu
         150
                         155
 Ser Val Thr Glu Gln Asp Ser Lys Asp Ser Thr Tyr Ser Leu Ser Ser
               165
                           170
Thr Leu Thr Leu Ser Lys Ala Asp Tyr Glu Lys His Lys Val Tyr Ala
           180
                            185
Cys Glu Val Thr His Gln Gly Leu Ser Ser Pro Val Thr Lys Ser Phe
     195
                          200
Asn Arg Gly Glu Cys
  210
<210> 216
<211> 450
<212> PRT
<213> Artificial Sequence
<220>
<223> Humanized antibody - VH Chain
<400> 216
Gln Val Thr Leu Arg Glu Ser Gly Pro Ala Leu Val Lys Pro Thr Gln
                                 10
Thr Leu Thr Leu Thr Cys Thr Phe Ser Gly Phe Ser Leu Ser Thr Pro
Gly Met Ser Val Gly Trp Ile Arg Gln Pro Pro Gly Lys Ala Leu Glu
Trp Leu Ala Asp Ile Trp Trp Asp Gly Lys Lys His Tyr Asn Pro Ser
                    55
Leu Lys Asp Arg Leu Thr Ile Ser Lys Asp Thr Ser Lys Asn Gln Val
                                     75
Val Leu Lys Val Thr Asn Met Asp Pro Ala Asp Thr Ala Thr Tyr Tyr
              85
                              90
Cys Ala Arg Asp Met Ile Phe Asn Trp Tyr Phe Asp Val Trp Gly Gln
                             105 110
Gly Thr Thr Val Thr Val Ser Ser Ala Ser Thr Lys Gly Pro Ser Val
                         120
Phe Pro Leu Ala Pro Ser Ser Lys Ser Thr Ser Gly Gly Thr Ala Ala
                      135
                                        140
Leu Gly Cys Leu Val Lys Asp Tyr Phe Pro Glu Pro Val Thr Val Ser
                  150
                                     155
Trp Asn Ser Gly Ala Leu Thr Ser Gly Val His Thr Phe Pro Ala Val
                                 170
Leu Gln Ser Ser Gly Leu Tyr Ser Leu Ser Ser Val Val Thr Val Pro
                              185
Ser Ser Ser Leu Gly Thr Gln Thr Tyr Ile Cys Asn Val Asn His Lys
                          200
Pro Ser Asn Thr Lys Val Asp Lys Arg Val Glu Pro Lys Ser Cys Asp
                     215
Lys Thr His Thr Cys Pro Pro Cys Pro Ala Pro Glu Leu Leu Gly Gly
                  230
Pro Ser Val Phe Leu Phe Pro Pro Lys Pro Lys Asp Thr Leu Met Ile
```

```
250
               245
Ser Arg Thr Pro Glu Val Thr Cys Val Val Val Asp Val Ser His Glu
           260
                              2.65
Asp Pro Glu Val Lys Phe Asn Trp Tyr Val Asp Gly Val Glu Val His
                          280
Asn Ala Lys Thr Lys Pro Arg Glu Glu Gln Tyr Asn Ser Thr Tyr Arg
                      295
Val Val Ser Val Leu Thr Val Leu His Gln Asp Trp Leu Asn Gly Lys
                   310
                                      315
Glu Tyr Lys Cys Lys Val Ser Asn Lys Ala Leu Pro Ala Pro Ile Glu
               325
                                  330
Lys Thr Ile Ser Lys Ala Lys Gly Gln Pro Arg Glu Pro Gln Val Tyr
                              345
Thr Leu Pro Pro Ser Arg Glu Glu Met Thr Lys Asn Gln Val Ser Leu
Thr Cys Leu Val Lys Gly Phe Tyr Pro Ser Asp Ile Ala Val Glu Trp
                      375
                                         380
Glu Ser Asn Gly Gln Pro Glu Asn Asn Tyr Lys Thr Thr Pro Pro Val
                                   395
Leu Asp Ser Asp Gly Ser Phe Phe Leu Tyr Ser Lys Leu Thr Val Asp
                                 410 415
              405
Lys Ser Arg Trp Gln Gln Gly Asn Val Phe Ser Cys Ser Val Met His
          420 425 430
Glu Ala Leu His Asn His Tyr Thr Gln Lys Ser Leu Ser Leu Ser Pro
Gly Lys
  450
<210> 217
<211> 213
<212> PRT
<213> Artificial Sequence
<220>
<223> Humanized antibody - VL Chain
<400> 217
Asp Ile Gln Met Thr Gln Ser Pro Ser Thr Leu Ser Ala Ser Val Gly
                                  10
Asp Arg Val Thr Ile Thr Cys Ser Pro Ser Ser Arg Val Gly Tyr Met
His Trp Tyr Gln Gln Lys Pro Gly Lys Ala Pro Lys Leu Leu Ile Tyr
                          40
Asp Thr Met Arg Leu Ala Ser Gly Val Pro Ser Arg Phe Ser Gly Ser
                      55
Gly Ser Gly Thr Glu Phe Thr Leu Thr Ile Ser Ser Leu Gln Pro Asp
                  70
                                     75
Asp Phe Ala Thr Tyr Tyr Cys Phe Gln Gly Ser Gly Tyr Pro Phe Thr
              8.5
                                 90
Phe Gly Gly Gly Thr Lys Val Glu Ile Lys Arg Thr Val Ala Ala Pro
                            105
           100
Ser Val Phe Ile Phe Pro Pro Ser Asp Glu Gln Leu Lys Ser Gly Thr
                         120
                                            125
Ala Ser Val Val Cys Leu Leu Asn Asn Phe Tyr Pro Arg Glu Ala Lys
                     135
                                         140
Val Gln Trp Lys Val Asp Asn Ala Leu Gln Ser Gly Asn Ser Gln Glu
               150
                                     155 160
Ser Val Thr Glu Gln Asp Ser Lys Asp Ser Thr Tyr Ser Leu Ser Ser
              165 170
Thr Leu Thr Leu Ser Lys Ala Asp Tyr Glu Lys His Lys Val Tyr Ala
                             185
```

```
Cys Glu Val Thr His Gln Gly Leu Ser Ser Pro Val Thr Lys Ser Phe
       195
                           200
Asn Arg Gly Glu Cys
  210
<210> 218
<211> 450
<212> PRT
<213> Artificial Sequence
<220>
<223> Humanized antibody - VH Chain
<400> 218
Gln Val Thr Leu Arg Glu Ser Gly Pro Ala Leu Val Lys Pro Thr Gln
                                 10
Thr Leu Thr Leu Thr Cys Thr Phe Ser Gly Phe Ser Leu Ser Thr Ala
                             25
Gly Met Ser Val Gly Trp Ile Arg Gln Pro Pro Gly Lys Ala Leu Glu
                       40
Trp Leu Ala Asp Ile Trp Trp Asp Gly Lys Lys His Tyr Asn Pro Ser
                   55
Leu Lys Asp Arg Leu Thr Ile Ser Lys Asp Thr Ser Lys Asn Gln Val
Val Leu Lys Val Thr Asn Met Asp Pro Ala Asp Thr Ala Thr Tyr Tyr
                                 90
Cys Ala Arg Asp Met Ile Phe Asn Trp Tyr Phe Asp Val Trp Gly Gln
               105
Gly Thr Thr Val Thr Val Ser Ser Ala Ser Thr Lys Gly Pro Ser Val
                         120
                              125
Phe Pro Leu Ala Pro Ser Ser Lys Ser Thr Ser Gly Gly Thr Ala Ala
                      135
                             140
Leu Gly Cys Leu Val Lys Asp Tyr Phe Pro Glu Pro Val Thr Val Ser
                  150
                          155
Trp Asn Ser Gly Ala Leu Thr Ser Gly Val His Thr Phe Pro Ala Val
                                 170 175
Leu Gln Ser Ser Gly Leu Tyr Ser Leu Ser Ser Val Val Thr Val Pro
                              185
Ser Ser Ser Leu Gly Thr Gln Thr Tyr Ile Cys Asn Val Asn His Lys
                          200
Pro Ser Asn Thr Lys Val Asp Lys Arg Val Glu Pro Lys Ser Cys Asp
                  215
                                        220
Lys Thr His Thr Cys Pro Pro Cys Pro Ala Pro Glu Leu Leu Gly Gly
                  230 235
Pro Ser Val Phe Leu Phe Pro Pro Lys Pro Lys Asp Thr Leu Met Ile
                                 250
Ser Arg Thr Pro Glu Val Thr Cys Val Val Val Asp Val Ser His Glu
                             265
Asp Pro Glu Val Lys Phe Asn Trp Tyr Val Asp Gly Val Glu Val His
                          280
Asn Ala Lys Thr Lys Pro Arg Glu Glu Gln Tyr Asn Ser Thr Tyr Arg
                     295
Val Val Ser Val Leu Thr Val Leu His Gln Asp Trp Leu Asn Gly Lys
                  310
Glu Tyr Lys Cys Lys Val Ser Asn Lys Ala Leu Pro Ala Pro Ile Glu
              325
                                 330
Lys Thr Ile Ser Lys Ala Lys Gly Gln Pro Arg Glu Pro Gln Val Tyr
Thr Leu Pro Pro Ser Arg Glu Glu Met Thr Lys Asn Gln Val Ser Leu
                          360
Thr Cys Leu Val Lys Gly Phe Tyr Pro Ser Asp Ile Ala Val Glu Trp
```

```
370
                      375
                                         380
Glu Ser Asn Gly Gln Pro Glu Asn Asn Tyr Lys Thr Thr Pro Pro Val
385 390
                                     395
Leu Asp Ser Asp Gly Ser Phe Phe Leu Tyr Ser Lys Leu Thr Val Asp
        405
                                 410 415
Lys Ser Arg Trp Gln Gln Gly Asn Val Phe Ser Cys Ser Val Met His
          420
                             425
Glu Ala Leu His Asn His Tyr Thr Gln Lys Ser Leu Ser Leu Ser Pro
Gly Lys
  450
<210> 219
<211> 213
<212> PRT
<213> Artificial Sequence
<220>
<223> Humanized antibody - VL Chain
<400> 219
Asp Ile Gln Met Thr Gln Ser Pro Ser Thr Leu Ser Ala Ser Val Gly
                                  10
Asp Arg Val Thr Ile Thr Cys Ser Leu Ser Ser Arg Val Gly Tyr Met
                              25
His Trp Tyr Gln Gln Lys Pro Gly Lys Ala Pro Lys Leu Leu Ile Tyr
                          4 ()
Asp Thr Phe Lys Leu Ser Ser Gly Val Pro Ser Arg Phe Ser Gly Ser
                      55
Gly Ser Gly Thr Glu Phe Thr Leu Thr Ile Ser Ser Leu Gln Pro Asp
                                     75
Asp Phe Ala Thr Tyr Tyr Cys Phe Gln Gly Ser Gly Tyr Pro Phe Thr
                                  90
Phe Gly Gly Thr Lys Val Glu Ile Lys Arg Thr Val Ala Ala Pro
                              105
Ser Val Phe Ile Phe Pro Pro Ser Asp Glu Gln Leu Lys Ser Gly Thr
                          120
                                            125
Ala Ser Val Val Cys Leu Leu Asn Asn Phe Tyr Pro Arg Glu Ala Lys
                      135
                                        140
Val Gln Trp Lys Val Asp Asn Ala Leu Gln Ser Gly Asn Ser Gln Glu
                  150
                                      155
Ser Val Thr Glu Gln Asp Ser Lys Asp Ser Thr Tyr Ser Leu Ser Ser
                                 170
Thr Leu Thr Leu Ser Lys Ala Asp Tyr Glu Lys His Lys Val Tyr Ala
                             185
Cys Glu Val Thr His Gln Gly Leu Ser Ser Pro Val Thr Lys Ser Phe
      195
                          200
Asn Arg Gly Glu Cys
   210
<210> 220
<211> 450
<212> PRT
<213> Artificial Sequence
<220>
<223> Humanized antibody - VH Chain
Gln Val Thr Leu Arg Glu Ser Gly Pro Ala Leu Val Lys Pro Thr Gln
```

```
Thr Leu Thr Leu Thr Cys Thr Phe Ser Gly Phe Ser Leu Ser Thr Ala
                          25
Gly Met Ser Val Gly Trp Ile Arg Gln Pro Pro Gly Lys Ala Leu Glu
Trp Leu Ala Asp Ile Trp Trp Asp Gly Lys Lys Asp Tyr Asn Pro Ser
                 55
Leu Lys Asp Arg Leu Thr Ile Ser Lys Asp Thr Ser Lys Asn Gln Val
              70
                                75
Val Leu Lys Val Thr Asn Met Asp Pro Ala Asp Thr Ala Thr Tyr Tyr
                           90
Cys Ala Arg Asp Met Ile Phe Asn Phe Tyr Phe Asp Val Trp Gly Gln
                       105
                                        110
Gly Thr Thr Val Thr Val Ser Ser Ala Ser Thr Lys Gly Pro Ser Val
                      120 125
Phe Pro Leu Ala Pro Ser Ser Lys Ser Thr Ser Gly Gly Thr Ala Ala
  130 135
                        140
Leu Gly Cys Leu Val Lys Asp Tyr Phe Pro Glu Pro Val Thr Val Ser
    150
                       155 160
Trp Asn Ser Gly Ala Leu Thr Ser Gly Val His Thr Phe Pro Ala Val
            165 170 175
Leu Gln Ser Ser Gly Leu Tyr Ser Leu Ser Ser Val Val Thr Val Pro
         180
                          185
Ser Ser Ser Leu Gly Thr Gln Thr Tyr Ile Cys Asn Val Asn His Lys
                      200
Pro Ser Asn Thr Lys Val Asp Lys Arg Val Glu Pro Lys Ser Cys Asp
   210 215
                                    220
Lys Thr His Thr Cys Pro Pro Cys Pro Ala Pro Glu Leu Leu Gly Gly
225 230
                                235
Pro Ser Val Phe Leu Phe Pro Pro Lys Pro Lys Asp Thr Leu Met Ile
             245
                             250 255
Ser Arg Thr Pro Glu Val Thr Cys Val Val Val Asp Val Ser His Glu
         260 265
Asp Pro Glu Val Lys Phe Asn Trp Tyr Val Asp Gly Val Glu Val His
 275 280
Asn Ala Lys Thr Lys Pro Arg Glu Glu Gln Tyr Asn Ser Thr Tyr Arg
   290 295
Val Val Ser Val Leu Thr Val Leu His Gln Asp Trp Leu Asn Gly Lys
305 310
                    315
Glu Tyr Lys Cys Lys Val Ser Asn Lys Ala Leu Pro Ala Pro Ile Glu
             325
                              330 335
Lys Thr Ile Ser Lys Ala Lys Gly Gln Pro Arg Glu Pro Gln Val Tyr
         340
                          345
Thr Leu Pro Pro Ser Arg Glu Glu Met Thr Lys Asn Gln Val Ser Leu
                       360
Thr Cys Leu Val Lys Gly Phe Tyr Pro Ser Asp Ile Ala Val Glu Trp
                   375
Glu Ser Asn Gly Gln Pro Glu Asn Asn Tyr Lys Thr Thr Pro Pro Val
385 390
                                 395
Leu Asp Ser Asp Gly Ser Phe Phe Leu Tyr Ser Lys Leu Thr Val Asp
                 410
            405
Lys Ser Arg Trp Gln Gln Gly Asn Val Phe Ser Cys Ser Val Met His
         420
                       425
Glu Ala Leu His Asn His Tyr Thr Gln Lys Ser Leu Ser Leu Ser Pro
                      440
Gly Lys
   450
<210> 221
<211> 213
<212> PRT
<213> Artificial Sequence
```

```
<220>
<223> Humanized antibody - VL Chain
Asp Ile Gln Met Thr Gln Ser Pro Ser Thr Leu Ser Ala Ser Val Gly
                                    10
Asp Arg Val Thr Ile Thr Cys Ser Ala Ser Ser Arg Val Gly Tyr Met
His Trp Tyr Gln Gln Lys Pro Gly Lys Ala Pro Lys Leu Leu Ile Tyr
                           40
Asp Thr Phe Lys Leu Ser Ser Gly Val Pro Ser Arg Phe Ser Gly Ser
Gly Ser Gly Thr Glu Phe Thr Leu Thr Ile Ser Ser Leu Gln Pro Asp
Asp Phe Ala Thr Tyr Tyr Cys Phe Gln Gly Ser Gly Tyr Pro Phe Thr
                85
                                    90
Phe Gly Gly Gly Thr Lys Val Glu Ile Lys Arg Thr Val Ala Ala Pro
            100
                                105
Ser Val Phe Ile Phe Pro Pro Ser Asp Glu Gln Leu Lys Ser Gly Thr
                           120
Ala Ser Val Val Cys Leu Leu Asn Asn Phe Tyr Pro Arg Glu Ala Lys
                        135
Val Gln Trp Lys Val Asp Asn Ala Leu Gln Ser Gly Asn Ser Gln Glu
                    150
Ser Val Thr Glu Gln Asp Ser Lys Asp Ser Thr Tyr Ser Leu Ser Ser
                165
                                    170
Thr Leu Thr Leu Ser Lys Ala Asp Tyr Glu Lys His Lys Val Tyr Ala
                               185
Cys Glu Val Thr His Gln Gly Leu Ser Ser Pro Val Thr Lys Ser Phe
                            200
Asn Arg Gly Glu Cys
   210
<210> 222
<211> 450
<212> PRT
<213> Artificial Sequence
<220>
<223> Humanized antibody - VH Chain
Gln Val Thr Leu Arg Glu Ser Gly Pro Ala Leu Val Lys Pro Thr Gln
Thr Leu Thr Leu Thr Cys Thr Phe Ser Gly Phe Ser Leu Ser Thr Ala
Gly Met Ser Val Gly Trp Ile Arg Gln Pro Pro Gly Lys Ala Leu Glu
Trp Leu Ala Asp Ile Trp Trp Asp Gly Lys Lys Ser Tyr Asn Pro Ser
                        55
Leu Lys Asp Arg Leu Thr Ile Ser Lys Asp Thr Ser Lys Asn Gln Val
                                        75
Val Leu Lys Val Thr Asn Met Asp Pro Ala Asp Thr Ala Thr Tyr Tyr
Cys Ala Arg Asp Met Ile Phe Asn Phe Tyr Phe Asp Val Trp Gly Gln
                               105
Gly Thr Thr Val Thr Val Ser Ser Ala Ser Thr Lys Gly Pro Ser Val
                          120
                                              125
Phe Pro Leu Ala Pro Ser Ser Lys Ser Thr Ser Gly Gly Thr Ala Ala
                        135
```

```
Leu Gly Cys Leu Val Lys Asp Tyr Phe Pro Glu Pro Val Thr Val Ser
                150
                                 155
Trp Asn Ser Gly Ala Leu Thr Ser Gly Val His Thr Phe Pro Ala Val
                             170
Leu Gln Ser Ser Gly Leu Tyr Ser Leu Ser Ser Val Val Thr Val Pro
                          185
Ser Ser Ser Leu Gly Thr Gln Thr Tyr Ile Cys Asn Val Asn His Lys
                       200
Pro Ser Asn Thr Lys Val Asp Lys Arg Val Glu Pro Lys Ser Cys Asp
         215
Lys Thr His Thr Cys Pro Pro Cys Pro Ala Pro Glu Leu Leu Gly Gly
      230 235 240
Pro Ser Val Phe Leu Phe Pro Pro Lys Pro Lys Asp Thr Leu Met Ile
            245 250
Ser Arg Thr Pro Glu Val Thr Cys Val Val Val Asp Val Ser His Glu
         260 265 270
Asp Pro Glu Val Lys Phe Asn Trp Tyr Val Asp Gly Val Glu Val His
          280
Asn Ala Lys Thr Lys Pro Arg Glu Glu Gln Tyr Asn Ser Thr Tyr Arg
                   295
Val Val Ser Val Leu Thr Val Leu His Gln Asp Trp Leu Asn Gly Lys
      310 315
Glu Tyr Lys Cys Lys Val Ser Asn Lys Ala Leu Pro Ala Pro Ile Glu
            325
                 330 335
Lys Thr Ile Ser Lys Ala Lys Gly Gln Pro Arg Glu Pro Gln Val Tyr
                         345 350
Thr Leu Pro Pro Ser Arg Glu Glu Met Thr Lys Asn Gln Val Ser Leu
                      360 365
Thr Cys Leu Val Lys Gly Phe Tyr Pro Ser Asp Ile Ala Val Glu Trp
                   375 380
Glu Ser Asn Gly Gln Pro Glu Asn Asn Tyr Lys Thr Thr Pro Pro Val
      390 395 400
Leu Asp Ser Asp Gly Ser Phe Phe Leu Tyr Ser Lys Leu Thr Val Asp
            405 410 415
Lys Ser Arg Trp Gln Gln Gly Asn Val Phe Ser Cys Ser Val Met His
        420 425 430
Glu Ala Leu His Asn His Tyr Thr Gln Lys Ser Leu Ser Leu Ser Pro
                      440
Gly Lys
  450
<210> 223
<211> 213
<212> PRT
<213> Artificial Sequence
<220>
<223> Humanized antibody - VL Chain
<400> 223
Asp Ile Gln Met Thr Gln Ser Pro Ser Thr Leu Ser Ala Ser Val Gly
   5
                          10
Asp Arg Val Thr Ile Thr Cys Ser Leu Ser Ser Arg Val Gly Tyr Met
His Trp Tyr Gln Gln Lys Pro Gly Lys Ala Pro Lys Leu Leu Ile Tyr
                      40
Asp Thr Met Tyr Gln Ser Ser Gly Val Pro Ser Arg Phe Ser Gly Ser
Gly Ser Gly Thr Glu Phe Thr Leu Thr Ile Ser Ser Leu Gln Pro Asp
                                75
Asp Phe Ala Thr Tyr Tyr Cys Phe Gln Gly Ser Gly Tyr Pro Phe Thr
```

```
90
            85
Phe Gly Gly Gly Thr Lys Val Glu Ile Lys Arg Thr Val Ala Ala Pro
        100 105
Ser Val Phe Ile Phe Pro Pro Ser Asp Glu Gln Leu Lys Ser Gly Thr
115 120 125
Ala Ser Val Val Cys Leu Leu Asn Asn Phe Tyr Pro Arg Glu Ala Lys
  130 135 140
Val Gln Trp Lys Val Asp Asn Ala Leu Gln Ser Gly Asn Ser Gln Glu
      150 155
Ser Val Thr Glu Gln Asp Ser Lys Asp Ser Thr Tyr Ser Leu Ser Ser
      165 170 175
Thr Leu Thr Leu Ser Lys Ala Asp Tyr Glu Lys His Lys Val Tyr Ala
        180 185 190
Cys Glu Val Thr His Gln Gly Leu Ser Ser Pro Val Thr Lys Ser Phe
                     200
Asn Arg Gly Glu Cys
<210> 224
<211> 450
<212> PRT
<213> Artificial Sequence
<220>
<223> Humanized antibody - VH Chain
<400> 224
Gln Val Thr Leu Arg Glu Ser Gly Pro Ala Leu Val Lys Pro Thr Gln
Thr Leu Thr Leu Thr Cys Thr Phe Ser Gly Phe Ser Leu Ser Thr Ala
Gly Met Ser Val Gly Trp Ile Arg Gln Pro Pro Gly Lys Ala Leu Glu
                      40
Trp Leu Ala Asp Ile Trp Trp Asp Asp Lys Lys Ser Tyr Asn Pro Ser
Leu Lys Asp Arg Leu Thr Ile Ser Lys Asp Thr Ser Lys Asn Gln Val
                    75
Val Leu Lys Val Thr Asn Met Asp Pro Ala Asp Thr Ala Thr Tyr Tyr
                            90 95
Cys Ala Arg Asp Met Ile Phe Asn Phe Tyr Phe Asp Val Trp Gly Gln
   100 105 110
Gly Thr Thr Val Thr Val Ser Ser Ala Ser Thr Lys Gly Pro Ser Val
                      120 125
Phe Pro Leu Ala Pro Ser Ser Lys Ser Thr Ser Gly Gly Thr Ala Ala
                   135
Leu Gly Cys Leu Val Lys Asp Tyr Phe Pro Glu Pro Val Thr Val Ser
               150
Trp Asn Ser Gly Ala Leu Thr Ser Gly Val His Thr Phe Pro Ala Val
            165
                            170 175
Leu Gln Ser Ser Gly Leu Tyr Ser Leu Ser Ser Val Val Thr Val Pro
                          185
         180
Ser Ser Ser Leu Gly Thr Gln Thr Tyr Ile Cys Asn Val Asn His Lys
                      200
Pro Ser Asn Thr Lys Val Asp Lys Arg Val Glu Pro Lys Ser Cys Asp
                   215
                                   220
Lys Thr His Thr Cys Pro Pro Cys Pro Ala Pro Glu Leu Leu Gly Gly
               230 235 240
Pro Ser Val Phe Leu Phe Pro Pro Lys Pro Lys Asp Thr Leu Met Ile
                             250
       245
Ser Arg Thr Pro Glu Val Thr Cys Val Val Val Asp Val Ser His Glu
```

```
265
          260
Asp Pro Glu Val Lys Phe Asn Trp Tyr Val Asp Gly Val Glu Val His
              280
Asn Ala Lys Thr Lys Pro Arg Glu Glu Gln Tyr Asn Ser Thr Tyr Arg
           295
Val Val Ser Val Leu Thr Val Leu His Gln Asp Trp Leu Asn Gly Lys
                       315
       310
Glu Tyr Lys Cys Lys Val Ser Asn Lys Ala Leu Pro Ala Pro Ile Glu
                     330 335
Lys Thr Ile Ser Lys Ala Lys Gly Gln Pro Arg Glu Pro Gln Val Tyr
                          345
Thr Leu Pro Pro Ser Arg Glu Glu Met Thr Lys Asn Gln Val Ser Leu
                       360
Thr Cys Leu Val Lys Gly Phe Tyr Pro Ser Asp Ile Ala Val Glu Trp
                   375
                           380
Glu Ser Asn Gly Gln Pro Glu Asn Asn Tyr Lys Thr Thr Pro Pro Val
                       395 400
                390
Leu Asp Ser Asp Gly Ser Phe Phe Leu Tyr Ser Lys Leu Thr Val Asp
                     410 415
Lys Ser Arg Trp Gln Gln Gly Asn Val Phe Ser Cys Ser Val Met His
                  425 430
Glu Ala Leu His Asn His Tyr Thr Gln Lys Ser Leu Ser Leu Ser Pro
                       440
Gly Lys
  450
<210> 225
<211> 213
<212> PRT
<213> Artificial Sequence
<220>
<223> Humanized antibody - VL Chain
<400> 225
Asp Ile Gln Met Thr Gln Ser Pro Ser Thr Leu Ser Ala Ser Val Gly
                               10
Asp Arg Val Thr Ile Thr Cys Leu Pro Ser Ser Arg Val Gly Tyr Met
His Trp Tyr Gln Gln Lys Pro Gly Lys Ala Pro Lys Leu Leu Ile Tyr
Asp Thr Met Tyr Gln Ser Ser Gly Val Pro Ser Arg Phe Ser Gly Ser
                     55
Gly Ser Gly Thr Glu Phe Thr Leu Thr Ile Ser Ser Leu Gln Pro Asp
                 70
Asp Phe Ala Thr Tyr Tyr Cys Phe Gln Gly Ser Gly Tyr Pro Phe Thr
Phe Gly Gly Gly Thr Lys Val Glu Ile Lys Arg Thr Val Ala Ala Pro
                            105
Ser Val Phe Ile Phe Pro Pro Ser Asp Glu Gln Leu Lys Ser Gly Thr
                        120
Ala Ser Val Val Cys Leu Leu Asn Asn Phe Tyr Pro Arg Glu Ala Lys
                     135
Val Gln Trp Lys Val Asp Asn Ala Leu Gln Ser Gly Asn Ser Gln Glu
                 150
                                   155 160
Ser Val Thr Glu Gln Asp Ser Lys Asp Ser Thr Tyr Ser Leu Ser Ser
                               170
              165
 Thr Leu Thr Leu Ser Lys Ala Asp Tyr Glu Lys His Lys Val Tyr Ala
                           185 190
        180
 Cys Glu Val Thr His Gln Gly Leu Ser Ser Pro Val Thr Lys Ser Phe
                         200
```

Asn Arg Gly Glu Cys 210 <210> 226 <211> 450 <212> PRT <213> Artificial Sequence <220> <223> Humanized antibody - VH Chain <400> 226 Gln Val Thr Leu Arg Glu Ser Gly Pro Ala Leu Val Lys Pro Thr Gln Thr Leu Thr Leu Thr Cys Thr Phe Ser Gly Phe Ser Leu Ser Thr Ala 25 Gly Met Ser Val Gly Trp Ile Arg Gln Pro Pro Gly Lys Ala Leu Glu 40 Trp Leu Ala Asp Ile Trp Trp Asp Asp Lys Lys His Tyr Asn Pro Ser 50 55 Leu Lys Asp Arg Leu Thr Ile Ser Lys Asp Thr Ser Lys Asn Gln Val 65 70 75 80 Val Leu Lys Val Thr Asn Met Asp Pro Ala Asp Thr Ala Thr Tyr Tyr 90 95 Cys Ala Arg Asp Met Ile Phe Asn Phe Tyr Phe Asp Val Trp Gly Gln 100 105 110 Gly Thr Thr Val Thr Val Ser Ser Ala Ser Thr Lys Gly Pro Ser Val 115 120 125 Phe Pro Leu Ala Pro Ser Ser Lys Ser Thr Ser Gly Gly Thr Ala Ala 130 135 140 Leu Gly Cys Leu Val Lys Asp Tyr Phe Pro Glu Pro Val Thr Val Ser 145 150 155 160 Trp Asn Ser Gly Ala Leu Thr Ser Gly Val His Thr Phe Pro Ala Val 165 170 175 Leu Gln Ser Ser Gly Leu Tyr Ser Leu Ser Ser Val Val Thr Val Pro 180 185 190 Ser Ser Ser Leu Gly Thr Gln Thr Tyr Ile Cys Asn Val Asn His Lys 195 200 205 Pro Ser Asn Thr Lys Val Asp Lys Arg Val Glu Pro Lys Ser Cys Asp 210 215 220 Lys Thr His Thr Cys Pro Pro Cys Pro Ala Pro Glu Leu Leu Gly Gly 225 230 235 240 Pro Ser Val Phe Leu Phe Pro Pro Lys Pro Lys Asp Thr Leu Met Ile 245 250 255 Ser Arg Thr Pro Glu Val Thr Cys Val Val Val Asp Val Ser His Glu 260 265 270 Asp Pro Glu Val Lys Phe Asn Trp Tyr Val Asp Gly Val Glu Val His 275 280 285 Asn Ala Lys Thr Lys Pro Arg Glu Glu Gln Tyr Asn Ser Thr Tyr Arg 290 295 300 Val Val Ser Val Leu Thr Val Leu His Gln Asp Trp Leu Asn Gly Lys 305 310 315 320 Glu Tyr Lys Cys Lys Val Ser Asn Lys Ala Leu Pro Ala Pro Ile Glu 330 Lys Thr Ile Ser Lys Ala Lys Gly Gln Pro Arg Glu Pro Gln Val Tyr 345 Thr Leu Pro Pro Ser Arg Glu Glu Met Thr Lys Asn Gln Val Ser Leu 360 Thr Cys Leu Val Lys Gly Phe Tyr Pro Ser Asp Ile Ala Val Glu Trp 375 380 Glu Ser Asn Gly Gln Pro Glu Asn Asn Tyr Lys Thr Thr Pro Pro Val

```
390
                                    395
385
Leu Asp Ser Asp Gly Ser Phe Phe Leu Tyr Ser Lys Leu Thr Val Asp
                       410
Lys Ser Arg Trp Gln Gln Gly Asn Val Phe Ser Cys Ser Val Met His
                  425 430
Glu Ala Leu His Asn His Tyr Thr Gln Lys Ser Leu Ser Leu Ser Pro
                        440
Gly Lys
  450
<210> 227
<211> 213
<212> PRT
<213> Artificial Sequence
<220>
<223> Humanized antibody - VL Chain
<400> 227
Asp Ile Gln Met Thr Gln Ser Pro Ser Thr Leu Ser Ala Ser Val Gly
                                 10
Asp Arg Val Thr Ile Thr Cys Ser Ala Ser Ser Arg Val Gly Tyr Met
His Trp Tyr Gln Gln Lys Pro Gly Lys Ala Pro Lys Leu Leu Ile Tyr
                         4.0
Asp Thr Phe Phe Leu Asp Ser Gly Val Pro Ser Arg Phe Ser Gly Ser
                      55
Gly Ser Gly Thr Glu Phe Thr Leu Thr Ile Ser Ser Leu Gln Pro Asp
Asp Phe Ala Thr Tyr Tyr Cys Phe Gln Gly Ser Gly Tyr Pro Phe Thr
Phe Gly Gly Gly Thr Lys Val Glu Ile Lys Arg Thr Val Ala Ala Pro
                             105
          100
Ser Val Phe Ile Phe Pro Pro Ser Asp Glu Gln Leu Lys Ser Gly Thr
                         120
Ala Ser Val Val Cys Leu Leu Asn Asn Phe Tyr Pro Arg Glu Ala Lys
                                        140
                     135
Val Gln Trp Lys Val Asp Asn Ala Leu Gln Ser Gly Asn Ser Gln Glu
                                  '155
                  150
Ser Val Thr Glu Gln Asp Ser Lys Asp Ser Thr Tyr Ser Leu Ser Ser
              165
                                170
Thr Leu Thr Leu Ser Lys Ala Asp Tyr Glu Lys His Lys Val Tyr Ala
          180 185 190
Cys Glu Val Thr His Gln Gly Leu Ser Ser Pro Val Thr Lys Ser Phe
                         200
      195
Asn Arg Gly Glu Cys
   210
<210> 228
<211> 450
<212> PRT
<213> Artificial Sequence
<220>
<223> Humanized antibody - VH Chain
Gln Val Thr Leu Arg Glu Ser Gly Pro Ala Leu Val Lys Pro Thr Gln
                                 10
Thr Leu Thr Leu Thr Cys Thr Phe Ser Gly Phe Ser Leu Ser Thr Ala
```

```
Gly Met Ser Val Gly Trp Ile Arg Gln Pro Pro Gly Lys Ala Leu Glu
         40 45
Trp Leu Ala Asp Ile Trp Trp Asp Asp Lys Lys Ser Tyr Asn Pro Ser
50 55
Leu Lys Asp Arg Leu Thr Ile Ser Lys Asp Thr Ser Lys Asn Gln Val
   70 75 80
Val Leu Lys Val Thr Asn Met Asp Pro Ala Asp Thr Ala Thr Tyr Tyr
          85 90 95
Cys Ala Arg Asp Met Ile Phe Asn Trp Tyr Phe Asp Val Trp Gly Gln
 100 105 110
Gly Thr Thr Val Thr Val Ser Ser Ala Ser Thr Lys Gly Pro Ser Val
 115 120 125
Phe Pro Leu Ala Pro Ser Ser Lys Ser Thr Ser Gly Gly Thr Ala Ala
130 135 140
Leu Gly Cys Leu Val Lys Asp Tyr Phe Pro Glu Pro Val Thr Val Ser
145 150 155 160
Trp Asn Ser Gly Ala Leu Thr Ser Gly Val His Thr Phe Pro Ala Val
   165 170 175
Leu Gln Ser Ser Gly Leu Tyr Ser Leu Ser Ser Val Val Thr Val Pro
 180 185 190
Ser Ser Ser Leu Gly Thr Gln Thr Tyr Ile Cys Asn Val Asn His Lys
 195 200 205
Pro Ser Asn Thr Lys Val Asp Lys Arg Val Glu Pro Lys Ser Cys Asp
 210 215 220
Lys Thr His Thr Cys Pro Pro Cys Pro Ala Pro Glu Leu Leu Gly Gly
225 230 235 240
Pro Ser Val Phe Leu Phe Pro Pro Lys Pro Lys Asp Thr Leu Met Ile
                        250 255
Ser Arg Thr Pro Glu Val Thr Cys Val Val Val Asp Val Ser His Glu
 260 265 270
Asp Pro Glu Val Lys Phe Asn Trp Tyr Val Asp Gly Val Glu Val His
 275 280
Asn Ala Lys Thr Lys Pro Arg Glu Glu Gln Tyr Asn Ser Thr Tyr Arg
 290 295 300
Val Val Ser Val Leu Thr Val Leu His Gln Asp Trp Leu Asn Gly Lys
                          315 320
305 310
Glu Tyr Lys Cys Lys Val Ser Asn Lys Ala Leu Pro Ala Pro Ile Glu
          325 330 335
Lys Thr Ile Ser Lys Ala Lys Gly Gln Pro Arg Glu Pro Gln Val Tyr
       340 345 350
Thr Leu Pro Pro Ser Arg Glu Glu Met Thr Lys Asn Gln Val Ser Leu
                  360 365
Thr Cys Leu Val Lys Gly Phe Tyr Pro Ser Asp Ile Ala Val Glu Trp
 370 375 380
Glu Ser Asn Gly Gln Pro Glu Asn Asn Tyr Lys Thr Thr Pro Pro Val
385 390 395 400
Leu Asp Ser Asp Gly Ser Phe Phe Leu Tyr Ser Lys Leu Thr Val Asp
   405 410 415
Lys Ser Arg Trp Gln Gln Gly Asn Val Phe Ser Cys Ser Val Met His
   420 425 430
Glu Ala Leu His Asn His Tyr Thr Gln Lys Ser Leu Ser Leu Ser Pro
                 440
Gly Lys
  450
<210> 229
<211> 213
<212> PRT
<213> Artificial Sequence
<220>
```

73

PCT/US2006/024717

WO 2007/002543 <223> Humanized antibody - VL Chain <400> 229 Asp Ile Gln Met Thr Gln Ser Pro Ser Thr Leu Ser Ala Ser Val Gly 10 Asp Arg Val Thr Ile Thr Cys Ser Pro Ser Ser Arg Val Gly Tyr Met His Trp Tyr Gln Gln Lys Pro Gly Lys Ala Pro Lys Leu Leu Ile Tyr 40 Asp Thr Arg Tyr Gln Ser Ser Gly Val Pro Ser Arg Phe Ser Gly Ser 55 Gly Ser Gly Thr Glu Phe Thr Leu Thr Ile Ser Ser Leu Gln Pro Asp 70 Asp Phe Ala Thr Tyr Tyr Cys Phe Gln Gly Ser Gly Tyr Pro Phe Thr 90 Phe Gly Gly Gly Thr Lys Val Glu Ile Lys Arg Thr Val Ala Ala Pro 100 105 Ser Val Phe Ile Phe Pro Pro Ser Asp Glu Gln Leu Lys Ser Gly Thr 120 Ala Ser Val Val Cys Leu Leu Asn Asn Phe Tyr Pro Arg Glu Ala Lys 135 150 165 170

Val Gln Trp Lys Val Asp Asn Ala Leu Gln Ser Gly Asn Ser Gln Glu Ser Val Thr Glu Gln Asp Ser Lys Asp Ser Thr Tyr Ser Leu Ser Ser Thr Leu Thr Leu Ser Lys Ala Asp Tyr Glu Lys His Lys Val Tyr Ala

185 Cys Glu Val Thr His Gln Gly Leu Ser Ser Pro Val Thr Lys Ser Phe 195 200

Asn Arg Gly Glu Cys 210

<210> 230

<211> 450

<212> PRT

<213> Artificial Sequence

<220>

<223> Humanized antibody - VH Chain

<400> 230

Gln Val Thr Leu Arg Glu Ser Gly Pro Ala Leu Val Lys Pro Thr Gln Thr Leu Thr Leu Thr Cys Thr Phe Ser Gly Phe Ser Leu Ser Thr Ser Gly Met Ser Val Gly Trp Ile Arg Gln Pro Pro Gly Lys Ala Leu Glu Trp Leu Ala Asp Ile Trp Trp Asp Asp Lys Lys Asp Tyr Asn Pro Ser Leu Lys Ser Arg Leu Thr Ile Ser Lys Asp Thr Ser Lys Asn Gln Val 75 Val Leu Lys Val Thr Asn Met Asp Pro Ala Asp Thr Ala Thr Tyr Tyr Cys Ala Arg Ser Met Ile Thr Asn Trp Tyr Phe Asp Val Trp Gly Gln 105 Gly Thr Thr Val Thr Val Ser Ser Ala Ser Thr Lys Gly Pro Ser Val 120 Phe Pro Leu Ala Pro Ser Ser Lys Ser Thr Ser Gly Gly Thr Ala Ala 135 140 Leu Gly Cys Leu Val Lys Asp Tyr Phe Pro Glu Pro Val Thr Val Ser 155

74

```
Trp Asn Ser Gly Ala Leu Thr Ser Gly Val His Thr Phe Pro Ala Val
           165 170 175
Leu Gln Ser Ser Gly Leu Tyr Ser Leu Ser Ser Val Val Thr Val Pro
                185
Ser Ser Ser Leu Gly Thr Gln Thr Tyr Ile Cys Asn Val Asn His Lys
           200
Pro Ser Asn Thr Lys Val Asp Lys Arg Val Glu Pro Lys Ser Cys Asp
 210 215 220
Lys Thr His Thr Cys Pro Pro Cys Pro Ala Pro Glu Leu Leu Gly Gly
             230 235 240
Pro Ser Val Phe Leu Phe Pro Pro Lys Pro Lys Asp Thr Leu Met Ile
                            250 255
Ser Arg Thr Pro Glu Val Thr Cys Val Val Val Asp Val Ser His Glu
Asp Pro Glu Val Lys Phe Asn Trp Tyr Val Asp Gly Val Glu Val His
                     280 285
Asn Ala Lys Thr Lys Pro Arg Glu Glu Gln Tyr Asn Ser Thr Tyr Arg
       295 300
Val Val Ser Val Leu Thr Val Leu His Gln Asp Trp Leu Asn Gly Lys
   310 315 320
Glu Tyr Lys Cys Lys Val Ser Asn Lys Ala Leu Pro Ala Pro Ile Glu
            325 330 335
Lys Thr Ile Ser Lys Ala Lys Gly Gln Pro Arg Glu Pro Gln Val Tyr
                         345 350
Thr Leu Pro Pro Ser Arg Glu Glu Met Thr Lys Asn Gln Val Ser Leu
                      360
Thr Cys Leu Val Lys Gly Phe Tyr Pro Ser Asp Ile Ala Val Glu Trp
                375 380
Glu Ser Asn Gly Gln Pro Glu Asn Asn Tyr Lys Thr Thr Pro Pro Val
385 390 395 400
Leu Asp Ser Asp Gly Ser Phe Phe Leu Tyr Ser Lys Leu Thr Val Asp
            405 410 415
Lys Ser Arg Trp Gln Gln Gly Asn Val Phe Ser Cys Ser Val Met His
   420 425 430
Glu Ala Leu His Asn His Tyr Thr Gln Lys Ser Leu Ser Leu Ser Pro
                     440
Gly Lys
  450
<210> 231
<211> 213
<212> PRT
<213> Artificial Sequence
<223> Humanized antibody - VL Chain
<400> 231
Asp Ile Gln Met Thr Gln Ser Pro Ser Thr Leu Ser Ala Ser Val Gly
                             10
Asp Arg Val Thr Ile Thr Cys Ser Ala Ser Ser Ser Val Gly Tyr Met
                          25
          20
His Trp Tyr Gln Gln Lys Pro Gly Lys Ala Pro Lys Leu Leu Ile Tyr
                       40
Asp Thr Ser Lys Leu Ala Ser Gly Val Pro Ser Arg Phe Ser Gly Ser
                   55
Gly Ser Gly Thr Glu Phe Thr Leu Thr Ile Ser Ser Leu Gln Pro Asp
                70
                                75
Asp Phe Ala Thr Tyr Tyr Cys Phe Gln Gly Ser Gly Tyr Pro Phe Thr
Phe Gly Gly Gly Thr Lys Val Glu Ile Lys Arg Thr Val Ala Ala Pro
```

```
105
          100
Ser Val Phe Ile Phe Pro Pro Ser Asp Glu Gln Leu Lys Ser Gly Thr
     115 120
Ala Ser Val Val Cys Leu Leu Asn Asn Phe Tyr Pro Arg Glu Ala Lys
         135
Val Gln Trp Lys Val Asp Asn Ala Leu Gln Ser Gly Asn Ser Gln Glu
                       155
        150
Ser Val Thr Glu Gln Asp Ser Lys Asp Ser Thr Tyr Ser Leu Ser Ser
                   170
             165
Thr Leu Thr Leu Ser Lys Ala Asp Tyr Glu Lys His Lys Val Tyr Ala
                  185
Cys Glu Val Thr His Gln Gly Leu Ser Ser Pro Val Thr Lys Ser Phe
                        200
Asn Arg Gly Glu Cys
<210> 232
<211> 450
<212> PRT
<213> Artificial Sequence
<220>
<223> Humanized antibody - VH Chain
<400> 232
Gln Val Thr Leu Arg Glu Ser Gly Pro Ala Leu Val Lys Pro Thr Gln
                                1.0
Thr Leu Thr Leu Thr Cys Thr Phe Ser Gly Phe Ser Leu Ser Thr Ala
                            25
Gly Met Ser Val Gly Trp Ile Arg Gln Pro Pro Gly Lys Ala Leu Glu
                         40
Trp Leu Ala Asp Ile Trp Trp Asp Asp Lys Lys Asp Tyr Asn Pro Ser
Leu Lys Ser Arg Leu Thr Ile Ser Lys Asp Thr Ser Lys Asn Gln Val
Val Leu Lys Val Thr Asn Met Asp Pro Ala Asp Thr Ala Thr Tyr Tyr
                                90
Cys Ala Arg Asp Met Ile Phe Asn Trp Tyr Phe Asp Val Trp Gly Gln
                            105
Gly Thr Thr Val Thr Val Ser Ser Ala Ser Thr Lys Gly Pro Ser Val
                         120
Phe Pro Leu Ala Pro Ser Ser Lys Ser Thr Ser Gly Gly Thr Ala Ala
                     135
                             140
Leu Gly Cys Leu Val Lys Asp Tyr Phe Pro Glu Pro Val Thr Val Ser
                 150 155 160
Trp Asn Ser Gly Ala Leu Thr Ser Gly Val His Thr Phe Pro Ala Val
                                170
Leu Gln Ser Ser Gly Leu Tyr Ser Leu Ser Ser Val Val Thr Val Pro
                             185
Ser Ser Ser Leu Gly Thr Gln Thr Tyr Ile Cys Asn Val Asn His Lys
                         200
Pro Ser Asn Thr Lys Val Asp Lys Arg Val Glu Pro Lys Ser Cys Asp
                     215
Lys Thr His Thr Cys Pro Pro Cys Pro Ala Pro Glu Leu Leu Gly Gly
                 230
                                    235
Pro Ser Val Phe Leu Phe Pro Pro Lys Pro Lys Asp Thr Leu Met Ile
              245
                                250
Ser Arg Thr Pro Glu Val Thr Cys Val Val Val Asp Val Ser His Glu
        260 265
Asp Pro Glu Val Lys Phe Asn Trp Tyr Val Asp Gly Val Glu Val His
                         280
```

```
Asn Ala Lys Thr Lys Pro Arg Glu Glu Gln Tyr Asn Ser Thr Tyr Arg
           295
Val Val Ser Val Leu Thr Val Leu His Gln Asp Trp Leu Asn Gly Lys
             310
                                315
Glu Tyr Lys Cys Lys Val Ser Asn Lys Ala Leu Pro Ala Pro Ile Glu
            325 330 335
Lys Thr Ile Ser Lys Ala Lys Gly Gln Pro Arg Glu Pro Gln Val Tyr
   340 345 350
Thr Leu Pro Pro Ser Arg Glu Glu Met Thr Lys Asn Gln Val Ser Leu
 355 360 365
Thr Cys Leu Val Lys Gly Phe Tyr Pro Ser Asp Ile Ala Val Glu Trp
370 375 380
Glu Ser Asn Gly Gln Pro Glu Asn Asn Tyr Lys Thr Thr Pro Pro Val
385 390 395 400
Leu Asp Ser Asp Gly Ser Phe Phe Leu Tyr Ser Lys Leu Thr Val Asp
           405 410 415
Lys Ser Arg Trp Gln Gln Gly Asn Val Phe Ser Cys Ser Val Met His
   420 425 430
Glu Ala Leu His Asn His Tyr Thr Gln Lys Ser Leu Ser Leu Ser Pro
            440
Gly Lys
  450
<210> 233
<211> 213
<212> PRT
<213> Artificial Sequence
<220>
<223> Humanized antibody - VL Chain
<400> 233
Asp Ile Gln Met Thr Gln Ser Pro Ser Thr Leu Ser Ala Ser Val Gly
                             10
Asp Arg Val Thr Ile Thr Cys Ser Ala Ser Ser Ser Val Gly Tyr Met
                         25
His Trp Tyr Gln Gln Lys Pro Gly Lys Ala Pro Lys Leu Leu Ile Tyr
                      40
Asp Thr Phe Lys Leu Ala Ser Gly Val Pro Ser Arg Phe Ser Gly Ser
                   55
Gly Ser Gly Thr Glu Phe Thr Leu Thr Ile Ser Ser Leu Gln Pro Asp
                                75
Asp Phe Ala Thr Tyr Cys Phe Gln Gly Ser Gly Tyr Pro Phe Thr
                             90
Phe Gly Gly Gly Thr Lys Val Glu Ile Lys Arg Thr Val Ala Ala Pro
                         105 110
Ser Val Phe Ile Phe Pro Pro Ser Asp Glu Gln Leu Lys Ser Gly Thr
                      120
Ala Ser Val Val Cys Leu Leu Asn Asn Phe Tyr Pro Arg Glu Ala Lys
                   135
Val Gln Trp Lys Val Asp Asn Ala Leu Gln Ser Gly Asn Ser Gln Glu
                                155 160
Ser Val Thr Glu Gln Asp Ser Lys Asp Ser Thr Tyr Ser Leu Ser Ser
                             170
            1.65
Thr Leu Thr Leu Ser Lys Ala Asp Tyr Glu Lys His Lys Val Tyr Ala
         180 185 190
Cys Glu Val Thr His Gln Gly Leu Ser Ser Pro Val Thr Lys Ser Phe
      195
                      200
Asn Arg Gly Glu Cys
   210
```

<210> 234

```
<211> 450
<212> PRT
<213> Artificial Sequence
<223> Humanized antibody - VH Chain
<400> 234
Gln Val Thr Leu Arg Glu Ser Gly Pro Ala Leu Val Lys Pro Thr Gln
               10
Thr Leu Thr Leu Thr Cys Thr Phe Ser Gly Phe Ser Leu Ser Thr Ala
              25
Gly Met Ser Val Gly Trp Ile Arg Gln Pro Pro Gly Lys Ala Leu Glu
         40
Trp Leu Ala Asp Ile Trp Trp Asp Asp Lys Lys Asp Tyr Asn Pro Ser
50 55
Leu Lys Ser Arg Leu Thr Ile Ser Lys Asp Thr Ser Lys Asn Gln Val
    70 75
Val Leu Lys Val Thr Asn Met Asp Pro Ala Asp Thr Ala Thr Tyr Tyr
               90 95
Cys Ala Arg Asp Met Ile Phe Asn Trp Tyr Phe Asp Val Trp Gly Gln
  100 105 110
Gly Thr Thr Val Thr Val Ser Ser Ala Ser Thr Lys Gly Pro Ser Val
115 120 125
Phe Pro Leu Ala Pro Ser Ser Lys Ser Thr Ser Gly Gly Thr Ala Ala
130 135 140
Leu Gly Cys Leu Val Lys Asp Tyr Phe Pro Glu Pro Val Thr Val Ser
    150 155 160
Trp Asn Ser Gly Ala Leu Thr Ser Gly Val His Thr Phe Pro Ala Val
     165 170 175
Leu Gln Ser Ser Gly Leu Tyr Ser Leu Ser Ser Val Val Thr Val Pro
 180 185 190
Ser Ser Ser Leu Gly Thr Gln Thr Tyr Ile Cys Asn Val Asn His Lys
195 200
Pro Ser Asn Thr Lys Val Asp Lys Arg Val Glu Pro Lys Ser Cys Asp
  210 215
                             220
Lys Thr His Thr Cys Pro Pro Cys Pro Ala Pro Glu Leu Leu Gly Gly
225 230 235 240
Pro Ser Val Phe Leu Phe Pro Pro Lys Pro Lys Asp Thr Leu Met Ile
  245 250 255
Ser Arg Thr Pro Glu Val Thr Cys Val Val Val Asp Val Ser His Glu
 260 265 270
Asp Pro Glu Val Lys Phe Asn Trp Tyr Val Asp Gly Val Glu Val His
275 280
Asn Ala Lys Thr Lys Pro Arg Glu Glu Gln Tyr Asn Ser Thr Tyr Arg
290 295 300
Val Val Ser Val Leu Thr Val Leu His Gln Asp Trp Leu Asn Gly Lys
305 310 315 320
Glu Tyr Lys Cys Lys Val Ser Asn Lys Ala Leu Pro Ala Pro Ile Glu
           325 330 335
Lys Thr Ile Ser Lys Ala Lys Gly Gln Pro Arg Glu Pro Gln Val Tyr
                      345 350
Thr Leu Pro Pro Ser Arg Glu Glu Met Thr Lys Asn Gln Val Ser Leu
                   360
Thr Cys Leu Val Lys Gly Phe Tyr Pro Ser Asp Ile Ala Val Glu Trp
                375
Glu Ser Asn Gly Gln Pro Glu Asn Asn Tyr Lys Thr Thr Pro Pro Val
             390 395 400
Leu Asp Ser Asp Gly Ser Phe Phe Leu Tyr Ser Lys Leu Thr Val Asp
                         410
Lys Ser Arg Trp Gln Gln Gly Asn Val Phe Ser Cys Ser Val Met His
```

```
420
                             425
Glu Ala Leu His Asn His Tyr Thr Gln Lys Ser Leu Ser Leu Ser Pro
Gly Lys
  450
<210> 235
<211> 213
<212> PRT
<213> Artificial Sequence
<220>
<223> Humanized antibody - VL Chain
<400> 235
Asp Ile Gln Met Thr Gln Ser Pro Ser Thr Leu Ser Ala Ser Val Gly
                             10
Asp Arg Val Thr Ile Thr Cys Ser Ala Ser Ser Ser Val Gly Tyr Met
                   25
His Trp Tyr Gln Gln Lys Pro Gly Lys Ala Pro Lys Leu Leu Ile Tyr
                        40
Asp Thr Tyr Lys Gln Thr Ser Gly Val Pro Ser Arg Phe Ser Gly Ser
                      55
Gly Ser Gly Thr Glu Phe Thr Leu Thr Ile Ser Ser Leu Gln Pro Asp
                 70
                                   75
Asp Phe Ala Thr Tyr Tyr Cys Phe Gln Gly Ser Gly Tyr Pro Phe Thr
             85
                                90
Phe Gly Gly Gly Thr Lys Val Glu Ile Lys Arg Thr Val Ala Ala Pro
         100 105
Ser Val Phe Ile Phe Pro Pro Ser Asp Glu Gln Leu Lys Ser Gly Thr
                        120
Ala Ser Val Val Cys Leu Leu Asn Asn Phe Tyr Pro Arg Glu Ala Lys
  130 135
                                       140
Val Gln Trp Lys Val Asp Asn Ala Leu Gln Ser Gly Asn Ser Gln Glu
       150 155
Ser Val Thr Glu Gln Asp Ser Lys Asp Ser Thr Tyr Ser Leu Ser Ser
             165
                                170
Thr Leu Thr Leu Ser Lys Ala Asp Tyr Glu Lys His Lys Val Tyr Ala
          180 185
Cys Glu Val Thr His Gln Gly Leu Ser Ser Pro Val Thr Lys Ser Phe
   195
                        200
Asn Arg Gly Glu Cys
  210
<210> 236
<211> 450
<212> PRT
<213> Artificial Sequence
<220>
<223> Humanized antibody - VH Chain
<400> 236
Gln Val Thr Leu Arg Glu Ser Gly Pro Ala Leu Val Lys Pro Thr Gln
Thr Leu Thr Leu Thr Cys Thr Phe Ser Gly Phe Ser Leu Ser Thr Ala
                             25
Gly Met Ser Val Gly Trp Ile Arg Gln Pro Pro Gly Lys Ala Leu Glu
                         40
Trp Leu Ala Asp Ile Trp Trp Asp Asp Lys Lys Asp Tyr Asn Pro Ser
```

```
55
Leu Lys Ser Arg Leu Thr Ile Ser Lys Asp Thr Ser Lys Asn Gln Val
                  70
                                    75
Val Leu Lys Val Thr Asn Met Asp Pro Ala Asp Thr Ala Thr Tyr Tyr
Cys Ala Arg Asp Met Ile Phe Asn Trp Tyr Phe Asp Val Trp Gly Gln
           100
                 105
Gly Thr Thr Val Thr Val Ser Ser Ala Ser Thr Lys Gly Pro Ser Val
                         120
Phe Pro Leu Ala Pro Ser Ser Lys Ser Thr Ser Gly Gly Thr Ala Ala
                      135
                              140
Leu Gly Cys Leu Val Lys Asp Tyr Phe Pro Glu Pro Val Thr Val Ser
                  150
                                   155
Trp Asn Ser Gly Ala Leu Thr Ser Gly Val His Thr Phe Pro Ala Val
              165
                                170 175
Leu Gln Ser Ser Gly Leu Tyr Ser Leu Ser Ser Val Val Thr Val Pro
           180
                             185
Ser Ser Ser Leu Gly Thr Gln Thr Tyr Ile Cys Asn Val Asn His Lys
                         200
Pro Ser Asn Thr Lys Val Asp Lys Arg Val Glu Pro Lys Ser Cys Asp
                      215
                             220
Lys Thr His Thr Cys Pro Pro Cys Pro Ala Pro Glu Leu Leu Gly Gly
                  230
                          235
Pro Ser Val Phe Leu Phe Pro Pro Lys Pro Lys Asp Thr Leu Met Ile
                                250
Ser Arg Thr Pro Glu Val Thr Cys Val Val Val Asp Val Ser His Glu
           260
                             265
Asp Pro Glu Val Lys Phe Asn Trp Tyr Val Asp Gly Val Glu Val His
                        280
Asn Ala Lys Thr Lys Pro Arg Glu Glu Gln Tyr Asn Ser Thr Tyr Arg
                     295
Val Val Ser Val Leu Thr Val Leu His Gln Asp Trp Leu Asn Gly Lys
                  310
                         315
Glu Tyr Lys Cys Lys Val Ser Asn Lys Ala Leu Pro Ala Pro Ile Glu
              325
                               330 335
Lys Thr Ile Ser Lys Ala Lys Gly Gln Pro Arg Glu Pro Gln Val Tyr
          340
                             345
Thr Leu Pro Pro Ser Arg Glu Glu Met Thr Lys Asn Gln Val Ser Leu
                         360
Thr Cys Leu Val Lys Gly Phe Tyr Pro Ser Asp Ile Ala Val Glu Trp
                     375
                                      380
Glu Ser Asn Gly Gln Pro Glu Asn Asn Tyr Lys Thr Thr Pro Pro Val
                      395 400
                  390
Leu Asp Ser Asp Gly Ser Phe Phe Leu Tyr Ser Lys Leu Thr Val Asp
                               410 415
Lys Ser Arg Trp Gln Gln Gly Asn Val Phe Ser Cys Ser Val Met His
                  425 430
Glu Ala Leu His Asn His Tyr Thr Gln Lys Ser Leu Ser Leu Ser Pro
                         440
Gly Lys
   450
<210> 237
<211> 213
<212> PRT
<213> Artificial Sequence
<220>
<223> Humanized antibody - VL Chain
<400> 237
```

```
Asp Ile Gln Met Thr Gln Ser Pro Ser Thr Leu Ser Ala Ser Val Gly
Asp Arg Val Thr Ile Thr Cys Ser Ala Ser Ser Ser Val Gly Tyr Met
                                25
His Trp Tyr Gln Gln Lys Pro Gly Lys Ala Pro Lys Leu Leu Ile Tyr
Asp Thr Arg Tyr Leu Ser Ser Gly Val Pro Ser Arg Phe Ser Gly Ser
Gly Ser Gly Thr Glu Phe Thr Leu Thr Ile Ser Ser Leu Gln Pro Asp
                                        75
Asp Phe Ala Thr Tyr Tyr Cys Phe Gln Gly Ser Gly Tyr Pro Phe Thr
                85
                                    90
Phe Gly Gly Gly Thr Lys Val Glu Ile Lys Arg Thr Val Ala Ala Pro
                                105
Ser Val Phe Ile Phe Pro Pro Ser Asp Glu Gln Leu Lys Ser Gly Thr
                            120
Ala Ser Val Val Cys Leu Leu Asn Asn Phe Tyr Pro Arg Glu Ala Lys
                       135
Val Gln Trp Lys Val Asp Asn Ala Leu Gln Ser Gly Asn Ser Gln Glu
                    150
                                        155
Ser Val Thr Glu Gln Asp Ser Lys Asp Ser Thr Tyr Ser Leu Ser Ser
                                    170
Thr Leu Thr Leu Ser Lys Ala Asp Tyr Glu Lys His Lys Val Tyr Ala
                                185
Cys Glu Val Thr His Gln Gly Leu Ser Ser Pro Val Thr Lys Ser Phe
                            200
Asn Arg Gly Glu Cys
  210
<210> 238
<211> 450
<212> PRT
<213> Artificial Sequence
<220>
<223> Humanized antibody - VH Chain
<400> 238
Gln Val Thr Leu Arg Glu Ser Gly Pro Ala Leu Val Lys Pro Thr Gln
                                    10
Thr Leu Thr Leu Thr Cys Thr Phe Ser Gly Phe Ser Leu Ser Thr Ala
Gly Met Ser Val Gly Trp Ile Arg Gln Pro Pro Gly Lys Ala Leu Glu
                           40
Trp Leu Ala Asp Ile Trp Trp Asp Asp Lys Lys Asp Tyr Asn Pro Ser
                       55
Leu Lys Ser Arg Leu Thr Ile Ser Lys Asp Thr Ser Lys Asn Gln Val
                   70
                                        75
Val Leu Lys Val Thr Asn Met Asp Pro Ala Asp Thr Ala Thr Tyr Tyr
               8.5
                                    90
Cys Ala Arg Asp Met Ile Thr Asn Phe Tyr Phe Asp Val Trp Gly Gln
            100
                                105
Gly Thr Thr Val Thr Val Ser Ser Ala Ser Thr Lys Gly Pro Ser Val
                           120
Phe Pro Leu Ala Pro Ser Ser Lys Ser Thr Ser Gly Gly Thr Ala Ala
                       135
                                           1.40
Leu Gly Cys Leu Val Lys Asp Tyr Phe Pro Glu Pro Val Thr Val Ser
                   150
                                        155
Trp Asn Ser Gly Ala Leu Thr Ser Gly Val His Thr Phe Pro Ala Val
                                   170
Leu Gln Ser Ser Gly Leu Tyr Ser Leu Ser Ser Val Val Thr Val Pro
```

```
180
                           185
Ser Ser Ser Leu Gly Thr Gln Thr Tyr Ile Cys Asn Val Asn His Lys
                 200
Pro Ser Asn Thr Lys Val Asp Lys Arg Val Glu Pro Lys Ser Cys Asp
                                     220
           215
Lys Thr His Thr Cys Pro Pro Cys Pro Ala Pro Glu Leu Leu Gly Gly
                         235
                230
Pro Ser Val Phe Leu Phe Pro Pro Lys Pro Lys Asp Thr Leu Met Ile
                               250
             245
Ser Arg Thr Pro Glu Val Thr Cys Val Val Val Asp Val Ser His Glu
       260 265
Asp Pro Glu Val Lys Phe Asn Trp Tyr Val Asp Gly Val Glu Val His
             280
Asn Ala Lys Thr Lys Pro Arg Glu Glu Gln Tyr Asn Ser Thr Tyr Arg
           295
                           300
Val Val Ser Val Leu Thr Val Leu His Gln Asp Trp Leu Asn Gly Lys
                310
                        315
Glu Tyr Lys Cys Lys Val Ser Asn Lys Ala Leu Pro Ala Pro Ile Glu
                    330 335
             325
Lys Thr Ile Ser Lys Ala Lys Gly Gln Pro Arg Glu Pro Gln Val Tyr
                          345
Thr Leu Pro Pro Ser Arg Glu Glu Met Thr Lys Asn Gln Val Ser Leu
                       360
Thr Cys Leu Val Lys Gly Phe Tyr Pro Ser Asp Ile Ala Val Glu Trp
                            380
                   375
Glu Ser Asn Gly Gln Pro Glu Asn Asn Tyr Lys Thr Thr Pro Pro Val
                390
                       395 400
Leu Asp Ser Asp Gly Ser Phe Phe Leu Tyr Ser Lys Leu Thr Val Asp
                     410 415
Lys Ser Arg Trp Gln Gln Gly Asn Val Phe Ser Cys Ser Val Met His
                 425 430
Glu Ala Leu His Asn His Tyr Thr Gln Lys Ser Leu Ser Leu Ser Pro
                       440
Gly Lys
  450
<210> 239
<211> 213
<212> PRT
<213> Artificial Sequence
<220>
<223> Humanized antibody - VL Chain
<400> 239
Asp Ile Gln Met Thr Gln Ser Pro Ser Thr Leu Ser Ala Ser Val Gly
                               10
Asp Arg Val Thr Ile Thr Cys Ser Ala Ser Ser Ser Val Gly Tyr Met
His Trp Tyr Gln Gln Lys Pro Gly Lys Ala Pro Lys Leu Leu Ile Tyr
Asp Thr Phe Lys Leu Ala Ser Gly Val Pro Ser Arg Phe Ser Gly Ser
                     55
Gly Ser Gly Thr Glu Phe Thr Leu Thr Ile Ser Ser Leu Gln Pro Asp
Asp Phe Ala Thr Tyr Tyr Cys Phe Gln Gly Ser Gly Tyr Pro Phe Thr
Phe Gly Gly Gly Thr Lys Val Glu Ile Lys Arg Thr Val Ala Ala Pro
          100 105
Ser Val Phe Ile Phe Pro Pro Ser Asp Glu Gln Leu Lys Ser Gly Thr
                        120
       115
```

```
Ala Ser Val Val Cys Leu Leu Asn Asn Phe Tyr Pro Arg Glu Ala Lys
         135 140
Val Gln Trp Lys Val Asp Asn Ala Leu Gln Ser Gly Asn Ser Gln Glu
      150 155
Ser Val Thr Glu Gln Asp Ser Lys Asp Ser Thr Tyr Ser Leu Ser Ser
   165 170 175
Thr Leu Thr Leu Ser Lys Ala Asp Tyr Glu Lys His Lys Val Tyr Ala
 180 185 190
Cys Glu Val Thr His Gln Gly Leu Ser Ser Pro Val Thr Lys Ser Phe
                     200
Asn Arg Gly Glu Cys
 210
<210> 240
<211> 450
<212> PRT
<213> Artificial Sequence
<220>
<223> Humanized antibody - VH Chain
<400> 240
Gln Val Thr Leu Arg Glu Ser Gly Pro Ala Leu Val Lys Pro Thr Gln
                      10 15
Thr Leu Thr Leu Thr Cys Thr Phe Ser Gly Phe Ser Leu Ser Thr Ala
                         25
Gly Met Ser Val Gly Trp Ile Arg Gln Pro Pro Gly Lys Ala Leu Glu
                     40
Trp Leu Ala Asp Ile Trp Trp Asp Asp Lys Lys Asp Tyr Asn Pro Ser
                 55
Leu Lys Ser Arg Leu Thr Ile Ser Lys Asp Thr Ser Lys Asn Gln Val
                    75 80
65 70
Val Leu Lys Val Thr Asn Met Asp Pro Ala Asp Thr Ala Thr Tyr Tyr
            85 90 95
Cys Ala Arg Ser Met Ile Thr Asn Phe Tyr Phe Asp Val Trp Gly Gln
       100 105 110
Gly Thr Thr Val Thr Val Ser Ser Ala Ser Thr Lys Gly Pro Ser Val
                     120 125
Phe Pro Leu Ala Pro Ser Ser Lys Ser Thr Ser Gly Gly Thr Ala Ala
                  135 140
Leu Gly Cys Leu Val Lys Asp Tyr Phe Pro Glu Pro Val Thr Val Ser
145 150 155 160
Trp Asn Ser Gly Ala Leu Thr Ser Gly Val His Thr Phe Pro Ala Val
           165 170 175
Leu Gln Ser Ser Gly Leu Tyr Ser Leu Ser Ser Val Val Thr Val Pro
                         185 190
Ser Ser Ser Leu Gly Thr Gln Thr Tyr Ile Cys Asn Val Asn His Lys
                      200
Pro Ser Asn Thr Lys Val Asp Lys Arg Val Glu Pro Lys Ser Cys Asp
Lys Thr His Thr Cys Pro Pro Cys Pro Ala Pro Glu Leu Leu Gly Gly
                               235 240
Pro Ser Val Phe Leu Phe Pro Pro Lys Pro Lys Asp Thr Leu Met Ile
            245
Ser Arg Thr Pro Glu Val Thr Cys Val Val Val Asp Val Ser His Glu
       2.60
                         2.65
Asp Pro Glu Val Lys Phe Asn Trp Tyr Val Asp Gly Val Glu Val His
   275 280
Asn Ala Lys Thr Lys Pro Arg Glu Glu Gln Tyr Asn Ser Thr Tyr Arg
                  295
                                   300
Val Val Ser Val Leu Thr Val Leu His Gln Asp Trp Leu Asn Gly Lys
```

```
310
                                  315
305
Glu Tyr Lys Cys Lys Val Ser Asn Lys Ala Leu Pro Ala Pro Ile Glu
                    330
       325
Lys Thr Ile Ser Lys Ala Lys Gly Gln Pro Arg Glu Pro Gln Val Tyr
                          345
Thr Leu Pro Pro Ser Arg Glu Glu Met Thr Lys Asn Gln Val Ser Leu
           360
Thr Cys Leu Val Lys Gly Phe Tyr Pro Ser Asp Ile Ala Val Glu Trp
   370 375
Glu Ser Asn Gly Gln Pro Glu Asn Asn Tyr Lys Thr Thr Pro Pro Val
                      395 400
       390
Leu Asp Ser Asp Gly Ser Phe Phe Leu Tyr Ser Lys Leu Thr Val Asp
                    410 415
          405
Lys Ser Arg Trp Gln Gln Gly Asn Val Phe Ser Cys Ser Val Met His
                 425 430
Glu Ala Leu His Asn His Tyr Thr Gln Lys Ser Leu Ser Leu Ser Pro
                      440
Gly Lys
 450
<210> 241
<211> 213
<212> PRT
<213> Artificial Sequence
<220>
<223> Humanized antibody - VH Chain
<400> 241
Asp Ile Gln Met Thr Gln Ser Pro Ser Thr Leu Ser Ala Ser Val Gly
                              10
Asp Arg Val Thr Ile Thr Cys Ser Ala Ser Ser Ser Val Gly Tyr Met
                        25
His Trp Tyr Gln Gln Lys Pro Gly Lys Ala Pro Lys Leu Leu Ile Tyr
Asp Thr Phe Lys Leu Ala Ser Gly Val Pro Ser Arg Phe Ser Gly Ser
                                     60
Gly Ser Gly Thr Glu Phe Thr Leu Thr Ile Ser Ser Leu Gln Pro Asp
                                  75 80
Asp Phe Ala Thr Tyr Tyr Cys Phe Gln Gly Ser Phe Tyr Pro Phe Thr
                              90
Phe Gly Gly Gly Thr Lys Val Glu Ile Lys Arg Thr Val Ala Ala Pro
                           105 110
Ser Val Phe Ile Phe Pro Pro Ser Asp Glu Gln Leu Lys Ser Gly Thr
                       120
Ala Ser Val Val Cys Leu Leu Asn Asn Phe Tyr Pro Arg Glu Ala Lys
                    135
                                     140
Val Gln Trp Lys Val Asp Asn Ala Leu Gln Ser Gly Asn Ser Gln Glu
                                  155 160
                150
Ser Val Thr Glu Gln Asp Ser Lys Asp Ser Thr Tyr Ser Leu Ser Ser
                              170
             165
Thr Leu Thr Leu Ser Lys Ala Asp Tyr Glu Lys His Lys Val Tyr Ala
         180 185 190
Cys Glu Val Thr His Gln Gly Leu Ser Ser Pro Val Thr Lys Ser Phe
                       200
Asn Arg Gly Glu Cys
   210
<210> 242
<211> 450
<212> PRT
```

<213> Artificial Sequence <220> <223> Humanized antibody - VH Chain <400> 242 Gln Val Thr Leu Arg Glu Ser Gly Pro Ala Leu Val Lys Pro Thr Gln 10 Thr Leu Thr Leu Thr Cys Thr Phe Ser Gly Phe Ser Leu Ser Thr Ala 25 Gly Met Ser Val Gly Trp Ile Arg Gln Pro Pro Gly Lys Ala Leu Glu 4.0 Trp Leu Ala Asp Ile Trp Trp Asp Asp Lys Lys Asp Tyr Asn Pro Ser 55 Leu Lys Ser Arg Leu Thr Ile Ser Lys Asp Thr Ser Lys Asn Gln Val Val Leu Lys Val Thr Asn Met Asp Pro Ala Asp Thr Ala Thr Tyr Tyr 8.5 90 Cys Ala Arg Asp Met Ile Thr Asn Phe Tyr Phe Asp Val Trp Gly Gln 100 105 Gly Thr Thr Val Thr Val Ser Ser Ala Ser Thr Lys Gly Pro Ser Val 120 Phe Pro Leu Ala Pro Ser Ser Lys Ser Thr Ser Gly Gly Thr Ala Ala 135 Leu Gly Cys Leu Val Lys Asp Tyr Phe Pro Glu Pro Val Thr Val Ser 150 Trp Asn Ser Gly Ala Leu Thr Ser Gly Val His Thr Phe Pro Ala Val 170 175 Leu Gln Ser Ser Gly Leu Tyr Ser Leu Ser Ser Val Val Thr Val Pro 180 185 Ser Ser Ser Leu Gly Thr Gln Thr Tyr Ile Cys Asn Val Asn His Lys 200 Pro Ser Asn Thr Lys Val Asp Lys Arg Val Glu Pro Lys Ser Cys Asp 215 Lys Thr His Thr Cys Pro Pro Cys Pro Ala Pro Glu Leu Leu Gly Gly 230 Pro Ser Val Phe Leu Phe Pro Pro Lys Pro Lys Asp Thr Leu Met Ile 245 250 Ser Arg Thr Pro Glu Val Thr Cys Val Val Val Asp Val Ser His Glu 265 Asp Pro Glu Val Lys Phe Asn Trp Tyr Val Asp Gly Val Glu Val His 280 Asn Ala Lys Thr Lys Pro Arg Glu Glu Gln Tyr Asn Ser Thr Tyr Arg 295 Val Val Ser Val Leu Thr Val Leu His Gln Asp Trp Leu Asn Gly Lys 310 315 Glu Tyr Lys Cys Lys Val Ser Asn Lys Ala Leu Pro Ala Pro Ile Glu 325 Lys Thr Ile Ser Lys Ala Lys Gly Gln Pro Arg Glu Pro Gln Val Tyr 340 Thr Leu Pro Pro Ser Arg Glu Glu Met Thr Lys Asn Gln Val Ser Leu Thr Cys Leu Val Lys Gly Phe Tyr Pro Ser Asp Ile Ala Val Glu Trp 380 370 375 Glu Ser Asn Gly Gln Pro Glu Asn Asn Tyr Lys Thr Thr Pro Pro Val 395 . 400 Leu Asp Ser Asp Gly Ser Phe Phe Leu Tyr Ser Lys Leu Thr Val Asp 410 415 Lys Ser Arg Trp Gln Gln Gly Asn Val Phe Ser Cys Ser Val Met His 425 Glu Ala Leu His Asn His Tyr Thr Gln Lys Ser Leu Ser Leu Ser Pro

85

```
445
                         440
      435
Gly Lys
  450
<210> 243
<211> 213
<212> PRT
<213> Artificial Sequence
<220>
<223> Humanized antibody - VL Chain
<400> 243
Asp Ile Gln Met Thr Gln Ser Pro Ser Thr Leu Ser Ala Ser Val Gly
                                  10
Asp Arg Val Thr Ile Thr Cys Ser Ala Ser Ser Ser Val Gly Tyr Met
                              25
His Trp Tyr Gln Gln Lys Pro Gly Lys Ala Pro Lys Leu Leu Ile Tyr
Asp Thr Phe Lys Leu Thr Ser Gly Val Pro Ser Arg Phe Ser Gly Ser
Gly Ser Gly Thr Glu Phe Thr Leu Thr Ile Ser Ser Leu Gln Pro Asp
                                      75
Asp Phe Ala Thr Tyr Tyr Cys Phe Gln Gly Ser Gly Tyr Pro Phe Thr
Phe Gly Gly Gly Thr Lys Val Glu Ile Lys Arg Thr Val Ala Ala Pro
                              105
Ser Val Phe Ile Phe Pro Pro Ser Asp Glu Gln Leu Lys Ser Gly Thr
                          120
Ala Ser Val Val Cys Leu Leu Asn Asn Phe Tyr Pro Arg Glu Ala Lys
                       135
Val Gln Trp Lys Val Asp Asn Ala Leu Gln Ser Gly Asn Ser Gln Glu
                  150
Ser Val Thr Glu Gln Asp Ser Lys Asp Ser Thr Tyr Ser Leu Ser Ser
                                  170
               165
Thr Leu Thr Leu Ser Lys Ala Asp Tyr Glu Lys His Lys Val Tyr Ala
           180 185
Cys Glu Val Thr His Gln Gly Leu Ser Ser Pro Val Thr Lys Ser Phe
       195
Asn Arg Gly Glu Cys
   210
<210> 244
<211> 450
<212> PRT
<213> Artificial Sequence
<220>
<223> Humanized antibody - VH Chain
<400> 244
Gln Val Thr Leu Arg Glu Ser Gly Pro Ala Leu Val Lys Pro Thr Gln
                                   10
Thr Leu Thr Leu Thr Cys Thr Phe Ser Gly Phe Ser Leu Ser Thr Ala
                               2.5
Gly Met Ser Val Gly Trp Ile Arg Gln Pro Pro Gly Lys Ala Leu Glu
                           40
Trp Leu Ala Asp Ile Trp Trp Asp Asp Lys Lys Asp Tyr Asn Pro Ser
                       55
                                           60
Leu Lys Ser Arg Leu Thr Ile Ser Lys Asp Thr Ser Lys Asn Gln Val
```

```
Val Leu Lys Val Thr Asn Met Asp Pro Ala Asp Thr Ala Thr Tyr Tyr
              90 95
Cys Ala Arg Asp Met Ile Thr Asn Phe Tyr Phe Asp Val Trp Gly Gln
   100 105 110
Gly Thr Thr Val Thr Val Ser Ser Ala Ser Thr Lys Gly Pro Ser Val
                  120 125
Phe Pro Leu Ala Pro Ser Ser Lys Ser Thr Ser Gly Gly Thr Ala Ala
130 135 140
Leu Gly Cys Leu Val Lys Asp Tyr Phe Pro Glu Pro Val Thr Val Ser
145 150 155 160
Trp Asn Ser Gly Ala Leu Thr Ser Gly Val His Thr Phe Pro Ala Val
          165 170 175
Leu Gln Ser Ser Gly Leu Tyr Ser Leu Ser Ser Val Val Thr Val Pro
 180 185 190
Ser Ser Ser Leu Gly Thr Gln Thr Tyr Ile Cys Asn Val Asn His Lys
 195 200 205
Pro Ser Asn Thr Lys Val Asp Lys Arg Val Glu Pro Lys Ser Cys Asp
210 215 220
Lys Thr His Thr Cys Pro Pro Cys Pro Ala Pro Glu Leu Leu Gly Gly
225 230 235 240
Pro Ser Val Phe Leu Phe Pro Pro Lys Pro Lys Asp Thr Leu Met Ile
                         250 255
 245
Ser Arg Thr Pro Glu Val Thr Cys Val Val Val Asp Val Ser His Glu
 260 265 270
Asp Pro Glu Val Lys Phe Asn Trp Tyr Val Asp Gly Val Glu Val His
 275 280 285
Asn Ala Lys Thr Lys Pro Arg Glu Glu Gln Tyr Asn Ser Thr Tyr Arg
 290 295 300
Val Val Ser Val Leu Thr Val Leu His Gln Asp Trp Leu Asn Gly Lys
   310 315 320
Glu Tyr Lys Cys Lys Val Ser Asn Lys Ala Leu Pro Ala Pro Ile Glu
           325 330 335
Lys Thr Ile Ser Lys Ala Lys Gly Gln Pro Arg Glu Pro Gln Val Tyr
      340 345 350
Thr Leu Pro Pro Ser Arg Glu Glu Met Thr Lys Asn Gln Val Ser Leu
                 360 365
Thr Cys Leu Val Lys Gly Phe Tyr Pro Ser Asp Ile Ala Val Glu Trp
                375 380
Glu Ser Asn Gly Gln Pro Glu Asn Asn Tyr Lys Thr Thr Pro Pro Val
385 , 390 395 400
Leu Asp Ser Asp Gly Ser Phe Phe Leu Tyr Ser Lys Leu Thr Val Asp
          405 410 415
Lys Ser Arg Trp Gln Gln Gly Asn Val Phe Ser Cys Ser Val Met His
   420 425 430
Glu Ala Leu His Asn His Tyr Thr Gln Lys Ser Leu Ser Leu Ser Pro
            440
Gly Lys
  450
<210> 245
<211> 213
<212> PRT
<213> Artificial Sequence
<223> Humanized antibody - VL Chain
<400> 245
Asp Ile Gln Met Thr Gln Ser Pro Ser Thr Leu Ser Ala Ser Val Gly
                         10
Asp Arg Val Thr Ile Thr Cys Ser Ala Ser Ser Arg Val Gly Tyr Met
```

```
25
His Trp Tyr Gln Gln Lys Pro Gly Lys Ala Pro Lys Leu Leu Ile Tyr
                         40
Asp Thr Phe Lys Leu Ala Ser Gly Val Pro Ser Arg Phe Ser Gly Ser
                      55
Gly Ser Gly Thr Glu Phe Thr Leu Thr Ile Ser Ser Leu Gln Pro Asp
                                     75
Asp Phe Ala Thr Tyr Tyr Cys Phe Gln Gly Ser Gly Tyr Pro Phe Thr
                                  90
Phe Gly Gly Gly Thr Lys Val Glu Ile Lys Arg Thr Val Ala Ala Pro
                             105
Ser Val Phe Ile Phe Pro Pro Ser Asp Glu Gln Leu Lys Ser Gly Thr
                         120
Ala Ser Val Val Cys Leu Leu Asn Asn Phe Tyr Pro Arg Glu Ala Lys
                      135
Val Gln Trp Lys Val Asp Asn Ala Leu Gln Ser Gly Asn Ser Gln Glu
                                     155
                  150
Ser Val Thr Glu Gln Asp Ser Lys Asp Ser Thr Tyr Ser Leu Ser Ser
              165
Thr Leu Thr Leu Ser Lys Ala Asp Tyr Glu Lys His Lys Val Tyr Ala
        180 185
Cys Glu Val Thr His Gln Gly Leu Ser Ser Pro Val Thr Lys Ser Phe
                         200
      195
Asn Arg Gly Glu Cys
 210
<210> 246
<211> 450
<212> PRT
<213> Artificial Sequence
<220>
<223> Humanized antibody - VH Chain
<400> 246
Gln Val Thr Leu Arg Glu Ser Gly Pro Ala Leu Val Lys Pro Thr Gln
Thr Leu Thr Leu Thr Cys Thr Phe Ser Gly Phe Ser Leu Ser Thr Ala
Gly Met Ser Val Gly Trp Ile Arg Gln Pro Pro Gly Lys Ala Leu Glu
                          40
Trp Leu Ala Asp Ile Trp Trp Asp Asp Lys Lys Asp Tyr Asn Pro Ser
                       55
Leu Lys Ser Arg Leu Thr Ile Ser Lys Asp Thr Ser Lys Asn Gln Val
                                      75
Val Leu Lys Val Thr Asn Met Asp Pro Ala Asp Thr Ala Thr Tyr Tyr
Cys Ala Arg Asp Met Ile Thr Asn Phe Tyr Phe Asp Val Trp Gly Gln
                              105
           100
Gly Thr Thr Val Thr Val Ser Ser Ala Ser Thr Lys Gly Pro Ser Val
                          120
Phe Pro Leu Ala Pro Ser Ser Lys Ser Thr Ser Gly Gly Thr Ala Ala
                                         140
                       135
Leu Gly Cys Leu Val Lys Asp Tyr Phe Pro Glu Pro Val Thr Val Ser
                                      1.55
                  150
Trp Asn Ser Gly Ala Leu Thr Ser Gly Val His Thr Phe Pro Ala Val
                                  170
               165
Leu Gln Ser Ser Gly Leu Tyr Ser Leu Ser Ser Val Val Thr Val Pro
                           185 190
 Ser Ser Ser Leu Gly Thr Gln Thr Tyr Ile Cys Asn Val Asn His Lys
                           200
```

```
Pro Ser Asn Thr Lys Val Asp Lys Arg Val Glu Pro Lys Ser Cys Asp
        215 220
Lys Thr His Thr Cys Pro Pro Cys Pro Ala Pro Glu Leu Leu Gly Gly
                       235
Pro Ser Val Phe Leu Phe Pro Pro Lys Pro Lys Asp Thr Leu Met Ile
                            250
            245
Ser Arg Thr Pro Glu Val Thr Cys Val Val Val Asp Val Ser His Glu
         260 265 270
Asp Pro Glu Val Lys Phe Asn Trp Tyr Val Asp Gly Val Glu Val His
                     280
Asn Ala Lys Thr Lys Pro Arg Glu Glu Gln Tyr Asn Ser Thr Tyr Arg
       295 300
Val Val Ser Val Leu Thr Val Leu His Gln Asp Trp Leu Asn Gly Lys
    310 315 320
Glu Tyr Lys Cys Lys Val Ser Asn Lys Ala Leu Pro Ala Pro Ile Glu
            325 330 335
Lys Thr Ile Ser Lys Ala Lys Gly Gln Pro Arg Glu Pro Gln Val Tyr
Thr Leu Pro Pro Ser Arg Glu Glu Met Thr Lys Asn Gln Val Ser Leu
                      360 365
Thr Cys Leu Val Lys Gly Phe Tyr Pro Ser Asp Ile Ala Val Glu Trp
                   375 380
Glu Ser Asn Gly Gln Pro Glu Asn Asn Tyr Lys Thr Thr Pro Pro Val
385 390 395 400
Leu Asp Ser Asp Gly Ser Phe Phe Leu Tyr Ser Lys Leu Thr Val Asp
            405 410 415
Lys Ser Arg Trp Gln Gln Gly Asn Val Phe Ser Cys Ser Val Met His
  420 425 430
Glu Ala Leu His Asn His Tyr Thr Gln Lys Ser Leu Ser Leu Ser Pro
                      440
 435
Gly Lys
   450
<210> 247
<211> 213
<212> PRT
<213> Artificial Sequence
<220>
<223> Humanized antibody - VL Chain
<400> 247
Asp Ile Gln Met Thr Gln Ser Pro Ser Thr Leu Ser Ala Ser Val Gly
                             10 15
Asp Arg Val Thr Ile Thr Cys Ser Ala Ser Ser Ser Val Gly Tyr Met
                          25
His Trp Tyr Gln Gln Lys Pro Gly Lys Ala Pro Lys Leu Leu Ile Tyr
                       40
Asp Thr Phe Arg Leu Ala Ser Gly Val Pro Ser Arg Phe Ser Gly Ser
                    55
Gly Ser Gly Thr Glu Phe Thr Leu Thr Ile Ser Ser Leu Gln Pro Asp
                                 75
                70
Asp Phe Ala Thr Tyr Tyr Cys Phe Gln Gly Ser Gly Tyr Pro Phe Thr
                             90
             85
Phe Gly Gly Gly Thr Lys Val Glu Ile Lys Arg Thr Val Ala Ala Pro
                          105
 Ser Val Phe Ile Phe Pro Pro Ser Asp Glu Gln Leu Lys Ser Gly Thr
                      120
 Ala Ser Val Val Cys Leu Leu Asn Asn Phe Tyr Pro Arg Glu Ala Lys
                   135
                                    140
Val Gln Trp Lys Val Asp Asn Ala Leu Gln Ser Gly Asn Ser Gln Glu
```

```
155
                   150
 Ser Val Thr Glu Gln Asp Ser Lys Asp Ser Thr Tyr Ser Leu Ser Ser
                        170
               165
 Thr Leu Thr Leu Ser Lys Ala Asp Tyr Glu Lys His Lys Val Tyr Ala
                    185
 Cys Glu Val Thr His Gln Gly Leu Ser Ser Pro Val Thr Lys Ser Phe
                          200
       195
 Asn Arq Gly Glu Cys
   210
 <210> 248
 <211> 450
 <212> PRT
 <213> Artificial Sequence
 <220>
 <223> Humanized antibody - VH Chain
 <400> 248
 Gln Val Thr Leu Arg Glu Ser Gly Pro Ala Leu Val Lys Pro Thr Gln
 Thr Leu Thr Leu Thr Cys Thr Phe Ser Gly Phe Ser Leu Ser Thr Ala
                               2.5
 Gly Met Ser Val Gly Trp Ile Arg Gln Pro Pro Gly Lys Ala Leu Glu
 Trp Leu Ala Asp Ile Trp Trp Asp Asp Lys, Lys His Tyr Asn Pro Ser
                       55
 Leu Lys Asp Arg Leu Thr Ile Ser Lys Asp Thr Ser Lys Asn Gln Val
                                       75
                    70
 Val Leu Lys Val Thr Asn Met Asp Pro Ala Asp Thr Ala Thr Tyr Tyr
                                  90
 Cys Ala Arg Asp Met Ile Phe Asn Trp Tyr Phe Asp Val Trp Gly Gln
                            105
 Gly Thr Thr Val Thr Val Ser Ser Ala Ser Thr Lys Gly Pro Ser Val
                          120
 Phe Pro Leu Ala Pro Ser Ser Lys Ser Thr Ser Gly Gly Thr Ala Ala
                                         140
          135
 Leu Gly Cys Leu Val Lys Asp Tyr Phe Pro Glu Pro Val Thr Val Ser
                            155
                   150
 Trp Asn Ser Gly Ala Leu Thr Ser Gly Val His Thr Phe Pro Ala Val
                                  170
                165
Leu Gln Ser Ser Gly Leu Tyr Ser Leu Ser Ser Val Val Thr Val Pro
             180
                               185
 Ser Ser Ser Leu Gly Thr Gln Thr Tyr Ile Cys Asn Val Asn His Lys
                            200
 Pro Ser Asn Thr Lys Val Asp Lys Arg Val Glu Pro Lys Ser Cys Asp
                                          220
                        215
 Lys Thr His Thr Cys Pro Pro Cys Pro Ala Pro Glu Leu Leu Gly Gly
                                      235
                   230
  Pro Ser Val Phe Leu Phe Pro Pro Lys Pro Lys Asp Thr Leu Met Ile
                                  250
                245
  Ser Arg Thr Pro Glu Val Thr Cys Val Val Val Asp Val Ser His Glu
                                265
             260
  Asp Pro Glu Val Lys Phe Asn Trp Tyr Val Asp Gly Val Glu Val His
                           280
  Asn Ala Lys Thr Lys Pro Arg Glu Glu Gln Tyr Asn Ser Thr Tyr Arg
                                           300
                        295
  Val Val Ser Val Leu Thr Val Leu His Gln Asp Trp Leu Asn Gly Lys
  305 310
                                      315
  Glu Tyr Lys Cys Lys Val Ser Asn Lys Ala Leu Pro Ala Pro Ile Glu
                                   330
```

```
Lys Thr Ile Ser Lys Ala Lys Gly Gln Pro Arg Glu Pro Gln Val Tyr
   340 345 350
Thr Leu Pro Pro Ser Arg Glu Glu Met Thr Lys Asn Gln Val Ser Leu
            360 365
Thr Cys Leu Val Lys Gly Phe Tyr Pro Ser Asp Ile Ala Val Glu Trp
  370 375 380
Glu Ser Asn Gly Gln Pro Glu Asn Asn Tyr Lys Thr Thr Pro Pro Val
385 390 395 400
Leu Asp Ser Asp Gly Ser Phe Phe Leu Tyr Ser Lys Leu Thr Val Asp
           405 410 415
Lys Ser Arg Trp Gln Gln Gly Asn Val Phe Ser Cys Ser Val Met His
   420 425 430
Glu Ala Leu His Asn His Tyr Thr Gln Lys Ser Leu Ser Leu Ser Pro
Gly Lys
 450
<210> 249
<211> 213
<212> PRT
<213> Artificial Sequence
<223> Humanized antibody - VL Chain
<400> 249
Asp Ile Gln Met Thr Gln Ser Pro Ser Thr Leu Ser Ala Ser Val Gly
                         10
Asp Arg Val Thr Ile Thr Cys Ser Pro Ser Ser Arg Val Gly Tyr Met
                         25
His Trp Tyr Gln Gln Lys Pro Gly Lys Ala Pro Lys Leu Leu Ile Tyr
                     40
Asp Thr Tyr Arg His Ser Ser Gly Val Pro Ser Arg Phe Ser Gly Ser
                   55
Gly Ser Gly Thr Glu Phe Thr Leu Thr Ile Ser Ser Leu Gln Pro Asp
                               75
Asp Phe Ala Thr Tyr Tyr Cys Phe Gln Gly Ser Gly Tyr Pro Phe Thr
Phe Gly Gly Gly Thr Lys Val Glu Ile Lys Arg Thr Val Ala Ala Pro
                       105 110
Ser Val Phe Ile Phe Pro Pro Ser Asp Glu Gln Leu Lys Ser Gly Thr
    115 120 125
Ala Ser Val Val Cys Leu Leu Asn Asn Phe Tyr Pro Arg Glu Ala Lys
                  135 140
Val Gln Trp Lys Val Asp Asn Ala Leu Gln Ser Gly Asn Ser Gln Glu
               150 155 160
Ser Val Thr Glu Gln Asp Ser Lys Asp Ser Thr Tyr Ser Leu Ser Ser
                            170
            165
Thr Leu Thr Leu Ser Lys Ala Asp Tyr Glu Lys His Lys Val Tyr Ala
    180 185 190
Cys Glu Val Thr His Gln Gly Leu Ser Ser Pro Val Thr Lys Ser Phe
  195
                     200
Asn Arg Gly Glu Cys
  210
<210> 250
<211> 450
<212> PRT
<213> Artificial Sequence
<220>
```

<223> Humanized antibody - VH Chain

<400> 250 Gln Val Thr Leu Arg Glu Ser Gly Pro Ala Leu Val Lys Pro Thr Gln Thr Leu Thr Leu Thr Cys Thr Phe Ser Gly Phe Ser Leu Ser Thr Ala Gly Met Ser Val Gly Trp Ile Arg Gln Pro Pro Gly Lys Ala Leu Glu Trp Leu Ala Asp Ile Trp Trp Asp Gly Lys Lys His Tyr Asn Pro Ser Leu Lys Asp Arg Leu Thr Ile Ser Lys Asp Thr Ser Lys Asn Gln Val 70 Val Leu Lys Val Thr Asn Met Asp Pro Ala Asp Thr Ala Thr Tyr Tyr 85 Cys Ala Arg Asp Met Ile Phe Asn Trp Tyr Phe Asp Val Trp Gly Gln 105 100 Gly Thr Thr Val Thr Val Ser Ser Ala Ser Thr Lys Gly Pro Ser Val 120 Phe Pro Leu Ala Pro Ser Ser Lys Ser Thr Ser Gly Gly Thr Ala Ala 135 Leu Gly Cys Leu Val Lys Asp Tyr Phe Pro Glu Pro Val Thr Val Ser 155 150 Trp Asn Ser Gly Ala Leu Thr Ser Gly Val His Thr Phe Pro Ala Val 170 165 Leu Gln Ser Ser Gly Leu Tyr Ser Leu Ser Ser Val Val Thr Val Pro 185 180 Ser Ser Ser Leu Gly Thr Gln Thr Tyr Ile Cys Asn Val Asn His Lys 200 Pro Ser Asn Thr Lys Val Asp Lys Arg Val Glu Pro Lys Ser Cys Asp 220 215 Lys Thr His Thr Cys Pro Pro Cys Pro Ala Pro Glu Leu Leu Gly Gly 235 230 Pro Ser Val Phe Leu Phe Pro Pro Lys Pro Lys Asp Thr Leu Met Ile 250 245 Ser Arg Thr Pro Glu Val Thr Cys Val Val Val Asp Val Ser His Glu 265 260 Asp Pro Glu Val Lys Phe Asn Trp Tyr Val Asp Gly Val Glu Val His 280 Asn Ala Lys Thr Lys Pro Arg Glu Glu Gln Tyr Asn Ser Thr Tyr Arg 295 Val Val Ser Val Leu Thr Val Leu His Gln Asp Trp Leu Asn Gly Lys 315 310 Glu Tyr Lys Cys Lys Val Ser Asn Lys Ala Leu Pro Ala Pro Ile Glu 330 325 Lys Thr Ile Ser Lys Ala Lys Gly Gln Pro Arg Glu Pro Gln Val Tyr 345 340 Thr Leu Pro Pro Ser Arg Glu Glu Met Thr Lys Asn Gln Val Ser Leu 360 Thr Cys Leu Val Lys Gly Phe Tyr Pro Ser Asp Ile Ala Val Glu Trp 380 375 Glu Ser Asn Gly Gln Pro Glu Asn Asn Tyr Lys Thr Thr Pro Pro Val 390 395 Leu Asp Ser Asp Gly Ser Phe Phe Leu Tyr Ser Lys Leu Thr Val Asp 410 Lys Ser Arg Trp Gln Gln Gly Asn Val Phe Ser Cys Ser Val Met His 425 430 Glu Ala Leu His Asn His Tyr Thr Gln Lys Ser Leu Ser Leu Ser Pro Gly Lys 450

92

```
<210> 251
 <211> 213
 <212> PRT
 <213> Artificial Sequence
 <220>
 <223> Humanized antibody - VL Chain
 <400> 251
 Asp Ile Gln Met Thr Gln Ser Pro Ser Thr Leu Ser Ala Ser Val Glv
                                    10
 Asp Arg Val Thr Ile Thr Cys Ser Leu Ser Ser Ser Val Gly Tyr Met
 His Trp Tyr Gln Gln Lys Pro Gly Lys Ala Pro Lys Leu Leu Ile Tyr
 Asp Thr Phe Phe His Arg Ser Gly Val Pro Ser Arg Phe Ser Gly Ser
 Gly Ser Gly Thr Glu Phe Thr Leu Thr Ile Ser Ser Leu Gln Pro Asp
                                        75
 Asp Phe Ala Thr Tyr Tyr Cys Phe Gln Gly Ser Gly Tyr Pro Phe Thr
 Phe Gly Gly Gly Thr Lys Val Glu Ile Lys Arg Thr Val Ala Ala Pro
                               105
 Ser Val Phe Ile Phe Pro Pro Ser Asp Glu Gln Leu Lys Ser Gly Thr
                           120
                                             125
 Ala Ser Val Val Cys Leu Leu Asn Asn Phe Tyr Pro Arg Glu Ala Lys
                        135
 Val Gln Trp Lys Val Asp Asn Ala Leu Gln Ser Gly Asn Ser Gln Glu
                                        155
 Ser Val Thr Glu Gln Asp Ser Lys Asp Ser Thr Tyr Ser Leu Ser Ser
                                    170
 Thr Leu Thr Leu Ser Lys Ala Asp Tyr Glu Lys His Lys Val Tyr Ala
                               185
 Cys Glu Val Thr His Gln Gly Leu Ser Ser Pro Val Thr Lys Ser Phe
 Asn Arg Gly Glu Cys
  210
 <210> 252
 <211> 450
 <212> PRT
<213> Artificial Sequence
 <220>
 <223> Humanized antibody - VH Chain
 <400> 252
 Gln Val Thr Leu Arg Glu Ser Gly Pro Ala Leu Val Lys Pro Thr Gln
                                    10
 Thr Leu Thr Leu Thr Cys Thr Phe Ser Gly Phe Ser Leu Ser Thr Ala
                                25
 Gly Met Ser Val Gly Trp Ile Arg Gln Pro Pro Gly Lys Ala Leu Glu
                           40
                                               4.5
 Trp Leu Ala Asp Ile Trp Trp Asp Asp Lys Lys His Tyr Asn Pro Ser
                        55
                                            60
 Leu Lys Asp Arg Leu Thr Ile Ser Lys Asp Thr Ser Lys Asn Gln Val
                    70
                                        75
 Val Leu Lys Val Thr Asn Met Asp Pro Ala Asp Thr Ala Thr Tyr Tyr
                                90
 Cys Ala Arg Asp Met Ile Phe Asn Phe Tyr Phe Asp Val Trp Gly Gln
             100
                                105
```

```
Gly Thr Thr Val Thr Val Ser Ser Ala Ser Thr Lys Gly Pro Ser Val
           120 125
 115
Phe Pro Leu Ala Pro Ser Ser Lys Ser Thr Ser Gly Gly Thr Ala Ala
                 135
Leu Gly Cys Leu Val Lys Asp Tyr Phe Pro Glu Pro Val Thr Val Ser
145 150 155 160
Trp Asn Ser Gly Ala Leu Thr Ser Gly Val His Thr Phe Pro Ala Val
           165 170 175
Leu Gln Ser Ser Gly Leu Tyr Ser Leu Ser Ser Val Val Thr Val Pro
                       185 190
         180
Ser Ser Ser Leu Gly Thr Gln Thr Tyr Ile Cys Asn Val Asn His Lys
                         205
                    200
 195
Pro Ser Asn Thr Lys Val Asp Lys Arg Val Glu Pro Lys Ser Cys Asp
 210 215 220
Lys Thr His Thr Cys Pro Pro Cys Pro Ala Pro Glu Leu Leu Gly Gly
                             235 240
   230
Pro Ser Val Phe Leu Phe Pro Pro Lys Pro Lys Asp Thr Leu Met Ile
                           250 255
           245
Ser Arg Thr Pro Glu Val Thr Cys Val Val Val Asp Val Ser His Glu
                       265 270
        260
Asp Pro Glu Val Lys Phe Asn Trp Tyr Val Asp Gly Val Glu Val His
  275 280 285
Asn Ala Lys Thr Lys Pro Arg Glu Glu Gln Tyr Asn Ser Thr Tyr Arg
       295 300
Val Val Ser Val Leu Thr Val Leu His Gln Asp Trp Leu Asn Gly Lys
    310
Glu Tyr Lys Cys Lys Val Ser Asn Lys Ala Leu Pro Ala Pro Ile Glu
           325 330 335
Lys Thr Ile Ser Lys Ala Lys Gly Gln Pro Arg Glu Pro Gln Val Tyr
                        345 350
        340
Thr Leu Pro Pro Ser Arg Glu Glu Met Thr Lys Asn Gln Val Ser Leu
                     360 365
Thr Cys Leu Val Lys Gly Phe Tyr Pro Ser Asp Ile Ala Val Glu Trp
                  375 380
Glu Ser Asn Gly Gln Pro Glu Asn Asn Tyr Lys Thr Thr Pro Pro Val
385 390 395 400
Leu Asp Ser Asp Gly Ser Phe Phe Leu Tyr Ser Lys Leu Thr Val Asp
            405 410 415
Lys Ser Arg Trp Gln Gln Gly Asn Val Phe Ser Cys Ser Val Met His
        420 425 430
Glu Ala Leu His Asn His Tyr Thr Gln Lys Ser Leu Ser Leu Ser Pro
   435 440
Gly Lys
  450
 <210> 253
<210    233
<211>    213
<212>    PRT
<213>    Artificial Sequence
 <223> Humanized antibody - VL Chain
 <400> 253
 Asp Ile Gln Met Thr Gln Ser Pro Ser Thr Leu Ser Ala Ser Val Gly
                            10
 Asp Arg Val Thr Ile Thr Cys Ser Ala Ser Ser Arg Val Gly Tyr Met
                         25
 His Trp Tyr Gln Gln Lys Pro Gly Lys Ala Pro Lys Leu Leu Ile Tyr
 Asp Thr Leu Leu Leu Asp Ser Gly Val Pro Ser Arg Phe Ser Gly Ser
```

```
55
Gly Ser Gly Thr Glu Phe Thr Leu Thr Ile Ser Ser Leu Gln Pro Asp
                 70
Asp Phe Ala Thr Tyr Tyr Cys Phe Gln Gly Ser Gly Tyr Pro Phe Thr
            85
Phe Gly Gly Gly Thr Lys Val Glu Ile Lys Arg Thr Val Ala Ala Pro
         100
                 105
Ser Val Phe Ile Phe Pro Pro Ser Asp Glu Gln Leu Lys Ser Gly Thr
                                        125
   115 120
Ala Ser Val Val Cys Leu Leu Asn Asn Phe Tyr Pro Arg Glu Ala Lys
        135
Val Gln Trp Lys Val Asp Asn Ala Leu Gln Ser Gly Asn Ser Gln Glu
      150
                        155
Ser Val Thr Glu Gln Asp Ser Lys Asp Ser Thr Tyr Ser Leu Ser Ser
                  170 175
      165
Thr Leu Thr Leu Ser Lys Ala Asp Tyr Glu Lys His Lys Val Tyr Ala
      180 185
Cys Glu Val Thr His Gln Gly Leu Ser Ser Pro Val Thr Lys Ser Phe
              200
 195
Asn Arg Gly Glu Cys
<210> 254
<211> 450
<212> PRT
<213> Artificial Sequence
<220>
<223> Humanized antibody - VH Chain
<400> 254
Gln Val Thr Leu Arg Glu Ser Gly Pro Ala Leu Val Lys Pro Thr Gln
                              10
Thr Leu Thr Leu Thr Cys Thr Phe Ser Gly Phe Ser Leu Ser Thr Ala
                          25
Gly Met Ser Val Gly Trp Ile Arg Gln Pro Pro Gly Lys Ala Leu Glu
                       40
Trp Leu Ala Asp Ile Trp Trp Asp Asp Lys Lys His Tyr Asn Pro Ser
                   55
Leu Lys Asp Arg Leu Thr Ile Ser Lys Asp Thr Ser Lys Asn Gln Val
                                 75
                70
Val Leu Lys Val Thr Asn Met Asp Pro Ala Asp Thr Ala Thr Tyr Tyr
                              90
Cys Ala Arg Asp Met Ile Phe Asn Phe Tyr Phe Asp Val Trp Gly Gln
                                 110
                          105
Gly Thr Thr Val Thr Val Ser Ser Ala Ser Thr Lys Gly Pro Ser Val
                       120
Phe Pro Leu Ala Pro Ser Ser Lys Ser Thr Ser Gly Gly Thr Ala Ala
                    135
                           140
Leu Gly Cys Leu Val Lys Asp Tyr Phe Pro Glu Pro Val Thr Val Ser
                        155
Trp Asn Ser Gly Ala Leu Thr Ser Gly Val His Thr Phe Pro Ala Val
                              170 175
Leu Gln Ser Ser Gly Leu Tyr Ser Leu Ser Ser Val Val Thr Val Pro
                           185
Ser Ser Ser Leu Gly Thr Gln Thr Tyr Ile Cys Asn Val Asn His Lys
                       200
Pro Ser Asn Thr Lys Val Asp Lys Arg Val Glu Pro Lys Ser Cys Asp
                 215
                                     220
Lys Thr His Thr Cys Pro Pro Cys Pro Ala Pro Glu Leu Leu Gly Gly
                230
```

```
Pro Ser Val Phe Leu Phe Pro Pro Lys Pro Lys Asp Thr Leu Met Ile
                               250
Ser Arg Thr Pro Glu Val Thr Cys Val Val Val Asp Val Ser His Glu
                           265
Asp Pro Glu Val Lys Phe Asn Trp Tyr Val Asp Gly Val Glu Val His
           280
Asn Ala Lys Thr Lys Pro Arg Glu Glu Gln Tyr Asn Ser Thr Tyr Arg
                    295
Val Val Ser Val Leu Thr Val Leu His Gln Asp Trp Leu Asn Gly Lys
   310 315
Glu Tyr Lys Cys Lys Val Ser Asn Lys Ala Leu Pro Ala Pro Ile Glu
            325 330 335
Lys Thr Ile Ser Lys Ala Lys Gly Gln Pro Arg Glu Pro Gln Val Tyr
                          345 350
Thr Leu Pro Pro Ser Arg Glu Glu Met Thr Lys Asn Gln Val Ser Leu
                        360
Thr Cys Leu Val Lys Gly Phe Tyr Pro Ser Asp Ile Ala Val Glu Trp
                   375
                                     380
Glu Ser Asn Gly Gln Pro Glu Asn Asn Tyr Lys Thr Thr Pro Pro Val
    390 395
Leu Asp Ser Asp Gly Ser Phe Phe Leu Tyr Ser Lys Leu Thr Val Asp
             405 410 415
Lys Ser Arg Trp Gln Gln Gly Asn Val Phe Ser Cys Ser Val Met His
   420 425 430
Glu Ala Leu His Asn His Tyr Thr Gln Lys Ser Leu Ser Leu Ser Pro
Gly Lys
  450
<210> 255
<211> 213
<212> PRT
<213> Artificial Sequence
<220>
<223> Humanized antibody - VL Chain
<400> 255
Asp Ile Gln Met Thr Gln Ser Pro Ser Thr Leu Ser Ala Ser Val Gly
                  10
Asp Arg Val Thr Ile Thr Cys Ser Ala Ser Ser Arg Val Gly Tyr Met
His Trp Tyr Gln Gln Lys Pro Gly Lys Ala Pro Lys Leu Leu Ile Tyr
          40
Asp Thr Ser Lys Leu Ala Ser Gly Val Pro Ser Arg Phe Ser Gly Ser
                    55
Gly Ser Gly Thr Glu Phe Thr Leu Thr Ile Ser Ser Leu Gln Pro Asp
                70
                                  75
Asp Phe Ala Thr Tyr Tyr Cys Phe Gln Gly Ser Gly Tyr Pro Phe Thr
                               90 95
Phe Gly Gly Gly Thr Lys Val Glu Ile Lys Arg Thr Val Ala Ala Pro
                          105
Ser Val Phe Ile Phe Pro Pro Ser Asp Glu Gln Leu Lys Ser Gly Thr
                       120
Ala Ser Val Val Cys Leu Leu Asn Asn Phe Tyr Pro Arg Glu Ala Lys
                    135
Val Gln Trp Lys Val Asp Asn Ala Leu Gln Ser Gly Asn Ser Gln Glu
                150
Ser Val Thr Glu Gln Asp Ser Lys Asp Ser Thr Tyr Ser Leu Ser Ser
                              170
Thr Leu Thr Leu Ser Lys Ala Asp Tyr Glu Lys His Lys Val Tyr Ala
```

```
180
                              185
Cys Glu Val Thr His Gln Gly Leu Ser Ser Pro Val Thr Lys Ser Phe
       195
                      200
Asn Arg Gly Glu Cys
  210
<210> 256
<211> 450
<212> PRT
<213> Artificial Sequence
<220>
<223> Humanized antibody - VH Chain
<400> 256
Gln Val Thr Leu Arg Glu Ser Gly Pro Ala Leu Val Lys Pro Thr Gln
                                   10
Thr Leu Thr Leu Thr Cys Thr Phe Ser Gly Phe Ser Leu Ser Thr Ala
                               25
Gly Met Ser Val Gly Trp Ile Arg Gln Pro Pro Gly Lys Ala Leu Glu
                           40
Trp Leu Ala Asp Ile Trp Trp Asp Asp Lys Lys His Tyr Asn Pro Ser
                      55
Leu Lys Asp Arg Leu Thr Ile Ser Lys Asp Thr Ser Lys Asn Gln Val
Val Leu Lys Val Thr Asn Met Asp Pro Ala Asp Thr Ala Thr Tyr Tyr
                                  90
Cys Ala Arg Asp Met Ile Phe Asn Phe Tyr Phe Asp Val Trp Gly Gln
                               105
Gly Thr Thr Val Thr Val Ser Ser Ala Ser Thr Lys Gly Pro Ser Val
                           120
Phe Pro Leu Ala Pro Ser Ser Lys Ser Thr Ser Gly Gly Thr Ala Ala
                       135
Leu Gly Cys Leu Val Lys Asp Tyr Phe Pro Glu Pro Val Thr Val Ser
                   150
                                      155
Trp Asn Ser Gly Ala Leu Thr Ser Gly Val His Thr Phe Pro Ala Val
                                   170
Leu Gln Ser Ser Gly Leu Tyr Ser Leu Ser Ser Val Val Thr Val Pro
                               185
Ser Ser Ser Leu Gly Thr Gln Thr Tyr Ile Cys Asn Val Asn His Lys
                           200
Pro Ser Asn Thr Lys Val Asp Lys Arg Val Glu Pro Lys Ser Cys Asp
                      215
Lys Thr His Thr Cys Pro Pro Cys Pro Ala Pro Glu Leu Leu Gly Gly
                   230
                                       235
Pro Ser Val Phe Leu Phe Pro Pro Lys Pro Lys Asp Thr Leu Met Ile
               245
                                   250
Ser Arg Thr Pro Glu Val Thr Cys Val Val Val Asp Val Ser His Glu
                               265
Asp Pro Glu Val Lys Phe Asn Trp Tyr Val Asp Gly Val Glu Val His
                           280
Asn Ala Lys Thr Lys Pro Arg Glu Glu Gln Tyr Asn Ser Thr Tyr Arg
                      295
Val Val Ser Val Leu Thr Val Leu His Gln Asp Trp Leu Asn Gly Lys
                   310
                                       315
Glu Tyr Lys Cys Lys Val Ser Asn Lys Ala Leu Pro Ala Pro Ile Glu
               325
                                   330
Lys Thr Ile Ser Lys Ala Lys Gly Gln Pro Arg Glu Pro Gln Val Tyr
                               345
Thr Leu Pro Pro Ser Arg Glu Glu Met Thr Lys Asn Gln Val Ser Leu
```

```
Thr Cys Leu Val Lys Gly Phe Tyr Pro Ser Asp Ile Ala Val Glu Trp
370 375 380
Glu Ser Asn Gly Gln Pro Glu Asn Asn Tyr Lys Thr Thr Pro Pro Val
            390 395 400
Leu Asp Ser Asp Gly Ser Phe Phe Leu Tyr Ser Lys Leu Thr Val Asp
         405 410 415
Lys Ser Arg Trp Gln Gln Gly Asn Val Phe Ser Cys Ser Val Met His
  420 425 430
Glu Ala Leu His Asn His Tyr Thr Gln Lys Ser Leu Ser Leu Ser Pro
                     440
Gly Lys
 450
<210> 257
<211> 213
<212> PRT
<213> Artificial Sequence
<223> Humanized antibody - VL Chain
<400> 257
Asp Ile Gln Met Thr Gln Ser Pro Ser Thr Leu Ser Ala Ser Val Gly
                    10 15
Asp Arg Val Thr Ile Thr Cys Ser Ala Ser Ser Arg Val Gly Tyr Met
His Trp Tyr Gln Gln Lys Pro Gly Lys Ala Pro Lys Leu Leu Ile Tyr
                      40
                          · 45
Asp Thr Ser Phe Leu Asp Ser Gly Val Pro Ser Arg Phe Ser Gly Ser
                   55
Gly Ser Gly Thr Glu Phe Thr Leu Thr Ile Ser Ser Leu Gln Pro Asp
               70
Asp Phe Ala Thr Tyr Tyr Cys Phe Gln Gly Ser Gly Tyr Pro Phe Thr
Phe Gly Gly Gly Thr Lys Val Glu Ile Lys Arg Thr Val Ala Ala Pro
                        105 110
         1.00
Ser Val Phe Ile Phe Pro Pro Ser Asp Glu Gln Leu Lys Ser Gly Thr
                      120 125
Ala Ser Val Val Cys Leu Leu Asn Asn Phe Tyr Pro Arg Glu Ala Lys
                   135 140
Val Gln Trp Lys Val Asp Asn Ala Leu Gln Ser Gly Asn Ser Gln Glu
              150
Ser Val Thr Glu Gln Asp Ser Lys Asp Ser Thr Tyr Ser Leu Ser Ser
                            170 175
            165
Thr Leu Thr Leu Ser Lys Ala Asp Tyr Glu Lys His Lys Val Tyr Ala
       180 185 190
Cys Glu Val Thr His Gln Gly Leu Ser Ser Pro Val Thr Lys Ser Phe
                      200
  195
Asn Arg Gly Glu Cys
  210
<210> 258
<211> 26
<212> DNA
<213> Artificial
<223> Description of Artificial Sequence: Primer
<400> 258
agtgtcttaa ccagcaaagt gttaga
```

26

```
<210> 259
<211> 26
<212> DNA
<213> Artificial Sequence
<220>
<223> Description of Artificial Sequence: Primer
<400> 259
tcattgactt gagatattga tgcatc
                                                                  26
<210> 260
<211> 15
<212> PRT
<213> Artificial Sequence
<223> Linker for constructing humanized antibodies
Gly Gly Gly Ser Gly Gly Gly Ser Gly Gly Gly Ser
                                   10
<210> 261
<211> 15
<212> PRT
<213> Artificial Sequence
<223> Linker for constructing humanized antibodies
Glu Ser Gly Arg Ser Gly Gly Gly Gly Ser Gly Gly Gly Ser
                                   10
<210> 262
<211> 14
<212> PRT
<213> Artificial Sequence
<223> Linker for constructing humanized antibodies
Glu Gly Lys Ser Ser Gly Ser Gly Ser Glu Ser Lys Ser Thr
<210> 263
<211> 15
<212> PRT
<213> Artificial Sequence
<220>
<223> Linker for constructing humanized antibodies
Glu Gly Lys Ser Ser Gly Ser Gly Ser Glu Ser Lys Ser Thr Gln
<210> 264
```

```
<211> 14
<212> PRT
<213> Artificial Sequence
<223> Linker for constructing humanized antibodies
<400> 264
Glu Gly Lys Ser Ser Gly Ser Glu Ser Lys Val Asp
<210> 265
<211> 14
<212> PRT
<213> Artificial Sequence
<223> Linker for constructing humanized antibodies
<400> 265
Gly Ser Thr Ser Gly Ser Gly Lys Ser Ser Glu Gly Lys Gly
<210> 266
<211> 18
<212> PRT
<213> Artificial Sequence
<223> Linker for constructing humanized antibodies
<400> 266
Lys Glu Ser Gly Ser Val Ser Ser Glu Gln Leu Ala Gln Phe Arg Ser
Leu Asp
<210> 267
<211> 16
<212> PRT
<213> Artificial Sequence
<220>
<223> Linker for constructing humanized antibodies
Glu Ser Gly Ser Val Ser Ser Glu Glu Leu Ala Phe Arg Ser Leu Asp
                                    10
<210> 268
<211> 4
<212> PRT
<213> Homo sapiens
<220>
<223> intrabody
<400> 268
Lys Asp Glu Leu
```

```
<210> 269
<211> 4
<212> PRT
<213> Homo sapiens
<220>
<223> intrabody
<400> 269
Asp Asp Glu Leu
<210> 270
<211> 4
<212> PRT
<213> Homo sapiens
<220>
<223> intrabody
<400> 270
Asp Glu Glu Leu
<210> 271
<211> 4
<212> PRT
<213> Homo sapiens
<220>
<223> intrabody
<400> 271
Gln Glu Asp Leu
<210> 272
<211> 4
<212> PRT
<213> Homo sapiens
<220>
<223> intrabody
<400> 272
Arg Asp Glu Leu
<210> 273
<211> 7
<212> PRT
<213> Homo sapiens
<220>
<223> intrabody
<400> 273
Pro Lys Lys Lys Arg Lys Val
```

```
<210> 274
<211> 7
<212> PRT
<213> Homo sapiens
<220>
<223> intrabody
<400> 274
Pro Gln Lys Lys Ile Lys Ser
<210> 275
<211> 5
<212> PRT
<213> Homo sapiens
<220>
<223> intrabody
<400> 275
Gln Pro Lys Lys Pro
<210> 276
<211> 4
<212> PRT
<213> Homo sapiens
<220>
<223> intrabody
<400> 276
Arg Lys Lys Arg
<210> 277
<211> 5
<212> PRT
<213> Homo sapiens
<220>
<223> intrabody
<400> 277
Lys Lys Lys Arg Lys
<210> 278
<211> 12
<212> PRT
<213> Homo sapiens
<220>
<223> intrabody
<400> 278
Arg Lys Lys Arg Arg Gln Arg Arg Arg Ala His Gln
<210> 279
```

```
<211> 16
<212> PRT
<213> Homo sapiens
<220>
<223> intrabody
<400> 279
Arg Gln Ala Arg Arg Asn Arg Arg Arg Arg Trp Arg Glu Arg Gln Arg
<210> 280
<211> 19
<212> PRT
<213> Homo sapiens
<220>
<223> intrabody
<400> 280
Met Pro Leu Thr Arg Arg Pro Ala Ala Ser Gln Ala Leu Ala Pro
Pro Thr Pro
<210> 281
<211> 15
<212> PRT
<213> Homo sapiens
<220>
<223> intrabody
 Met Asp Asp Gln Arg Asp Leu Ile Ser Asn Asn Glu Gln Leu Pro
                                10
<210> 282
<211> 32
<212> PRT
<213> Homo sapiens
 <220>
 <223> intrabody
 <220>
 <221> misc_feature <222> 7, 8, 32,
 <223> Xaa can be any naturally occurring amino acid
 Met Leu Phe Asn Leu Arg Xaa Xaa Leu Asn Asn Ala Ala Phe Arg His
                                       10
 Gly His Asn Phe Met Val Arg Asn Phe Arg Cys Gly Gln Pro Leu Xaa
                                    25
 <210> 283
 <211> 3
 <212> PRT
```

```
<213> Homo sapiens
<220>
<223> intrabody
<400> 283
Ala Lys Leu
<210> 284
<211> 6
<212> PRT
<213> Homo sapiens
<220>
<223> intrabody
<400> 284
Ser Asp Tyr Gln Arg Leu
<210> 285
<211> 8
<212> PRT
<213> Homo sapiens
<220>
<223> intrabody
<400> 285
Gly Cys Val Cys Ser Ser Asn Pro
<210> 286
<211> 8
<212> PRT
<213> Homo sapiens
<220>
<223> intrabody
<400> 286
Gly Gln Thr Val Thr Thr Pro Leu
<210> 287
<211> 8
<212> PRT
<213> Homo sapiens
<220>
<223> intrabody
 <400> 287
 Gly Gln Glu Leu Ser Gln His Glu
 <210> 288
 <211> 8
 <212> PRT
```

```
<213> Homo sapiens
<220>
<223> intrabody
<400> 288
Gly Asn Ser Pro Ser Tyr Asn Pro
<210> 289
<211> 8
<212> PRT
<213> Homo sapiens
<220>
<223> intrabody
<400> 289
Gly Val Ser Gly Ser Lys Gly Gln
1 5
<210> 290
<211> 8
<212> PRT
<213> Homo sapiens
<220>
<223> intrabody
<400> 290
Gly Gln Thr Ile Thr Thr Pro Leu
<210> 291
<211> 8
<212> PRT
<213> Homo sapiens
 <220>
 <223> intrabody
 <400> 291
 Gly Gln Thr Leu Thr Thr Pro Leu
 <210> 292
<211> 8
<212> PRT
<213> Homo sapiens
 <220>
 <223> intrabody
 <400> 292
 Gly Gln Ile Phe Ser Arg Ser Ala
 <210> 293
<211> 8
<212> PRT
 <213> Homo sapiens
```

```
<220>
<223> intrabody
<400> 293
Gly Gln Ile His Gly Leu Ser Pro
<210> 294
<211> 8
<212> PRT
<213> Homo sapiens
<220>
<223> intrabody
<400> 294
Gly Ala Arg Ala Ser Val Leu Ser
        5
<210> 295
<211> 8
<212> PRT
<213> Homo sapiens
<220>
<223> intrabody
<400> 295
Gly Cys Thr Leu Ser Ala Glu Glu
<210> 296
<211> 16
<212> PRT
<213> Homo sapiens
<220>
<223> intrabody
Ala Ala Val Ala Leu Leu Pro Ala Val Leu Leu Ala Leu Leu Ala Pro
<210> 297
<211> 12
<212> PRT
<213> Homo sapiens
 <220>
 <223> intrabody
 Ala Ala Val Leu Leu Pro Val Leu Leu Ala Ala Pro
 1 5
 <210> 298
<211> 15
<212> PRT
<213> Homo sapiens
```

```
<220>
<223> intrabody
<400> 298
Val Thr Val Leu Ala Leu Gly Ala Leu Ala Gly Val Gly
<210> 299
<211> 30
<212> DNA
<213> Artificial Sequence
<223> Description of Artificial Sequence: Primer
<400> 299
                                                                    30
ccagcagtac cactteettg cectgegeeg
<210> 300
<211> 30
<212> DNA
<213> Artificial Sequence
<220>
<223> Description of Artificial Sequence: Primer
<400> 300
                                                                    30
gccgcgtccc gttccttcac catgacgacc
<210> 301
<211> 31
<212> DNA
<213> Artificial Sequence
<223> Description of Artificial Sequence: Primer
<400> 301
                                                                    31
ccaqcagtac cgcttccttg ccctgcggcc g
<210> 302
 <211> 30
 <212> DNA
 <213> Artificial Sequence
 <223> Description of Artificial Sequence: Primer
 <400> 302
                                                                     30
 geogegteec gtteetteac catgacgace
 <210> 303
 <211> 450
<212> PRT
<213> Artificial Sequence
 <223> Humanized antibody - VH Chain
 <400> 303
 Gln Val Thr Leu Arg Glu Ser Gly Pro Ala Leu Val Lys Pro Thr Gln
```

```
10
Thr Leu Thr Leu Thr Cys Thr Phe Ser Gly Phe Ser Leu Ser Thr Ala
                    25
Gly Met Ser Val Gly Trp Ile Arg Gln Pro Pro Gly Lys Ala Leu Glu
                       40
Trp Leu Ala Asp Ile Trp Trp Gly Asp Lys Gly His Tyr Asn Pro Ser
Leu Lys Asp Arg Leu Thr Ile Ser Lys Asp Thr Ser Lys Asn Gln Val
                                75
Val Leu Lys Val Thr Asn Met Asp Pro Ala Asp Thr Ala Thr Tyr Tyr
                             90
Cys Ala Arg Asp Met Ile Phe Asn Trp Tyr Phe Asp Val Trp Gly Gln
                          105
Gly Thr Thr Val Thr Val Ser Ser Ala Ser Thr Lys Gly Pro Ser Val
                               125
                       120
Phe Pro Leu Ala Pro Ser Ser Lys Ser Thr Ser Gly Gly Thr Ala Ala
 130 135
                           140
Leu Gly Cys Leu Val Lys Asp Tyr Phe Pro Glu Pro Val Thr Val Ser
                        155 160
                150
Trp Asn Ser Gly Ala Leu Thr Ser Gly Val His Thr Phe Pro Ala Val
             165 170 175
Leu Gln Ser Ser Gly Leu Tyr Ser Leu Ser Ser Val Val Thr Val Pro
                          185
Ser Ser Ser Leu Gly Thr Gln Thr Tyr Ile Cys Asn Val Asn His Lys
                       200
   195
Pro Ser Asn Thr Lys Val Asp Lys Arg Val Glu Pro Lys Ser Cys Asp
       215
Lys Thr His Thr Cys Pro Pro Cys Pro Ala Pro Glu Leu Leu Gly Gly
    230
                                 235
Pro Ser Val Phe Leu Phe Pro Pro Lys Pro Lys Asp Thr Leu Met Ile
                              250 255
             245
Ser Arg Thr Pro Glu Val Thr Cys Val Val Val Asp Val Ser His Glu
          260
                          265
Asp Pro Glu Val Lys Phe Asn Trp Tyr Val Asp Gly Val Glu Val His
                       280
Asn Ala Lys Thr Lys Pro Arg Glu Glu Gln Tyr Asn Ser Thr Tyr Arg
                  295
Val Val Ser Val Leu Thr Val Leu His Gln Asp Trp Leu Asn Gly Lys
       310
Glu Tyr Lys Cys Lys Val Ser Asn Lys Ala Leu Pro Ala Pro Ile Glu
                              330 335
             325
Lys Thr Ile Ser Lys Ala Lys Gly Gln Pro Arg Glu Pro Gln Val Tyr
                           345
          340
Thr Leu Pro Pro Ser Arg Glu Glu Met Thr Lys Asn Gln Val Ser Leu
                        360
 Thr Cys Leu Val Lys Gly Phe Tyr Pro Ser Asp Ile Ala Val Glu Trp
                    375
 Glu Ser Asn Gly Gln Pro Glu Asn Asn Tyr Lys Thr Thr Pro Pro Val
              390
 Leu Asp Ser Asp Gly Ser Phe Phe Leu Tyr Ser Lys Leu Thr Val Asp
             405 410 415
 Lys Ser Arg Trp Gln Gln Gly Asn Val Phe Ser Cys Ser Val Met His
          420 425 430
 Glu Ala Leu His Asn His Tyr Thr Gln Lys Ser Leu Ser Leu Ser Pro
                       440
 Gly Lys
    450
 <210> 304
<211> 120
 <212> PRT
```

```
<213> Artificial Sequence
<220>
<223> Humanized antibody - VH Domain
<400> 304
Gln Val Thr Leu Arg Glu Ser Gly Pro Ala Leu Val Lys Pro Thr Gln
Thr Leu Thr Leu Thr Cys Thr Phe Ser Gly Phe Ser Leu Ser Thr Ala
                               25
Gly Met Ser Val Gly Trp Ile Arg Gln Pro Pro Gly Lys Ala Leu Glu
                                               45
                           40
Trp Leu Ala Asp Ile Trp Trp Gly Asp Lys Gly His Tyr Asn Pro Ser
                      55
Leu Lys Asp Arg Leu Thr Ile Ser Lys Asp Thr Ser Lys Asn Gln Val
                                      75
Val Leu Lys Val Thr Asn Met Asp Pro Ala Asp Thr Ala Thr Tyr Tyr
                               90
Cys Ala Arg Asp Met Ile Phe Asn Trp Tyr Phe Asp Val Trp Gly Gln
                   105
Gly Thr Thr Val Thr Val Ser Ser
       115
<210> 305
<211> 16
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
     antibody and further modified by amino acid
      substitutions
Asp Ile Trp Trp Gly Asp Lys Gly His Tyr Asn Pro Ser Leu Lys Asp
<210> 306
<211> 213
<212> PRT
<213> Artificial Sequence
<220>
<223> Humanized antibody - VL Chain
<400> 306
Asp Ile Gln Met Thr Gln Ser Pro Ser Thr Leu Ser Ala Ser Val Gly
                                    10
Asp Arg Val Thr Ile Thr Cys Ser Ala Ser Ser Ser Val Gly Tyr Met
 His Trp Tyr Gln Gln Lys Pro Gly Lys Ala Pro Lys Leu Leu Ile Tyr
                            40
 Asp Thr Phe Tyr Leu His Ser Gly Val Pro Ser Arg Phe Ser Gly Ser
                        55
 Gly Ser Gly Thr Glu Phe Thr Leu Thr Ile Ser Ser Leu Gln Pro Asp
                                        75
                    70
 Asp Phe Ala Thr Tyr Tyr Cys Phe Gln Gly Ser Gly Tyr Pro Phe Thr
                                    90
 Phe Gly Gly Gly Thr Lys Val Glu Ile Lys Arg Thr Val Ala Ala Pro
                               105
            100
 Ser Val Phe Ile Phe Pro Pro Ser Asp Glu Gln Leu Lys Ser Gly Thr
                            120
```

```
Ala Ser Val Val Cys Leu Leu Asn Asn Phe Tyr Pro Arg Glu Ala Lys
                       135
                                           140
Val Gln Trp Lys Val Asp Asn Ala Leu Gln Ser Gly Asn Ser Gln Glu
                   150
                                       155
Ser Val Thr Glu Gln Asp Ser Lys Asp Ser Thr Tyr Ser Leu Ser Ser
                                   170
Thr Leu Thr Leu Ser Lys Ala Asp Tyr Glu Lys His Lys Val Tyr Ala
                               185
Cys Glu Val Thr His Gln Gly Leu Ser Ser Pro Val Thr Lys Ser Phe
                            200
                                               205
Asn Arg Gly Glu Cys
   210
<210> 307
<211> 106
<212> PRT
<213> Artificial Sequence
<220>
<223> Humanized antibody - VL Domain
<400> 307
Asp Ile Gln Met Thr Gln Ser Pro Ser Thr Leu Ser Ala Ser Val Gly
               5
                                   10
Asp Arg Val Thr Ile Thr Cys Ser Ala Ser Ser Ser Val Gly Tyr Met
His Trp Tyr Gln Gln Lys Pro Gly Lys Ala Pro Lys Leu Leu Ile Tyr
                            40
Asp Thr Phe Tyr Leu His Ser Gly Val Pro Ser Arg Phe Ser Gly Ser
                        55
Gly Ser Gly Thr Glu Phe Thr Leu Thr Ile Ser Ser Leu Gln Pro Asp
                    70
                                       75
Asp Phe Ala Thr Tyr Tyr Cys Phe Gln Gly Ser Gly Tyr Pro Phe Thr
                85
                                   90
Phe Gly Gly Gly Thr Lys Val Glu Ile Lys
<210> 308
<211> 7
<212> PRT
<213> Artificial Sequence
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 308
Asp Thr Phe Tyr Leu His Ser
1
<210> 309
<211> 450
<212> PRT
<213> Artificial Sequence
<220>
<223> Humanized antibody - VH Chain
<400> 309
Gln Val Thr Leu Arg Glu Ser Gly Pro Ala Leu Val Lys Pro Thr Gln
```

```
1.0
Thr Leu Thr Leu Thr Cys Thr Phe Ser Gly Phe Ser Leu Ser Thr Ala
                  25
Gly Met Ser Val Gly Trp Ile Arg Gln Pro Pro Gly Lys Ala Leu Glu
                 40
Trp Leu Ala Asp Ile Trp Trp Asp Asp Lys Lys Ser Tyr Asn Pro Ser
                   55
Leu Lys Asp Arg Leu Thr Ile Ser Lys Asp Thr Ser Lys Asn Gln Val
                                 75
                70
Val Leu Lys Val Thr Asn Met Asp Pro Ala Asp Thr Ala Thr Tyr Tyr
                              90
Cys Ala Arg Asp Met Ile Thr Asn Trp Tyr Phe Asp Val Trp Gly Gln
                 105
Gly Thr Thr Val Thr Val Ser Ser Ala Ser Thr Lys Gly Pro Ser Val
                       120
Phe Pro Leu Ala Pro Ser Ser Lys Ser Thr Ser Gly Gly Thr Ala Ala
                           140
         135
Leu Gly Cys Leu Val Lys Asp Tyr Phe Pro Glu Pro Val Thr Val Ser
                        155 160
       150
Trp Asn Ser Gly Ala Leu Thr Ser Gly Val His Thr Phe Pro Ala Val
             165 170 175
Leu Gln Ser Ser Gly Leu Tyr Ser Leu Ser Ser Val Val Thr Val Pro
                           185
Ser Ser Ser Leu Gly Thr Gln Thr Tyr Ile Cys Asn Val Asn His Lys
                       200
Pro Ser Asn Thr Lys Val Asp Lys Arg Val Glu Pro Lys Ser Cys Asp
                    215
Lys Thr His Thr Cys Pro Pro Cys Pro Ala Pro Glu Leu Leu Gly Gly
                                 235
Pro Ser Val Phe Leu Phe Pro Pro Lys Pro Lys Asp Thr Leu Met Ile
                              250 255
             245
Ser Arg Thr Pro Glu Val Thr Cys Val Val Val Asp Val Ser His Glu
                         265
Asp Pro Glu Val Lys Phe Asn Trp Tyr Val Asp Gly Val Glu Val His
                       280
Asn Ala Lys Thr Lys Pro Arg Glu Glu Gln Tyr Asn Ser Thr Tyr Arg
                                      300
                    295
Val Val Ser Val Leu Thr Val Leu His Gln Asp Trp Leu Asn Gly Lys
                                  315 320
                310
Glu Tyr Lys Cys Lys Val Ser Asn Lys Ala Leu Pro Ala Pro Ile Glu
                               330 335
Lys Thr Ile Ser Lys Ala Lys Gly Gln Pro Arg Glu Pro Gln Val Tyr
                           345
Thr Leu Pro Pro Ser Arg Glu Glu Met Thr Lys Asn Gln Val Ser Leu
                        360
Thr Cys Leu Val Lys Gly Phe Tyr Pro Ser Asp Ile Ala Val Glu Trp
                    375
Glu Ser Asn Gly Gln Pro Glu Asn Asn Tyr Lys Thr Thr Pro Pro Val
               390
Leu Asp Ser Asp Gly Ser Phe Phe Leu Tyr Ser Lys Leu Thr Val Asp
                              410 415
Lys Ser Arg Trp Gln Gln Gly Asn Val Phe Ser Cys Ser Val Met His
          420 425 430
Glu Ala Leu His Asn His Tyr Thr Gln Lys Ser Leu Ser Leu Ser Pro
                       440
Gly Lys
   450
<210> 310
<211> 120
<212> PRT
```

```
<213> Artificial Sequence
<220>
<223> Humanized antibody - VH Domain
<400> 310
Gln Val Thr Leu Arg Glu Ser Gly Pro Ala Leu Val Lys Pro Thr Gln
                                  10
Thr Leu Thr Leu Thr Cys Thr Phe Ser Gly Phe Ser Leu Ser Thr Ala
Gly Met Ser Val Gly Trp Ile Arg Gln Pro Pro Gly Lys Ala Leu Glu
                           40
                                              45
Trp Leu Ala Asp Ile Trp Trp Asp Asp Lys Lys Ser Tyr Asn Pro Ser
                       55
Leu Lys Asp Arg Leu Thr Ile Ser Lys Asp Thr Ser Lys Asn Gln Val
                                       75
                   70
Val Leu Lys Val Thr Asn Met Asp Pro Ala Asp Thr Ala Thr Tyr Tyr
                                  90
               8.5
Cys Ala Arg Asp Met Ile Thr Asn Trp Tyr Phe Asp Val Trp Gly Gln
           100 105
Gly Thr Thr Val Thr Val Ser Ser
    115
<210> 311
<211> 10
<212> PRT
<213> Artificial Sequence
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 311
Asp Met Ile Thr Asn Trp Tyr Phe Asp Val
<210> 312
<211> 213
<212> PRT
<213> Artificial Sequence
<220>
<223> Humanized antibody - VL Chain
<400> 312
Asp Ile Gln Met Thr Gln Ser Pro Ser Thr Leu Ser Ala Ser Val Gly
                                    10
Asp Arg Val Thr Ile Thr Cys Leu Leu Ser Ser Arg Val Gly Tyr Met
                                25
 His Trp Tyr Gln Gln Lys Pro Gly Lys Ala Pro Lys Leu Leu Ile Tyr
                            40
 Asp Thr Tyr Tyr Gln Thr Ser Gly Val Pro Ser Arg Phe Ser Gly Ser
                        55
 Gly Ser Gly Thr Glu Phe Thr Leu Thr Ile Ser Ser Leu Gln Pro Asp
                    70
                                        75
 Asp Phe Ala Thr Tyr Tyr Cys Phe Gln Gly Ser Gly Tyr Pro Phe Thr
                                   90
 Phe Gly Gly Gly Thr Lys Val Glu Ile Lys Arg Thr Val Ala Ala Pro
             100
                                105
 Ser Val Phe Ile Phe Pro Pro Ser Asp Glu Gln Leu Lys Ser Gly Thr
```

```
125
                          120
Ala Ser Val Val Cys Leu Leu Asn Asn Phe Tyr Pro Arg Glu Ala Lys
                              140
                     135
Val Gln Trp Lys Val Asp Asn Ala Leu Gln Ser Gly Asn Ser Gln Glu
                           155
                 150
Ser Val Thr Glu Gln Asp Ser Lys Asp Ser Thr Tyr Ser Leu Ser Ser
              165
                        170
Thr Leu Thr Leu Ser Lys Ala Asp Tyr Glu Lys His Lys Val Tyr Ala
                                      190
          180 185
Cys Glu Val Thr His Gln Gly Leu Ser Ser Pro Val Thr Lys Ser Phe
                         200
      195
Asn Arg Gly Glu Cys
   210
<210> 313
<211> 106
<212> PRT
<213> Artificial Sequence
<220>
<223> Humanized antibody - VL Domain
<400> 313
Asp Ile Gln Met Thr Gln Ser Pro Ser Thr Leu Ser Ala Ser Val Gly
                                  10
Asp Arg Val Thr Ile Thr Cys Leu Leu Ser Ser Arg Val Gly Tyr Met
                              25
His Trp Tyr Gln Gln Lys Pro Gly Lys Ala Pro Lys Leu Leu Ile Tyr
                          40
Asp Thr Tyr Tyr Gln Thr Ser Gly Val Pro Ser Arg Phe Ser Gly Ser
                       55
Gly Ser Gly Thr Glu Phe Thr Leu Thr Ile Ser Ser Leu Gln Pro Asp
                                     75
                  70
Asp Phe Ala Thr Tyr Tyr Cys Phe Gln Gly Ser Gly Tyr Pro Phe Thr
              85
Phe Gly Gly Gly Thr Lys Val Glu Ile Lys
<210> 314
<211> 10
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
Leu Leu Ser Ser Arg Val Gly Tyr Met His
               5
<210> 315
<211> 7
<212> PRT
 <213> Artificial Sequence
 <223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
```

```
<400> 315
Asp Thr Tyr Tyr Gln Thr Ser
<210> 316
<211> 450
<212> PRT
<213> Artificial Sequence
<220>
<223> Humanized antibody - VH Chain
<400> 316
Gln Val Thr Leu Arg Glu Ser Gly Pro Ala Leu Val Lys Pro Thr Gln
Thr Leu Thr Leu Thr Cys Thr Phe Ser Gly Phe Ser Leu Ser Thr Ala
Gly Met Ser Val Gly Trp Ile Arg Gln Pro Pro Gly Lys Ala Leu Glu
Trp Leu Ala Asp Ile Trp Trp Asp Asp Lys Lys His Tyr Asn Pro Ser
                 55
Leu Lys Asp Arg Leu Thr Ile Ser Lys Asp Thr Ser Lys Asn Gln Val
Val Leu Lys Val Thr Asn Met Asp Pro Ala Asp Thr Ala Thr Tyr Tyr
Cys Ala Arg Asp Met Ile Phe Asn Trp Tyr Phe Asp Val Trp Gly Gln
                 105 110
Gly Thr Thr Val Thr Val Ser Ser Ala Ser Thr Lys Gly Pro Ser Val
                     120 125
Phe Pro Leu Ala Pro Ser Ser Lys Ser Thr Ser Gly Gly Thr Ala Ala
                   135 140
Leu Gly Cys Leu Val Lys Asp Tyr Phe Pro Glu Pro Val Thr Val Ser
               150 155
Trp Asn Ser Gly Ala Leu Thr Ser Gly Val His Thr Phe Pro Ala Val
            165 170 175
Leu Gln Ser Ser Gly Leu Tyr Ser Leu Ser Ser Val Val Thr Val Pro
                       185 190
Ser Ser Ser Leu Gly Thr Gln Thr Tyr Ile Cys Asn Val Asn His Lys
           200
Pro Ser Asn Thr Lys Val Asp Lys Arg Val Glu Pro Lys Ser Cys Asp
                   215
Lys Thr His Thr Cys Pro Pro Cys Pro Ala Pro Glu Leu Leu Gly Gly
225 230 235 240
Pro Ser Val Phe Leu Phe Pro Pro Lys Pro Lys Asp Thr Leu Met Ile
                 250 255
Ser Arg Thr Pro Glu Val Thr Cys Val Val Val Asp Val Ser His Glu
         260 265 270
Asp Pro Glu Val Lys Phe Asn Trp Tyr Val Asp Gly Val Glu Val His
                   280
Asn Ala Lys Thr Lys Pro Arg Glu Glu Gln Tyr Asn Ser Thr Tyr Arg
                   295
                          300
Val Val Ser Val Leu Thr Val Leu His Gln Asp Trp Leu Asn Gly Lys
                                315
       310
Glu Tyr Lys Cys Lys Val Ser Asn Lys Ala Leu Pro Ala Pro Ile Glu
             325 330 335
Lys Thr Ile Ser Lys Ala Lys Gly Gln Pro Arg Glu Pro Gln Val Tyr
                         345 350
Thr Leu Pro Pro Ser Arg Glu Glu Met Thr Lys Asn Gln Val Ser Leu
                      360
Thr Cys Leu Val Lys Gly Phe Tyr Pro Ser Asp Ile Ala Val Glu Trp
```

```
375
   370
Glu Ser Asn Gly Gln Pro Glu Asn Asn Tyr Lys Thr Thr Pro Pro Val
               390
                            395
Leu Asp Ser Asp Gly Ser Phe Phe Leu Tyr Ser Lys Leu Thr Val Asp
                       410
              405
Lys Ser Arg Trp Gln Gln Gly Asn Val Phe Ser Cys Ser Val Met His
                   425
Glu Ala Leu His Asn His Tyr Thr Gln Lys Ser Leu Ser Leu Ser Pro
                 440
Gly Lys
  450
<210> 317
<211> 120
<212> PRT
<213> Artificial Sequence
<220>
<223> Humanized antibody - VH Domain
<400> 317
Gln Val Thr Leu Arg Glu Ser Gly Pro Ala Leu Val Lys Pro Thr Gln
                                  10
Thr Leu Thr Leu Thr Cys Thr Phe Ser Gly Phe Ser Leu Ser Thr Ala
                              25
Gly Met Ser Val Gly Trp Ile Arg Gln Pro Pro Gly Lys Ala Leu Glu
                          40
Trp Leu Ala Asp Ile Trp Trp Asp Asp Lys Lys His Tyr Asn Pro Ser
                                         60
Leu Lys Asp Arg Leu Thr Ile Ser Lys Asp Thr Ser Lys Asn Gln Val
                                     75
                   70
Val Leu Lys Val Thr Asn Met Asp Pro Ala Asp Thr Ala Thr Tyr Tyr
                                 90
Cys Ala Arg Asp Met Ile Phe Asn Trp Tyr Phe Asp Val Trp Gly Gln
Gly Thr Thr Val Thr Val Ser Ser
<210> 318
<211> 213
<212> PRT
<213> Artificial Sequence
<220>
<223> Humanized antibody - VL Chain
Asp Ile Gln Met Thr Gln Ser Pro Ser Thr Leu Ser Ala Ser Val Gly
Asp Arg Val Thr Ile Thr Cys Leu Leu Ser Ser Arg Val Gly Tyr Met
His Trp Tyr Gln Gln Lys Pro Gly Lys Ala Pro Lys Leu Leu Ile Tyr
Asp Thr Met Tyr Gln Ala Ser Gly Val Pro Ser Arg Phe Ser Gly Ser
                       55
Gly Ser Gly Thr Glu Phe Thr Leu Thr Ile Ser Ser Leu Gln Pro Asp
Asp Phe Ala Thr Tyr Tyr Cys Phe Gln Gly Ser Gly Tyr Pro Phe Thr
                                  90
               85
Phe Gly Gly Gly Thr Lys Val Glu Ile Lys Arg Thr Val Ala Ala Pro
                              105
```

```
Ser Val Phe Ile Phe Pro Pro Ser Asp Glu Gln Leu Lys Ser Gly Thr
                            120
                                                125
Ala Ser Val Val Cys Leu Leu Asn Asn Phe Tyr Pro Arg Glu Ala Lys
                        135
Val Gln Trp Lys Val Asp Asn Ala Leu Gln Ser Gly Asn Ser Gln Glu
                    150
                                        155
Ser Val Thr Glu Gln Asp Ser Lys Asp Ser Thr Tyr Ser Leu Ser Ser
                165
                                    170
Thr Leu Thr Leu Ser Lys Ala Asp Tyr Glu Lys His Lys Val Tyr Ala
                                185
                                                    190
Cys Glu Val Thr His Gln Gly Leu Ser Ser Pro Val Thr Lys Ser Phe
                            200
Asn Arg Gly Glu Cys
    210
<210> 319
<211> 106
<212> PRT
<213> Artificial Sequence
<220>
<223> Humanized antibody - VL Domain
<400> 319
Asp Ile Gln Met Thr Gln Ser Pro Ser Thr Leu Ser Ala Ser Val Gly
                                    10
Asp Arg Val Thr Ile Thr Cys Leu Leu Ser Ser Arg Val Gly Tyr Met
His Trp Tyr Gln Gln Lys Pro Gly Lys Ala Pro Lys Leu Leu Ile Tyr
        35
                            40
                                                45
Asp Thr Met Tyr Gln Ala Ser Gly Val Pro Ser Arg Phe Ser Gly Ser
                        55
                                            60
Gly Ser Gly Thr Glu Phe Thr Leu Thr Ile Ser Ser Leu Gln Pro Asp
                                        75
Asp Phe Ala Thr Tyr Tyr Cys Phe Gln Gly Ser Gly Tyr Pro Phe Thr
Phe Gly Gly Gly Thr Lys Val Glu Ile Lys
            100
<210> 320
<211> 10
<212> PRT
<213> Artificial Sequence
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 320
Leu Leu Ser Ser Arg Val Gly Tyr Met His
                 5
<210> 321
<211>
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
     antibody and further modified by amino acid
```

substitutions

```
<400> 321
Asp Thr Met Tyr Gln Ala Ser
<210> 322
<211> 450
<212> PRT
<213> Artificial Sequence
<220>
<223> Humanized antibody - VH Chain
<400> 322
Gln Val Thr Leu Arg Glu Ser Gly Pro Ala Leu Val Lys Pro Thr Gln
Thr Leu Thr Leu Thr Cys Thr Phe Ser Gly Phe Ser Leu Ser Thr Ala
Gly Met Ser Val Gly Trp Ile Arg Gln Pro Pro Gly Lys Ala Leu Glu
Trp Leu Ala Asp Ile Trp Trp Asp Asp Lys Lys His Tyr Asn Pro Ser
                       55
Leu Lys Asp Arg Leu Thr Ile Ser Lys Asp Thr Ser Lys Asn Gln Val
                   70
Val Leu Lys Val Thr Asn Met Asp Pro Ala Asp Thr Ala Thr Tyr Tyr
Cys Ala Arg Asp Met Ile Phe Asn Trp Tyr Phe Asp Val Trp Gly Gln
           100
                               105
Gly Thr Thr Val Thr Val Ser Ser Ala Ser Thr Lys Gly Pro Ser Val
                           120
                                              125
Phe Pro Leu Ala Pro Ser Ser Lys Ser Thr Ser Gly Gly Thr Ala Ala
                       135
                                          140
Leu Gly Cys Leu Val Lys Asp Tyr Phe Pro Glu Pro Val Thr Val Ser
                   150
                                       155
Trp Asn Ser Gly Ala Leu Thr Ser Gly Val His Thr Phe Pro Ala Val
               165
                                   170
Leu Gln Ser Ser Gly Leu Tyr Ser Leu Ser Ser Val Val Thr Val Pro
                               185
Ser Ser Ser Leu Gly Thr Gln Thr Tyr Ile Cys Asn Val Asn His Lys
                           200
Pro Ser Asn Thr Lys Val Asp Lys Arg Val Glu Pro Lys Ser Cys Asp
Lys Thr His Thr Cys Pro Pro Cys Pro Ala Pro Glu Leu Leu Gly Gly
                   230
                                       235
Pro Ser Val Phe Leu Phe Pro Pro Lys Pro Lys Asp Thr Leu Met Ile
                                   250
Ser Arg Thr Pro Glu Val Thr Cys Val Val Val Asp Val Ser His Glu
                               265
Asp Pro Glu Val Lys Phe Asn Trp Tyr Val Asp Gly Val Glu Val His
                           280
Asn Ala Lys Thr Lys Pro Arg Glu Glu Gln Tyr Asn Ser Thr Tyr Arg
                       295
Val Val Ser Val Leu Thr Val Leu His Gln Asp Trp Leu Asn Gly Lys
                   310
                                      315
Glu Tyr Lys Cys Lys Val Ser Asn Lys Ala Leu Pro Ala Pro Ile Glu
               325
                                   330 335
Lys Thr Ile Ser Lys Ala Lys Gly Gln Pro Arg Glu Pro Gln Val Tyr
                              345
Thr Leu Pro Pro Ser Arg Glu Glu Met Thr Lys Asn Gln Val Ser Leu
                           360
```

```
Thr Cys Leu Val Lys Gly Phe Tyr Pro Ser Asp Ile Ala Val Glu Trp
                        375
Glu Ser Asn Gly Gln Pro Glu Asn Asn Tyr Lys Thr Thr Pro Pro Val
                    390
                                        395
Leu Asp Ser Asp Gly Ser Phe Phe Leu Tyr Ser Lys Leu Thr Val Asp
                405
                                    410
Lys Ser Arg Trp Gln Gln Gly Asn Val Phe Ser Cys Ser Val Met His
            420
                                425
Glu Ala Leu His Asn His Tyr Thr Gln Lys Ser Leu Ser Leu Ser Pro
                            440
Gly Lys
    450
<210> 323
<211> 120
<212> PRT
<213> Artificial Sequence
<220>
<223> Humanized antibody - VH Domain
<400> 323
Gln Val Thr Leu Arg Glu Ser Gly Pro Ala Leu Val Lys Pro Thr Gln
                                    10
Thr Leu Thr Leu Thr Cys Thr Phe Ser Gly Phe Ser Leu Ser Thr Ala
Gly Met Ser Val Gly Trp Ile Arg Gln Pro Pro Gly Lys Ala Leu Glu
                           4.0
Trp Leu Ala Asp Ile Trp Trp Asp Asp Lys Lys His Tyr Asn Pro Ser
Leu Lys Asp Arg Leu Thr Ile Ser Lys Asp Thr Ser Lys Asn Gln Val
                    70
                                        75
Val Leu Lys Val Thr Asn Met Asp Pro Ala Asp Thr Ala Thr Tyr Tyr
                                    90
Cys Ala Arg Asp Met Ile Phe Asn Trp Tyr Phe Asp Val Trp Gly Gln
                                105
Gly Thr Thr Val Thr Val Ser Ser
<210> 324
<211> 213
<212> PRT
<213> Artificial Sequence
<220>
<223> Humanized antibody - VL Chain
<400> 324
Asp Ile Gln Met Thr Gln Ser Pro Ser Thr Leu Ser Ala Ser Val Gly
                                    10
Asp Arg Val Thr Ile Thr Cys Ser Leu Ser Ser Arg Val Gly Tyr Met
His Trp Tyr Gln Gln Lys Pro Gly Lys Ala Pro Lys Leu Leu Ile Tyr
                            40
Asp Thr Tyr Tyr Leu Pro Ser Gly Val Pro Ser Arg Phe Ser Gly Ser
Gly Ser Gly Thr Glu Phe Thr Leu Thr Ile Ser Ser Leu Gln Pro Asp
                    70
                                        75
Asp Phe Ala Thr Tyr Tyr Cys Phe Gln Gly Ser Gly Tyr Pro Phe Thr
Phe Gly Gly Gly Thr Lys Val Glu Ile Lys Arg Thr Val Ala Ala Pro
```

```
100
                                105
Ser Val Phe Ile Phe Pro Pro Ser Asp Glu Gln Leu Lys Ser Gly Thr
                           120
                                                125
Ala Ser Val Val Cys Leu Leu Asn Asn Phe Tyr Pro Arg Glu Ala Lys
                        135
                                           140
Val Gln Trp Lys Val Asp Asn Ala Leu Gln Ser Gly Asn Ser Gln Glu
                    150
                                        155
Ser Val Thr Glu Gln Asp Ser Lys Asp Ser Thr Tyr Ser Leu Ser Ser
                165
                                    170
Thr Leu Thr Leu Ser Lys Ala Asp Tyr Glu Lys His Lys Val Tyr Ala
                               185
Cys Glu Val Thr His Gln Gly Leu Ser Ser Pro Val Thr Lys Ser Phe
       195
                            200
                                               205
Asn Arg Gly Glu Cys
   210
<210> 325
<211> 106
<212> PRT
<213> Artificial Sequence
<220>
<223> Humanized antibody - VL Domain
<400> 325
Asp Ile Gln Met Thr Gln Ser Pro Ser Thr Leu Ser Ala Ser Val Gly
                                    10
Asp Arg Val Thr Ile Thr Cys Ser Leu Ser Ser Arg Val Gly Tyr Met
                                25
His Trp Tyr Gln Gln Lys Pro Gly Lys Ala Pro Lys Leu Leu Ile Tyr
       35
                            40
Asp Thr Tyr Tyr Leu Pro Ser Gly Val Pro Ser Arg Phe Ser Gly Ser
                        55
Gly Ser Gly Thr Glu Phe Thr Leu Thr Ile Ser Ser Leu Gln Pro Asp
                    70
Asp Phe Ala Thr Tyr Tyr Cys Phe Gln Gly Ser Gly Tyr Pro Phe Thr
               85
                                    90
Phe Gly Gly Thr Lys Val Glu Ile Lys
           100
<210> 326
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
     antibody and further modified by amino acid
     substitutions
<400> 326
Asp Thr Tyr Tyr Leu Pro Ser
<210> 327
<211> 450
<212> PRT
<213> Artificial Sequence
<220>
<223> Humanized antibody - VH Chain
```

<400> 327 Gln Val Thr Leu Arg Glu Ser Gly Pro Ala Leu Val Lys Pro Thr Gln															
1				5					10					15	
			20	Thr				25					30		
_		35		Gly			40					45			
_	50			Ile		55					60				
65				Leu	70					75					80
				Thr 85					90					95	
			100	Met				105					110		
_		115		Thr			120					125			
	130			Pro		135					140				
145				Val	150					155					160
				Ala 165					170					175	
			180					185					190		
		195		Gly			200					205			
	210			Lys		215					220				
225				Cys	230					235					240
				Leu 245					250					255	
	_		260					265					270		
_		275		Lys			280					285			
	290			Lys		295					300				
305					310					315					Lys 320
	_			Lys 325					330	ı				335	
_			340)				345					350		Tyr
		355	,				360					365			Leu -
	370)				375	i				380	1			Trp
385	,				390					395	;				Val 400
	_			405	;				410)				415	
			420)				425	5				430	1	His
Glu	ı Ala	435		s Asr	n His	Туг	Thr 440		ı Lys	s Ser	: Leu	Ser 445		. Ser	Pro
Gl	л Lys 45(

120

```
<210> 328
<211> 120
<212> PRT
<213> Artificial Sequence
<220>
<223> Humanized antibody - VH Domain
<400> 328
Gln Val Thr Leu Arg Glu Ser Gly Pro Ala Leu Val Lys Pro Thr Gln
Thr Leu Thr Leu Thr Cys Thr Phe Ser Gly Phe Ser Leu Ser Thr Ala
Gly Met Ser Val Gly Trp Ile Arg Gln Pro Pro Gly Lys Ala Leu Glu
                           40
Trp Leu Ala Asp Ile Trp Trp Asp Asp Lys Lys Asp Tyr Asn Pro Ser
Leu Lys Asp Arg Leu Thr Ile Ser Lys Asp Thr Ser Lys Asn Gln Val
Val Leu Lys Val Thr Asn Met Asp Pro Ala Asp Thr Ala Thr Tyr Tyr
                                  90
               85
Cys Ala Arg Asp Met Ile Phe Asn Trp Tyr Phe Asp Val Trp Gly Gln
                   105
           100
Gly Thr Thr Val Thr Val Ser Ser
       115
                          120
<210> 329
<211> 16
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
Asp Ile Trp Trp Asp Asp Lys Lys Asp Tyr Asn Pro Ser Leu Lys Asp
<210> 330
 <211> 213
<212> PRT
<213> Artificial Sequence
 <220>
 <223> Humanized antibody - VL Chain
 <400> 330
 Asp Ile Gln Met Thr Gln Ser Pro Ser Thr Leu Ser Ala Ser Val Gly
                                    10
 Asp Arg Val Thr Ile Thr Cys Ser Leu Ser Ser Arg Val Gly Tyr Met
                                25
 His Trp Tyr Gln Gln Lys Pro Gly Lys Ala Pro Lys Leu Leu Ile Tyr
                            40
 Asp Thr Phe Arg His Thr Ser Gly Val Pro Ser Arg Phe Ser Gly Ser
                        55
 Gly Ser Gly Thr Glu Phe Thr Leu Thr Ile Ser Ser Leu Gln Pro Asp
                                       75
 Asp Phe Ala Thr Tyr Tyr Cys Phe Gln Gly Ser Gly Tyr Pro Phe Thr
```

```
Phe Gly Gly Gly Thr Lys Val Glu Ile Lys Arg Thr Val Ala Ala Pro
                            105
          100
Ser Val Phe Ile Phe Pro Pro Ser Asp Glu Gln Leu Lys Ser Gly Thr
                        120
Ala Ser Val Val Cys Leu Leu Asn Asn Phe Tyr Pro Arg Glu Ala Lys
                  135
Val Gln Trp Lys Val Asp Asn Ala Leu Gln Ser Gly Asn Ser Gln Glu
               150
                                    155
Ser Val Thr Glu Gln Asp Ser Lys Asp Ser Thr Tyr Ser Leu Ser Ser
                               170
              165
Thr Leu Thr Leu Ser Lys Ala Asp Tyr Glu Lys His Lys Val Tyr Ala
                     185 190
Cys Glu Val Thr His Gln Gly Leu Ser Ser Pro Val Thr Lys Ser Phe
                         200
Asn Arg Gly Glu Cys
 210
<210> 331
<211> 106
<212> PRT
<213> Artificial Sequence
<220>
<223> Humanized antibody - VL Domain
<400> 331
Asp Ile Gln Met Thr Gln Ser Pro Ser Thr Leu Ser Ala Ser Val Gly
                    10
Asp Arg Val Thr Ile Thr Cys Ser Leu Ser Ser Arg Val Gly Tyr Met
                      25
His Trp Tyr Gln Gln Lys Pro Gly Lys Ala Pro Lys Leu Leu Ile Tyr
                      40
Asp Thr Phe Arg His Thr Ser Gly Val Pro Ser Arg Phe Ser Gly Ser
                     55
Gly Ser Gly Thr Glu Phe Thr Leu Thr Ile Ser Ser Leu Gln Pro Asp
                 70
                      75
Asp Phe Ala Thr Tyr Tyr Cys Phe Gln Gly Ser Gly Tyr Pro Phe Thr
              85
                      90
Phe Gly Gly Thr Lys Val Glu Ile Lys
<210> 332
<211> 7
<212> PRT
<213> Artificial Sequence
<223> Amino acid sequence derived from Murine monoclonal
     antibody and further modified by amino acid
      substitutions
<400> 332
Asp Thr Phe Arg His Thr Ser
<210> 333
<211> 213
<212> PRT
<213> Artificial Sequence
```

```
<220>
<223> Humanized antibody - VL Chain
<400> 333
Asp Ile Gln Met Thr Gln Ser Pro Ser Thr Leu Ser Ala Ser Val Gly
                             10
Asp Arg Val Thr Ile Thr Cys Ser Pro Ser Ser Ser Val Gly Tyr Met
                      25
His Trp Tyr Gln Gln Lys Pro Gly Lys Ala Pro Lys Leu Leu Ile Tyr
                        40
Asp Thr Tyr Tyr Leu Ala Ser Gly Val Pro Ser Arg Phe Ser Gly Ser
Gly Ser Gly Thr Glu Phe Thr Leu Thr Ile Ser Ser Leu Gln Pro Asp
                          75
Asp Phe Ala Thr Tyr Tyr Cys Phe Gln Gly Ser Gly Tyr Pro Phe Thr
                                90
Phe Gly Gly Gly Thr Lys Val Glu Ile Lys Arg Thr Val Ala Ala Pro
                            105
Ser Val Phe Ile Phe Pro Pro Ser Asp Glu Gln Leu Lys Ser Gly Thr
                        120 125
Ala Ser Val Val Cys Leu Leu Asn Asn Phe Tyr Pro Arg Glu Ala Lys
                            140
                     135
Val Gln Trp Lys Val Asp Asn Ala Leu Gln Ser Gly Asn Ser Gln Glu
                          155 160
                 150
Ser Val Thr Glu Gln Asp Ser Lys Asp Ser Thr Tyr Ser Leu Ser Ser
                      170 175
             1.65
Thr Leu Thr Leu Ser Lys Ala Asp Tyr Glu Lys His Lys Val Tyr Ala
          180 185 190
Cys Glu Val Thr His Gln Gly Leu Ser Ser Pro Val Thr Lys Ser Phe
                        200
      195
Asn Arg Gly Glu Cys
  210
<210> 334
<211> 106
<212> PRT
<213> Artificial Sequence
<220>
<223> Humanized antibody - VL Domain
<400> 334
Asp Ile Gln Met Thr Gln Ser Pro Ser Thr Leu Ser Ala Ser Val Gly
Asp Arg Val Thr Ile Thr Cys Ser Pro Ser Ser Ser Val Gly Tyr Met
His Trp Tyr Gln Gln Lys Pro Gly Lys Ala Pro Lys Leu Leu Ile Tyr
                          40
Asp Thr Tyr Tyr Leu Ala Ser Gly Val Pro Ser Arg Phe Ser Gly Ser
                      55
Gly Ser Gly Thr Glu Phe Thr Leu Thr Ile Ser Ser Leu Gln Pro Asp
                                    75
Asp Phe Ala Thr Tyr Tyr Cys Phe Gln Gly Ser Gly Tyr Pro Phe Thr
              85
 Phe Gly Gly Gly Thr Lys Val Glu Ile Lys
           1.00
 <210> 335
 <211> 10
<212> PRT
```

```
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
     antibody and further modified by amino acid
      substitutions
<400> 335
Ser Pro Ser Ser Ser Val Gly Tyr Met His
               5
<210> 336
<211> 7
<212> PRT
<213> Artificial Sequence
<223> Amino acid sequence derived from Murine monoclonal.
      antibody and further modified by amino acid
      substitutions
<400> 336
Asp Thr Tyr Tyr Leu Ala Ser
            5
<210> 337
<211> 365
<212> PRT
<213> Homo sapiens
<220>
<223> Human FcRn
<400> 337
Met Gly Val Pro Arg Pro Gln Pro Trp Ala Leu Gly Leu Leu Phe
                                    10
Leu Leu Pro Gly Ser Leu Gly Ala Glu Ser His Leu Ser Leu Leu Tyr
                                25
His Leu Thr Ala Val Ser Ser Pro Ala Pro Gly Thr Pro Ala Phe Trp
                            40
Val Ser Gly Trp Leu Gly Pro Gln Gln Tyr Leu Ser Tyr Asn Ser Leu
Arg Gly Glu Ala Glu Pro Cys Gly Ala Trp Val Trp Glu Asn Gln Val
                                        75
                    70
Ser Trp Tyr Trp Glu Lys Glu Thr Thr Asp Leu Arg Ile Lys Glu Lys
Leu Phe Leu Glu Ala Phe Lys Ala Leu Gly Gly Lys Gly Pro Tyr Thr
                                105
 Leu Gln Gly Leu Leu Gly Cys Glu Leu Gly Pro Asp Asn Thr Ser Val
                            120
 Pro Thr Ala Lys Phe Ala Leu Asn Gly Glu Glu Phe Met Asn Phe Asp
                                            140
                        135
 Leu Lys Gln Gly Thr Trp Gly Gly Asp Trp Pro Glu Ala Leu Ala Ile
                    150
                                        155
 Ser Gln Arg Trp Gln Gln Gln Asp Lys Ala Ala Asn Lys Glu Leu Thr
                                    170
                165
 Phe Leu Leu Phe Ser Cys Pro His Arg Leu Arg Glu His Leu Glu Arg
                                185
            180
 Gly Arg Gly Asn Leu Glu Trp Lys Glu Pro Pro Ser Met Arg Leu Lys
                            200
                                                205
 Ala Arg Pro Ser Ser Pro Gly Phe Ser Val Leu Thr Cys Ser Ala Phe
```

```
215
Ser Phe Tyr Pro Pro Glu Leu Gln Leu Arg Phe Leu Arg Asn Gly Leu
                                  235
               230
Ala Ala Gly Thr Gly Gln Gly Asp Phe Gly Pro Asn Ser Asp Gly Ser
                               250
             245
Phe His Ala Ser Ser Ser Leu Thr Val Lys Ser Gly Asp Glu His His
         260
                 265
Tyr Cys Cys Ile Val Gln His Ala Gly Leu Ala Gln Pro Leu Arg Val
                       280
Glu Leu Glu Ser Pro Ala Lys Ser Ser Val Leu Val Val Gly Ile Val
   290 295
                                      300
Ile Gly Val Leu Leu Thr Ala Ala Ala Val Gly Gly Ala Leu Leu
                                  315
       310
Trp Arg Arg Met Arg Ser Gly Leu Pro Ala Pro Trp Ile Ser Leu Arg
                               330
             325
Gly Asp Asp Thr Gly Val Leu Leu Pro Thr Pro Gly Glu Ala Gln Asp
         340 345 350
Ala Asp Leu Lys Asp Val Asn Val Ile Pro Ala Thr Ala
                       360
<210> 338
<211> 365
<212> PRT
<213> Murine
<400> 338
Met Gly Met Pro Leu Pro Trp Ala Leu Ser Leu Leu Leu Val Leu Leu
                               10
Pro Gln Thr Trp Gly Ser Glu Thr Arg Pro Pro Leu Met Tyr His Leu
Thr Ala Val Ser Asn Pro Ser Thr Gly Leu Pro Ser Phe Trp Ala Thr
                         40
Gly Trp Leu Gly Pro Gln Gln Tyr Leu Thr Tyr Asn Ser Leu Arg Gln
                     55
Glu Ala Asp Pro Cys Gly Ala Trp Val Trp Glu Asn Gln Val Ser Trp
                 70
                                    75
Tyr Trp Glu Lys Glu Thr Thr Asp Leu Lys Ser Lys Glu Gln Leu Phe
                               90
Leu Glu Ala Leu Lys Thr Leu Glu Lys Ile Leu Asn Gly Thr Tyr Thr
                           105
Leu Gln Gly Leu Leu Gly Cys Glu Leu Ala Ser Asp Asn Ser Ser Val
                        120
Pro Thr Ala Val Phe Ala Leu Asn Gly Glu Glu Phe Met Lys Phe Asn
                     135
                                       140
Pro Arg Ile Gly Asn Trp Thr Gly Glu Trp Pro Glu Thr Glu Ile Val
                                   155 160
                 150
Ala Asn Leu Trp Met Lys Gln Pro Asp Ala Ala Arg Lys Glu Ser Glu
                                170
              165
Phe Leu Leu Asn Ser Cys Pro Glu Arg Leu Leu Gly His Leu Glu Arg
                            185
Gly Arg Arg Asn Leu Glu Trp Lys Glu Pro Pro Ser Met Arg Leu Lys
                         200
Ala Arg Pro Gly Asn Ser Gly Ser Ser Val Leu Thr Cys Ala Ala Phe
                     215
                                       220
Ser Phe Tyr Pro Pro Glu Leu Lys Phe Arg Phe Leu Arg Asn Gly Leu
                 230
                                   235
Ala Ser Gly Ser Gly Asn Cys Ser Thr Gly Pro Asn Gly Asp Gly Ser
                               250
              245
Phe His Ala Trp Ser Leu Leu Glu Val Lys Arg Gly Asp Glu His His
                            265
Tyr Gln Cys Gln Val Glu His Glu Gly Leu Ala Gln Pro Leu Thr Val
```

```
280
Asp Leu Asp Ser Ser Ala Arg Ser Ser Val Pro Val Val Gly Ile Val
                                       300
                  295
Leu Gly Leu Leu Leu Val Val Val Ala Ile Ala Gly Gly Val Leu Leu
                                    315
    310
Trp Gly Arg Met Arg Ser Gly Leu Pro Ala Pro Trp Leu Ser Leu Ser
              325 330 335
Gly Asp Asp Ser Gly Asp Leu Leu Pro Gly Gly Asn Leu Pro Pro Glu
        340 345
Ala Glu Pro Gln Gly Ala Asn Ala Phe Pro Ala Thr Ser
                         360
<210> 339
<211> 110
<212> PRT
<213> Homo sapiens
<400> 339
Ala Pro Glu Leu Leu Gly Gly Pro Ser Val Phe Leu Phe Pro Pro Lys
                                 10
Pro Lys Asp Thr Leu Met Ile Ser Arg Thr Pro Glu Val Thr Cys Val
Val Val Asp Val Ser His Glu Asp Pro Glu Val Lys Phe Asn Trp Tyr
                         40
Val Asp Gly Val Glu Val His Asn Ala Lys Thr Lys Pro Arg Glu Glu
                      55
Gln Tyr Asn Ser Thr Tyr Arg Val Val Ser Val Leu Thr Val Leu His
                                     75
                70
Gln Asp Trp Leu Asn Gly Lys Glu Tyr Lys Cys Lys Val Ser Asn Lys
                                 90
            85
Ala Leu Pro Ala Pro Ile Glu Lys Thr Ile Ser Lys Ala Lys
                             105
<210> 340
<211> 107
<212> PRT
<213> Homo sapiens
<400> 340
Gly Gln Pro Arg Glu Pro Gln Val Tyr Thr Leu Pro Pro Ser Arg Glu
                                 10
Glu Met Thr Lys Asn Gln Val Ser Leu Thr Cys Leu Val Lys Gly Phe
          20
                              25
Tyr Pro Ser Asp Ile Ala Val Glu Trp Glu Ser Asn Gly Gln Pro Glu
                          4.0
Asn Asn Tyr Lys Thr Thr Pro Pro Val Leu Asp Ser Asp Gly Ser Phe
                      55
Phe Leu Tyr Ser Lys Leu Thr Val Asp Lys Ser Arg Trp Gln Gln Gly
                  70
                                     75
Asn Val Phe Ser Cys Ser Val Met His Glu Ala Leu His Asn His Tyr
                                 90
              85
 Thr Gln Lys Ser Leu Ser Leu Ser Pro Gly Lys
          100 105
 <210> 341
 <211> 15
 <212> PRT
 <213> Homo sapiens
 <400> 341
 Glu Pro Lys Ser Cys Asp Lys Thr His Thr Cys Pro Pro Cys Pro
```

```
1
                                10
                                                    15
<210> 342
<211> 232
<212> PRT
<213> Homo sapiens
<220>
<223> Human hinge Fc region
<400> 342
Glu Pro Lys Ser Cys Asp Lys Thr His Thr Cys Pro Pro Cys Pro Ala
Pro Glu Leu Leu Gly Gly Pro Ser Val Phe Leu Phe Pro Pro Lys Pro
Lys Asp Thr Leu Met Ile Ser Arg Thr Pro Glu Val Thr Cys Val Val
Val Asp Val Ser His Glu Asp Pro Glu Val Lys Phe Asn Trp Tyr Val
Asp Gly Val Glu Val His Asn Ala Lys Thr Lys Pro Arg Glu Glu Gln
Tyr Asn Ser Thr Tyr Arg Val Val Ser Val Leu Thr Val Leu His Gln
Asp Trp Leu Asn Gly Lys Glu Tyr Lys Cys Lys Val Ser Asn Lys Ala
                             105
Leu Pro Ala Pro Ile Glu Lys Thr Ile Ser Lys Ala Lys Gly Gln Pro
                         120
Arg Glu Pro Gln Val Tyr Thr Leu Pro Pro Ser Arg Glu Glu Met Thr
                     135
                                        140
Lys Asn Gln Val Ser Leu Thr Cys Leu Val Lys Gly Phe Tyr Pro Ser
                 150
                                  155
Asp Ile Ala Val Glu Trp Glu Ser Asn Gly Gln Pro Glu Asn Asn Tyr
             165
                                170
Lys Thr Thr Pro Pro Val Leu Asp Ser Asp Gly Ser Phe Phe Leu Tyr
          180
                             185
Ser Lys Leu Thr Val Asp Lys Ser Arg Trp Gln Gln Gly Asn Val Phe
                         200 205
Ser Cys Ser Val Met His Glu Ala Leu His Asn His Tyr Thr Gln Lys
  210 215
Ser Leu Ser Leu Ser Pro Gly Lys
<210> 343
<211> 120
<212> PRT
<213> Artificial Sequence
<220>
<223> Humanized antibody - VH Domain
Gln Val Thr Leu Arg Glu Ser Gly Pro Ala Leu Val Lys Pro Thr Gln
                                 10
Thr Leu Thr Leu Thr Cys Thr Phe Ser Gly Phe Ser Leu Ser Thr Ser
Gly Met Ser Val Gly Trp Ile Arg Gln Pro Pro Gly Lys Ala Leu Glu
                          40
Trp Leu Ala Asp Ile Trp Trp Asp Asp Lys Lys Asp Tyr Asn Pro Ser
                      55
Leu Lys Ser Arg Leu Thr Ile Ser Lys Asp Thr Ser Lys Asn Gln Val
```

```
Val Leu Lys Val Thr Asn Met Asp Pro Ala Asp Thr Ala Thr Tyr Tyr
                                    90
               85
Cys Ala Arg Ser Met Ile Thr Asn Trp Tyr Phe Asp Val Trp Gly Gln
                               105
Gly Thr Thr Val Thr Val Ser Ser
        115
<210> 344
<211> 7
<212> PRT
<213> Artificial Sequence
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 344
Val Leu His Gln Asp Trp Leu
<210> 345
<211> 6
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 345
Leu Met Ile Ser Arg Thr
<210> 346
<211> 9
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 346
Met His Glu Ala Leu His Asn His Tyr
<210> 347
<211> 5
<212> PRT
<213> Artificial Sequence
 <223> Amino acid sequence derived from Murine monoclonal
       antibody and further modified by amino acid
       substitutions
 <400> 347
```

```
Gly Gln Pro Glu Asn
<210> 348
<211> 6
<212> PRT
<213> Artificial Sequence
<223> Mutants isolated from Library by panning
<400> 348
Leu Tyr Ile Thr Arg Glu
<210> 349
<211> 6
<212> PRT
<213> Artificial Sequence
<223> Mutants isolated from Library by panning
<400> 349
Leu Tyr Ile Ser Arg Thr
<210> 350
<211> 6
<212> PRT
<213> Artificial Sequence
<223> Mutants isolated from Library by panning
<400> 350
Leu Tyr Ile Ser Arg Ser
<210> 351
<211> 6
<212> PRT
<213> Artificial Sequence
<223> Mutants isolated from Library by panning
<400> 351
Leu Tyr Ile Ser Arg Arg
<210> 352
 <211> 6
 <212> PRT
 <213> Artificial Sequence
 <220>
 <223> Mutants isolated from Library by panning
 <400> 352
 Leu Tyr Ile Ser Arg Gln
```

```
5
1
<210> 353
<211> 6
<212> PRT
<213> Artificial Sequence
<220>
<223> Mutants isolated from Library by panning
Leu Trp Ile Ser Arg Thr
<210> 354
<211> 6
<212> PRT
<213> Artificial Sequence
<223> Mutants isolated from Library by panning
<400> 354
Leu Tyr Ile Ser Leu Gln
<210> 355
<211> 6
<212> PRT
<213> Artificial Sequence
<223> Mutants isolated from Library by panning
<400> 355
Leu Phe Ile Ser Arg Asp
<210> 356
<211> 6
<212> PRT
<213> Artificial Sequence
<223> Mutants isolated from Library by panning
<400> 356
Leu Phe Ile Ser Arg Thr
<210> 357
<211> 6
 <212> PRT
 <213> Artificial Sequence
 <223> Mutants isolated from Library by panning
 <400> 357
 Leu Phe Ile Ser Arg Arg
            5
```

```
<210> 358
<211> 6
<212> PRT
<213> Artificial Sequence
<223> Mutants isolated from Library by panning
<400> 358
Leu Phe Ile Thr Gly Ala
<210> 359
<211> 6
<212> PRT
<213> Artificial Sequence
<223> Mutants isolated from Library by panning
<400> 359
Leu Ser Ile Ser Arg Glu
<210> 360
<211> 6
<212> PRT
<213> Artificial Sequence
<223> Mutants isolated from Library by panning
<400> 360
Arg Thr Ile Ser Ile Ser
<210> 361
<211> 7
<212> PRT
<213> Artificial Sequence
<223> Mutants isolated from Library by panning
<400> 361
Thr Pro His Ser Asp Trp Leu
<210> 362
<211> 7
<212> PRT
<213> Artificial Sequence
<223> Mutants isolated from Library by panning
 <400> 362
 Ile Pro His Glu Asp Trp Leu
```

```
<210> 363
<211> 5
<211> 5
<212> PRT
<213> Artificial Sequence
<220>
<223> Mutants isolated from Library by panning
<400> 363
Arg Thr Arg Glu Pro
<210> 364
<211> 5
<212> PRT
<213> Artificial Sequence
<220>
<223> Mutants isolated from Library by panning
<400> 364
Asp Pro Pro Glu Ser
<210> 365
<211> 5
<212> PRT
<213> Artificial Sequence
<220>
<223> Mutants isolated from Library by panning
<400> 365
Ser Asp Pro Glu Pro
<210> 366
<211> 5
<212> PRT
<213> Artificial Sequence
<223> Mutants isolated from Library by panning
<400> 366
Thr Ser His Glu Asn
<210> 367
<211> 5
 <212> PRT
<213> Artificial Sequence
<223> Mutants isolated from Library by panning
 <400> 367
 Ser Lys Ser Glu Asn
                 5
```

```
<210> 368
<211> 5
<212> PRT
<213> Artificial Sequence
<223> Mutants isolated from Library by panning
<400> 368
His Arg Ser Glu Asn
<210> 369
<211> 5
<211> 5
<212> PRT
<213> Artificial Sequence
<223> Mutants isolated from Library by panning
<400> 369
Lys Ile Arg Glu Asn
<210> 370
<211> 5
<212> PRT
<213> Artificial Sequence
<223> Mutants isolated from Library by panning
<400> 370
Gly Ile Thr Glu Ser
<210> 371
<211> 5
<212> PRT
<213> Artificial Sequence
 <223> Mutants isolated from Library by panning
 <400> 371
 Ser Met Ala Glu Pro
 <210> 372
<211> 9
<212> PRT
 <213> Artificial Sequence
 <223> Mutants isolated from Library by panning
 <400> 372
 Met His Glu Ala Leu Arg Tyr His His
```

```
<210> 373
<211> 9
<212> PRT
<213> Artificial Sequence
<223> Mutants isolated from Library by panning
<400> 373
Met His Glu Ala Leu His Phe His His
<210> 374
<211> 9
<212> PRT
<213> Artificial Sequence
<220>
<223> Mutants isolated from Library by panning
<400> 374
Met His Glu Ala Leu Lys Phe His His
<210> 375
<211> 9
<212> PRT
<213> Artificial Sequence
<220>
<223> Mutants isolated from Library by panning
<400> 375
Met His Glu Ala Leu Ser Tyr His Arg
<210> 376
<211> 9
<212> PRT
<213> Artificial Sequence
<220>
<223> Mutants isolated from Library by panning
<400> 376
Thr His Glu Ala Leu His Tyr His Thr
<210> 377
 <211> 9
 <212> PRT
 <213> Artificial Sequence
 <223> Mutants isolated from Library by panning
 <400> 377
Met His Glu Ala Leu His Tyr His Tyr
                5
 <210> 378
```

```
<211> 54
<212> DNA
<213> Artificial Sequence
<220>
<223> Degenerate oligoes used to combine with
      TAAssDNA template to make Library 1
<220>
<221> misc_feature
<222> (1)...(54)
<223> N = A, C, G, or T; S = G or C
<400> 378
                                                                    54
catgtgacct caggsnnsnn snngatsnns nnggtgtcct tgggttttgg gggg
<210> 379
<211> 53
<212> DNA
<213> Artificial Sequence
<220>
<223> Degenerate oligoes used to combine with
      TAAssDNA template to make Library 2
<220>
<221> misc_feature
<222> (1)...(53)
<223> N = A, C, G, or T; S = G or C
<400> 379
gcacttgtac tccttgccat tsnnccasnn snngtgsnns nnggtgagga cgc
                                                                    53
<210> 380
<211> 38
 <212> DNA
 <213> Artificial Sequence
 <220>
 <223> Degenerate oligoes used to combine with
      TAAssDNA template to make Library 3
 <220>
 <221> misc feature
 <222> (1)...(38)
 <223> N = A, C, G or T; S = G or C
 <400> 380
                                                                     38
 ggtcttgtag ttsnnctcsn nsnnsnnatt gctctccc
 <210> 381
 <211> 53
 <212> DNA
 <213> Artificial Sequence
 <220>
 <223> Degenerate oligoes used to combine with
       TAAssDNA template to make Library 4
 <220>
```

```
<221> misc feature
<222> (1) ... (53)
<223> N = A, C, G or T; S = G or C
<400> 381
ggctcttctg cgtsnngtgs nnsnncagag cctcatgsnn cacggagcat gag
                                                                   53
<210> 382
<211> 45
<212> DNA
<213> Artificial Sequence
<220>
<223> site-directed mutagenesis primer
<400> 382
qcatgtgacc tcaggttccc gagtgatata gagggtgtcc ttggg
                                                                   45
<210> 383
<211> 45
<212> DNA
<213> Artificial Sequence
<220>
<223> site-directed mutagenesis primer
                                                                   45
cccaaggaca ccctctatat cactcgggaa cctgaggtca catgc
<210> 384
<211> 16
<212> PRT
<213> Artificial Sequence
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 384
Asp Ile Trp Trp Asp Asp Lys Gly Asp Tyr Asn Pro Ser Leu Lys Ser
                                    10
<210> 385
<211> 16
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 385
Asp Ile Trp Trp Asp Asp Lys Gly Asp Tyr Asn Pro Ser Leu Lys Asp
<210> 386
```

```
<211> 16
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 386
Asp Ile Trp Trp Asp Asp Lys Gly His Tyr Asn Pro Ser Leu Lys Ser
                        10
<210> 387
<211> 16
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 387
Asp Ile Trp Trp Asp Asp Lys Gly His Tyr Asn Pro Ser Leu Lys Asp
                                    10
<210> 388
<211> 16
<212> PRT
<213> Artificial Sequence
<220>
 <223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
 <400> 388
 Asp Ile Trp Trp Asp Asp Lys Gly Ser Tyr Asn Pro Ser Leu Lys Ser
 <210> 389
 <211> 16
 <212> PRT
 <213> Artificial Sequence
 <220>
 <223> Amino acid sequence derived from Murine monoclonal
       antibody and further modified by amino acid
       substitutions
 <400> 389
 Asp Ile Trp Trp Asp Asp Lys Gly Ser Tyr Asn Pro Ser Leu Lys Asp
                                 10
 <210> 390
 <211> 16
 <212> PRT
 <213> Artificial Sequence
 <220>
```

```
<223> Amino acid sequence derived from Murine monoclonal
     antibody and further modified by amino acid
     substitutions
Asp Ile Trp Trp Asp Gly Lys Gly Asp Tyr Asn Pro Ser Leu Lys Ser
                                  10
<210> 391
<211> 16
<212> PRT
<213> Artificial Sequence
<223> Amino acid sequence derived from Murine monoclonal
     antibody and further modified by amino acid
      substitutions
Asp Ile Trp Trp Asp Gly Lys Gly Asp Tyr Asn Pro Ser Leu Lys Asp
                                   10
<210> 392
<211> 16
<212> PRT
<213> Artificial Sequence
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
Asp Ile Trp Trp Asp Gly Lys Gly His Tyr Asn Pro Ser Leu Lys Ser
                                   10
<210> 393
<211> 16
<212> PRT
<213> Artificial Sequence
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
Asp Ile Trp Trp Asp Gly Lys Gly His Tyr Asn Pro Ser Leu Lys Asp
                     10
                 5
<210> 394
<211> 16
<212> PRT
<213> Artificial Sequence
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 394
```

```
Asp Ile Trp Trp Asp Gly Lys Gly Ser Tyr Asn Pro Ser Leu Lys Ser
     5 10
<210> 395
<211> 16
<212> PRT
<213> Artificial Sequence
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 395
Asp Ile Trp Trp Asp Gly Lys Gly Ser Tyr Asn Pro Ser Leu Lys Asp
                           10
<210> 396
<211> 10
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 396
Lys Cys Gln Leu Phe Val Gly Tyr Met His
<210> 397
<211> 10
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 397
Lys Cys Gln Ser Phe Val Gly Tyr Met His
                5
<210> 398
 <211> 10
 <212> PRT
<213> Artificial Sequence
 <220>
 <223> Amino acid sequence derived from Murine monoclonal
       antibody and further modified by amino acid
       substitutions
 <400> 398
 Lys Cys Gln Val Ser Val Gly Tyr Met His
 <210> 399
```

139

```
<211> 10
<212> PRT
<213> Artificial Sequence
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 399
Lys Cys Gln Val Arg Val Gly Tyr Met His
                 5
<210> 400
<211> 10
<212> PRT
<213> Artificial Sequence
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 400
Lys Cys Gln Val Phe Val Gly Tyr Met His
<210> 401
<211> 10
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 401
Lys Cys Ser Leu Phe Val Gly Tyr Met His
                 5
<210> 402
<211> 10
<212> PRT
<213> Artificial Sequence
<220>
 <223> Amino acid sequence derived from Murine monoclonal
       antibody and further modified by amino acid
       substitutions
 <400> 402
 Lys Cys Ser Ser Phe Val Gly Tyr Met His
 <210> 403
 <211> 10
 <212> PRT
 <213> Artificial Sequence
 <220>
```

```
<223> Amino acid sequence derived from Murine monoclonal
     antibody and further modified by amino acid
     substitutions
<400> 403
Lys Cys Ser Val Ser Val Gly Tyr Met His
<210> 404
<211> 10
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 404
Lys Cys Ser Val Arg Val Gly Tyr Met His
<210> 405
<211> 10
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
Lys Cys Ser Val Phe Val Gly Tyr Met His
                 5
<210> 406
<211> 10
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
Lys Ala Gln Leu Phe Val Gly Tyr Met His
                 5
<210> 407
<211> 10
<212> PRT
<213> Artificial Sequence
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 407
```

```
Lys Ala Gln Ser Phe Val Gly Tyr Met His
       5
<210> 408
<211> 10
<212> PRT
<213> Artificial Sequence
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 408
Lys Ala Gln Val Ser Val Gly Tyr Met His
<210> 409
<211> 10
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 409
Lys Ala Gln Val Arg Val Gly Tyr Met His
<210> 410
<211> 10
<212> PRT
<213> Artificial Sequence
<2200>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 410
Lys Ala Gln Val Phe Val Gly Tyr Met His
 <210> 411
 <211> 10
 <212> PRT
 <213> Artificial Sequence
 <220>
 <223> Amino acid sequence derived from Murine monoclonal
       antibody and further modified by amino acid
       substitutions
 <400> 411
 Lys Ala Ser Leu Phe Val Gly Tyr Met His
 <210> 412
 <211> 10
```

```
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
Lys Ala Ser Ser Phe Val Gly Tyr Met His
                5
<210> 413
<211> 10
<212> PRT
<213> Artificial Sequence
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
Lys Ala Ser Val Ser Val Gly Tyr Met His
                 5
<210> 414
<211> 10
<212> PRT
<213> Artificial Sequence
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
       substitutions
<400> 414
Lys Ala Ser Val Arg Val Gly Tyr Met His
<210> 415
 <211> 10
 <212> PRT
 <213> Artificial Sequence
 <223> Amino acid sequence derived from Murine monoclonal
       antibody and further modified by amino acid
       substitutions
 <400> 415
 Lys Ala Ser Val Phe Val Gly Tyr Met His
                 5
 <210> 416
 <211> 10
 <212> PRT
 <213> Artificial Sequence
 <223> Amino acid sequence derived from Murine monoclonal
```

antibody and further modified by amino acid substitutions <400> 416 Lys Leu Gln Leu Phe Val Gly Tyr Met His <210> 417 <211> 10 <212> PRT <213> Artificial Sequence <223> Amino acid sequence derived from Murine monoclonal antibody and further modified by amino acid substitutions <400> 417 Lys Leu Gln Ser Phe Val Gly Tyr Met His 5 <210> 418 <211> 10 <212> PRT <213> Artificial Sequence <220> <223> Amino acid sequence derived from Murine monoclonal antibody and further modified by amino acid substitutions <400> 418 Lys Leu Gln Val Ser Val Gly Tyr Met His <210> 419 <211> 10 <212> PRT <213> Artificial Sequence <220> <223> Amino acid sequence derived from Murine monoclonal antibody and further modified by amino acid substitutions <400> 419 Lys Leu Gln Val Arg Val Gly Tyr Met His <210> 420 <211> 10 <212> PRT <213> Artificial Sequence <220> <223> Amino acid sequence derived from Murine monoclonal antibody and further modified by amino acid

<400> 420 Lys Leu Gln Val Phe Val Gly Tyr Met His

substitutions

```
1
                  5
                                    10
<210> 421
<211> 10
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 421
Lys Leu Ser Leu Phe Val Gly Tyr Met His
                 5
<210> 422
<211> 10
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 422
Lys Leu Ser Ser Phe Val Gly Tyr Met His
                 5
<210> 423
<211> 10
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 423
Lys Leu Ser Val Ser Val Gly Tyr Met His
<210> 424
<211> 10
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 424
Lys Leu Ser Val Arg Val Gly Tyr Met His
                 5
<210> 425
<211> 10
```

```
<212> PRT
<213> Artificial Sequence
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 425
Lys Leu Ser Val Phe Val Gly Tyr Met His
<210> 426
<211> 10
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 426
Lys Pro Gln Leu Phe Val Gly Tyr Met His
                 5
<210> 427
<211> 10
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
       substitutions
 <400> 427
 Lys Pro Gln Ser Phe Val Gly Tyr Met His
 <210> 428
 <211> 10
 <212> PRT
 <213> Artificial Sequence
 <220>
 <223> Amino acid sequence derived from Murine monoclonal
       antibody and further modified by amino acid
       substitutions
 <400> 428
 Lys Pro Gln Val Ser Val Gly Tyr Met His
                 5
 <210> 429
 <211> 10
 <212> PRT
 <213> Artificial Sequence
 <223> Amino acid sequence derived from Murine monoclonal
```

antibody and further modified by amino acid substitutions <400> 429 Lys Pro Gln Val Arg Val Gly Tyr Met His 5 <210> 430 <211> 10 <212> PRT <213> Artificial Sequence <220> <223> Amino acid sequence derived from Murine monoclonal antibody and further modified by amino acid substitutions <400> 430 Lys Pro Gln Val Phe Val Gly Tyr Met His <210> 431 <211> 10 <212> PRT <213> Artificial Sequence <220> <223> Amino acid sequence derived from Murine monoclonal antibody and further modified by amino acid substitutions <400> 431 Lys Pro Ser Leu Phe Val Gly Tyr Met His 5 <210> 432 <211> 10 <212> PRT <213> Artificial Sequence <220> <223> Amino acid sequence derived from Murine monoclonal antibody and further modified by amino acid substitutions <400> 432 Lys Pro Ser Ser Phe Val Gly Tyr Met His <210> 433 <211> 10 <212> PRT <213> Artificial Sequence <220> <223> Amino acid sequence derived from Murine monoclonal antibody and further modified by amino acid substitutions

147

<400> 433

```
Lys Pro Ser Val Ser Val Gly Tyr Met His
                5
<210> 434
<211> 10
<212> PRT
<213> Artificial Sequence
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 434
Lys Pro Ser Val Arg Val Gly Tyr Met His
                5
<210> 435
<211> 10
<212> PRT
<213> Artificial Sequence
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 435
{
m Lys} Pro Ser Val Phe Val Gly Tyr Met His
                 5
<210> 436
<211> 10
<212> PRT
<213> Artificial Sequence
 <223> Amino acid sequence derived from Murine monoclonal
       antibody and further modified by amino acid
       substitutions
 <400> 436
 Ser Cys Gln Leu Phe Val Gly Tyr Met His
 <210> 437
 <211> 10
 <212> PRT
 <213> Artificial Sequence
 <220>
 <223> Amino acid sequence derived from Murine monoclonal
       antibody and further modified by amino acid
       substitutions
 <400> 437
 Ser Cys Gln Ser Phe Val Gly Tyr Met His
                  5
 <210> 438
 <211> 10
```

```
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
     antibody and further modified by amino acid
     substitutions
Ser Cys Gln Val Ser Val Gly Tyr Met His
<210> 439
<211> 10
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 439
Ser Cys Gln Val Arg Val Gly Tyr Met His
<210> 440
<211> 10
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
Ser Cys Gln Val Phe Val Gly Tyr Met His
                 5
<210> 441
<211> 10
<212> PRT
<213> Artificial Sequence
 <220>
 <223> Amino acid sequence derived from Murine monoclonal
       antibody and further modified by amino acid
       substitutions
 Ser Cys Ser Leu Phe Val Gly Tyr Met His
                 5
 <210> 442
 <211> 10
 <212> PRT
 <213> Artificial Sequence
 <223> Amino acid sequence derived from Murine monoclonal
```

antibody and further modified by amino acid substitutions

```
<400> 442
Ser Cys Ser Ser Phe Val Gly Tyr Met His
                 5
<210> 443
<211> 10
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 443
Ser Cys Ser Val Ser Val Gly Tyr Met His
     5
<210> 444
<211> 10
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 444
Ser Cys Ser Val Arg Val Gly Tyr Met His
            5
<210> 445
<211> 10
<212> PRT
<213> Artificial Sequence
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 445
Ser Cys Ser Val Phe Val Gly Tyr Met His
<210> 446
<211> 10
<212> PRT
<213> Artificial Sequence
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 446
Ser Ala Gln Leu Phe Val Gly Tyr Met His
```

150

```
10
 1
                 5
<210> 447
<211> 10
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
Ser Ala Gln Ser Phe Val Gly Tyr Met His
<210> 448
<211> 10
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
Ser Ala Gln Val Ser Val Gly Tyr Met His
                 5
<210> 449
<211> 10
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
 Ser Ala Gln Val Arg Val Gly Tyr Met His
                 5
 <210> 450
 <211> 10
 <212> PRT
 <213> Artificial Sequence
 <220>
 <223> Amino acid sequence derived from Murine monoclonal
       antibody and further modified by amino acid
       substitutions
 Ser Ala Gln Val Phe Val Gly Tyr Met His
                 5
 <210> 451
 <211> 10
 <212> PRT
```

```
<213> Artificial Sequence
<223> Amino acid sequence derived from Murine monoclonal
     antibody and further modified by amino acid
     substitutions
<400> 451
Ser Ala Ser Leu Phe Val Gly Tyr Met His
                5
<210> 452
<211> 10
<212> PRT
<213> Artificial Sequence
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
Ser Ala Ser Ser Phe Val Gly Tyr Met His
                 5
<210> 453
<211> 10
<212> PRT
<213> Artificial Sequence
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 453
Ser Ala Ser Val Ser Val Gly Tyr Met His
<210> 454
<211> 10
 <212> PRT
<213> Artificial Sequence
 <220>
 <223> Amino acid sequence derived from Murine monoclonal
       antibody and further modified by amino acid
       substitutions
 <400> 454
 Ser Ala Ser Val Arg Val Gly Tyr Met His
                  5
 <210> 455
 <211> 10
 <212> PRT
 <213> Artificial Sequence
 <223> Amino acid sequence derived from Murine monoclonal
```

antibody and further modified by amino acid substitutions

```
<400> 455
Ser Ala Ser Val Phe Val Gly Tyr Met His
                5
<210> 456
<211> 10
<212> PRT
<213> Artificial Sequence
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 456
Ser Leu Gln Leu Phe Val Gly Tyr Met His
                5
<210> 457
<211> 10
<212> PRT
<213> Artificial Sequence
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 457
Ser Leu Gln Ser Phe Val Gly Tyr Met His
                 5
<210> 458
<211> 10
<212> PRT
<213> Artificial Sequence
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
       substitutions
<400> 458
Ser Leu Gln Val Ser Val Gly Tyr Met His
                 5
<210> 459
 <211> 10
 <212> PRT
 <213> Artificial Sequence
 <220>
 <223> Amino acid sequence derived from Murine monoclonal
       antibody and further modified by amino acid
       substitutions
 <400> 459
 Ser Leu Gln Val Arg Val Gly Tyr Met His
```

```
5
                                    10
1
<210> 460
<211> 10
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
Ser Leu Gln Val Phe Val Gly Tyr Met His
                 5
<210> 461
<211> 10
<212> PRT
<213> Artificial Sequence
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
Ser Leu Ser Leu Phe Val Gly Tyr Met His
                 5
<210> 462
<211> 10
<212> PRT
<213> Artificial Sequence
<223> Amino acid sequence derived from Murine monoclonal
       antibody and further modified by amino acid
       substitutions
<400> 462
Ser Leu Ser Ser Phe Val Gly Tyr Met His
 <210> 463
 <211> 10
 <212> PRT
 <213> Artificial Sequence
 <223> Amino acid sequence derived from Murine monoclonal
       antibody and further modified by amino acid
       substitutions
 <400> 463
 Ser Leu Ser Val Ser Val Gly Tyr Met His
                  5
 <210> 464
 <211> 10
 <212> PRT
```

```
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 464
Ser Leu Ser Val Arg Val Gly Tyr Met His
               5
<210> 465
<211> 10
<212> PRT
<213> Artificial Sequence
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 465
Ser Leu Ser Val Phe Val Gly Tyr Met His
                 5
<210> 466
<211> 10
<212> PRT
<213> Artificial Sequence
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
       substitutions
<400> 466
Ser Pro Gln Leu Phe Val Gly Tyr Met His
<210> 467
<211> 10
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
       antibody and further modified by amino acid
       substitutions
 <400> 467
 Ser Pro Gln Ser Phe Val Gly Tyr Met His
 <210> 468
 <211> 10
 <212> PRT
 <213> Artificial Sequence
 <220>
 <223> Amino acid sequence derived from Murine monoclonal
       antibody and further modified by amino acid
```

substitutions

```
<400> 468
Ser Pro Gln Val Ser Val Gly Tyr Met His
<210> 469
<211> 10
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 469
Ser Pro Gln Val Arg Val Gly Tyr Met His
                5
<210> 470
<211> 10
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 470
Ser Pro Gln Val Phe Val Gly Tyr Met His
                5
<210> 471
<211> 10
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
       substitutions
 <400> 471
 Ser Pro Ser Leu Phe Val Gly Tyr Met His
                 5
 <210> 472
 <211> 10
 <212> PRT
 <213> Artificial Sequence
 <220>
 <223> Amino acid sequence derived from Murine monoclonal
       antibody and further modified by amino acid
       substitutions
 Ser Pro Ser Ser Phe Val Gly Tyr Met His
  1
```

```
<210> 473
<211> 10
<212> PRT
<213> Artificial Sequence
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 473
Ser Pro Ser Val Ser Val Gly Tyr Met His
               5
<210> 474
<211> 10
<212> PRT
<213> Artificial Sequence
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 474
Ser Pro Ser Val Arg Val Gly Tyr Met His
<210> 475
<211> 10
<212> PRT
<213> Artificial Sequence
<223> Amino acid sequence derived from Murine monoclonal
       antibody and further modified by amino acid
       substitutions
 <400> 475
 Ser Pro Ser Val Phe Val Gly Tyr Met His
 <210> 476
 <211> 10
 <212> PRT
 <213> Artificial Sequence
 <223> Amino acid sequence derived from Murine monoclonal
       antibody and further modified by amino acid
       substitutions
 <400> 476
 Leu Cys Gln Leu Phe Val Gly Tyr Met His
 <210> 477
 <211> 10
 <212> PRT
 <213> Artificial Sequence
```

```
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
Leu Cys Gln Ser Phe Val Gly Tyr Met His
<210> 478
<211> 10
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 478
Leu Cys Gln Val Ser Val Gly Tyr Met His
<210> 479
<211> 10
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
Leu Cys Gln Val Arg Val Gly Tyr Met His
<210> 480
<211> 10
<212> PRT
<213> Artificial Sequence
 <220>
 <223> Amino acid sequence derived from Murine monoclonal
       antibody and further modified by amino acid
       substitutions
 <400> 480
 Leu Cys Gln Val Phe Val Gly Tyr Met His
                 5
 <210> 481
 <211> 10
 <212> PRT
 <213> Artificial Sequence
 <223> Amino acid sequence derived from Murine monoclonal
       antibody and further modified by amino acid
       substitutions
```

```
<400> 481
Leu Cys Ser Leu Phe Val Gly Tyr Met His
      5
<210> 482
<211> 10
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
Leu Cys Ser Ser Phe Val Gly Tyr Met His
<210> 483
<211> 10
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
Leu Cys Ser Val Ser Val Gly Tyr Met His
                5
<210> 484
<211> 10
 <212> PRT
 <213> Artificial Sequence
 <220>
 <223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
       substitutions
 Leu Cys Ser Val Arg Val Gly Tyr Met His
 <210> 485
 <211> 10
 <212> PRT
 <213> Artificial Sequence
 <220>
 <223> Amino acid sequence derived from Murine monoclonal
       antibody and further modified by amino acid
       substitutions
 Leu Cys Ser Val Phe Val Gly Tyr Met His
                 5
```

```
<210> 486
<211> 10
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 486
Leu Ala Gln Leu Ser Val Gly Tyr Met His
     5
<210> 487
<211> 10
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 487
Leu Ala Gln Leu Arg Val Gly Tyr Met His
      5
<210> 488
<211> 10
<212> PRT
<213> Artificial Sequence
 <223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
 <400> 488
 Leu Ala Gln Leu Phe Val Gly Tyr Met His
            5
 <210> 489
 <211> 10
 <212> PRT
 <213> Artificial Sequence
 <223> Amino acid sequence derived from Murine monoclonal
       antibody and further modified by amino acid
       substitutions
 <400> 489
 Leu Ala Gln Ser Ser Val Gly Tyr Met His
 <210> 490
 <211> 10
 <212> PRT
 <213> Artificial Sequence
```

```
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 490
Leu Ala Gln Ser Arg Val Gly Tyr Met His
           5
<210> 491
<211> 10
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 491
Leu Ala Gln Ser Phe Val Gly Tyr Met His
                5
<210> 492
<211> 10
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
Leu Ala Gln Val Ser Val Gly Tyr Met His
              5
<210> 493
<211> 10
<212> PRT
<213> Artificial Sequence
 <220>
 <223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
       substitutions
 Leu Ala Gln Val Arg Val Gly Tyr Met His
                5
 <210> 494
 <211> 10
 <212> PRT
 <213> Artificial Sequence
 <220>
 <223> Amino acid sequence derived from Murine monoclonal
       antibody and further modified by amino acid
       substitutions
```

```
<400> 494
Leu Ala Gln Val Phe Val Gly Tyr Met His
      5
<210> 495
<211> 10
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 495
Leu Ala Ser Leu Ser Val Gly Tyr Met His
    5
<210> 496
<211> 10
<212> PRT
<213> Artificial Sequence
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 496
Leu Ala Ser Leu Arg Val Gly Tyr Met His
           5
<210> 497
<211> 10
<212> PRT
<213> Artificial Sequence
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
 <400> 497
Leu Ala Ser Leu Phe Val Gly Tyr Met His
                5
 <210> 498
 <211> 10
 <212> PRT
 <213> Artificial Sequence
 <223> Amino acid sequence derived from Murine monoclonal
       antibody and further modified by amino acid
       substitutions
 <400> 498
 Leu Ala Ser Ser Ser Val Gly Tyr Met His
                 5
```

```
<210> 499
<211> 10
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
Leu Ala Ser Ser Arg Val Gly Tyr Met His
                5
<210> 500
<211> 10
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
Leu Ala Ser Ser Phe Val Gly Tyr Met His
                 5
<210> 501
<211> 10
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
 Leu Ala Ser Val Ser Val Gly Tyr Met His
                 5
 <210> 502
 <211> 10
 <212> PRT
 <213> Artificial Sequence
 <220>
 <223> Amino acid sequence derived from Murine monoclonal
       antibody and further modified by amino acid
       substitutions
 Leu Ala Ser Val Arg Val Gly Tyr Met His
                  5
 <210> 503
 <211> 10
 <212> PRT
 <213> Artificial Sequence
```

```
<220>
<223> Amino acid sequence derived from Murine monoclonal
     antibody and further modified by amino acid
     substitutions
<400> 503
Leu Ala Ser Val Phe Val Gly Tyr Met His
      5
<210> 504
<211> 10
<212> PRT
<213> Artificial Sequence
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 504
Leu Leu Gln Leu Ser Val Gly Tyr Met His
           5
<210> 505
<211> 10
<212> PRT
<213> Artificial Sequence
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 505
Leu Leu Gln Leu Arg Val Gly Tyr Met His
            5
<210> 506
<211> 10
<212> PRT
<213> Artificial Sequence
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
 <400> 506
Leu Leu Gln Leu Phe Val Gly Tyr Met His
                5
 <210> 507
 <211> 10
 <212> PRT
 <213> Artificial Sequence
 <220>
 <223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
       substitutions
```

```
<400> 507
Leu Leu Gln Ser Ser Val Gly Tyr Met His
<210> 508
<211> 10
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 508
Leu Leu Gln Ser Arg Val Gly Tyr Met His
<210> 509
<211> 10
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 509
Leu Leu Gln Ser Phe Val Gly Tyr Met His
                 5
<210> 510
<211> 10
<212> PRT
<213> Artificial Sequence
 <220>
 <223> Amino acid sequence derived from Murine monoclonal
       antibody and further modified by amino acid
       substitutions
 Leu Leu Gln Val Ser Val Gly Tyr Met His
                 5
 <210> 511
 <211> 10
 <212> PRT
 <213> Artificial Sequence
 <220>
 <223> Amino acid sequence derived from Murine monoclonal
       antibody and further modified by amino acid
       substitutions
 Leu Leu Gln Val Arg Val Gly Tyr Met His
                  5
  1
 <210> 512
```

```
<211> 10
<212> PRT
<213> Artificial Sequence
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 512
Leu Leu Gln Val Phe Val Gly Tyr Met His
               5
<210> 513
<211> 10
<212> PRT
<213> Artificial Sequence
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 513
Leu Leu Ser Leu Ser Val Gly Tyr Met His
     5
<210> 514
<211> 10
<212> PRT
<213> Artificial Sequence
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
 <400> 514
Leu Leu Ser Leu Arg Val Gly Tyr Met His
 <210> 515
 <211> 10
 <212> PRT
 <213> Artificial Sequence
 <220>
 <223> Amino acid sequence derived from Murine monoclonal
       antibody and further modified by amino acid
       substitutions
 <400> 515
 Leu Leu Ser Leu Phe Val Gly Tyr Met His
 <210> 516
 <211> 10
 <212> PRT
 <213> Artificial Sequence
 <220>
```

```
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 516
Leu Leu Ser Ser Ser Val Gly Tyr Met His
                5
<210> 517
<211> 10
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
Leu Leu Ser Ser Arg Val Gly Tyr Met His
                 5
<210> 518
<211> 10
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
Leu Leu Ser Ser Phe Val Gly Tyr Met His
<210> 519
<211> 10
<212> PRT
<213> Artificial Sequence
<220>
 <223> Amino acid sequence derived from Murine monoclonal
       antibody and further modified by amino acid
       substitutions
 <400> 519
 Leu Leu Ser Val Ser Val Gly Tyr Met His
 <210> 520
 <211> 10
 <212> PRT
 <213> Artificial Sequence
 <223> Amino acid sequence derived from Murine monoclonal
       antibody and further modified by amino acid
       substitutions
```

```
<400> 520
Leu Leu Ser Val Arg Val Gly Tyr Met His
             5
<210> 521
<211> 10
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
Leu Leu Ser Val Phe Val Gly Tyr Met His
<210> 522
<211> 10
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
Leu Pro Gln Leu Phe Val Gly Tyr Met His
<210> 523
<211> 10
<212> PRT
<213> Artificial Sequence
 <220>
 <223> Amino acid sequence derived from Murine monoclonal
       antibody and further modified by amino acid
       substitutions
 Leu Pro Gln Ser Phe Val Gly Tyr Met His
                 5
  1
 <210> 524
 <211> 10
 <212> PRT
 <213> Artificial Sequence
 <220>
 <223> Amino acid sequence derived from Murine monoclonal
       antibody and further modified by amino acid
       substitutions
 Leu Pro Gln Val Ser Val Gly Tyr Met His
                                      10
                  5
```

```
<21.0> 525
<211> 10
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 525
Leu Pro Gln Val Arg Val Gly Tyr Met His
                5
<210> 526
<211> 10
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 526
Leu Pro Gln Val Phe Val Gly Tyr Met His
                5
<210> 527
<211> 10
<212> PRT
<213> Artificial Sequence
<220>
 <223> Amino acid sequence derived from Murine monoclonal
       antibody and further modified by amino acid
       substitutions
 Leu Pro Ser Leu Phe Val Gly Tyr Met His
                  5
 <210> 528
 <211> 10
 <212> PRT
 <213> Artificial Sequence
 <220>
 <223> Amino acid sequence derived from Murine monoclonal
       antibody and further modified by amino acid
       substitutions
 Leu Pro Ser Ser Phe Val Gly Tyr Met His
                  5
 <210> 529
 <211> 10
 <212> PRT
 <213> Artificial Sequence
```

```
<220>
<223> Amino acid sequence derived from Murine monoclonal
     antibody and further modified by amino acid
     substitutions
<400> 529
Leu Pro Ser Val Ser Val Gly Tyr Met His
             5
<210> 530
<211> 10
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 530
Leu Pro Ser Val Arg Val Gly Tyr Met His
             5
<210> 531
<211> 10
<212> PRT
<213> Artificial Sequence
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 531
Leu Pro Ser Val Phe Val Gly Tyr Met His
            5
<210> 532
<211> 7
<212> PRT
<213> Artificial Sequence
 <223> Amino acid sequence derived from Murine monoclonal
       antibody and further modified by amino acid
       substitutions
 <400> 532
 Asp Thr Ser Lys Leu Lys Ser
                5
 <210> 533
 <211> 7
 <212> PRT
 <213> Artificial Sequence
 <223> Amino acid sequence derived from Murine monoclonal
       antibody and further modified by amino acid
       substitutions
```

```
<400> 533
Asp Thr Ser Lys Leu Arg Ser
<210> 534
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 534
Asp Thr Ser Lys Leu His Ser
<210> 535
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 535
Asp Thr Ser Lys Leu Thr Ser
<210> 536
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 536
Asp Thr Ser Lys His Ala Ser
 1
                5
 <210> 537
 <211> 7
 <212> PRT
 <213> Artificial Sequence
 <220>
 <223> Amino acid sequence derived from Murine monoclonal
       antibody and further modified by amino acid
       substitutions
 <400> 537
 Asp Thr Ser Lys His Ser Ser
  1
 <210> 538
```

```
<211> 7
<212> PRT
<213> Artificial Sequence
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 538
Asp Thr Ser Lys His Lys Ser
<210> 539
<211> 7
<212> PRT
<213> Artificial Sequence
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 539
Asp Thr Ser Lys His Arg Ser
<210> 540
 <211> 7
 <212> PRT
 <213> Artificial Sequence
 <223> Amino acid sequence derived from Murine monoclonal
       antibody and further modified by amino acid
       substitutions
 <400> 540
 Asp Thr Ser Lys His His Ser
 <210> 541
 <211> 7
 <212> PRT
 <213> Artificial Sequence
 <223> Amino acid sequence derived from Murine monoclonal
       antibody and further modified by amino acid
       substitutions
 <400> 541
 Asp Thr Ser Lys His Pro Ser
 <210> 542
 <211> 7
 <212> PRT
 <213> Artificial Sequence
 <220>
```

```
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 542
Asp Thr Ser Lys His Thr Ser
<210> 543
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 543
Asp Thr Ser Lys His Asp Ser
<210> 544
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
       antibody and further modified by amino acid
       substitutions
<400> 544
Asp Thr Ser Lys Gln Lys Ser
<210> 545
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
 <223> Amino acid sequence derived from Murine monoclonal
       antibody and further modified by amino acid
       substitutions
 <400> 545
 Asp Thr Ser Lys Gln Arg Ser
 <210> 546
 <211> 7
 <212> PRT
 <213> Artificial Sequence
 <220>
 <223> Amino acid sequence derived from Murine monoclonal
       antibody and further modified by amino acid
       substitutions
 <400> 546
```

```
Asp Thr Ser Lys Gln His Ser
               5
1
<210> 547
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 547
Asp Thr Ser Lys Gln Pro Ser
             5
<210> 548
<211> 7
<212> PRT
<213> Artificial Sequence
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 548
Asp Thr Ser Lys Gln Thr Ser
           5
<210> 549
<211> 7
<212> PRT
<213> Artificial Sequence
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
       substitutions
 <400> 549
 Asp Thr Ser Lys Gln Asp Ser
 <210> 550
 <211> 7
 <212> PRT
 <213> Artificial Sequence
 <223> Amino acid sequence derived from Murine monoclonal
       antibody and further modified by amino acid
       substitutions
 <400> 550
 Asp Thr Ser Gly Leu Ser Ser
  1
 <210> 551
 <211> 7
```

```
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 551
Asp Thr Ser Gly Leu Lys Ser
<210> 552
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 552
Asp Thr Ser Gly Leu Arg Ser
<210> 553
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
       antibody and further modified by amino acid
       substitutions
 <400> 553
 Asp Thr Ser Gly Leu His Ser
                 5
 <210> 554
 <211> 7
 <212> PRT
 <213> Artificial Sequence
 <220>
 <223> Amino acid sequence derived from Murine monoclonal
       antibody and further modified by amino acid
       substitutions
 <400> 554
 Asp Thr Ser Gly Leu Thr Ser
  1
 <210> 555
 <211> 7
 <212> PRT
 <213> Artificial Sequence
 <223> Amino acid sequence derived from Murine monoclonal
```

antibody and further modified by amino acid substitutions

```
<400> 555
Asp Thr Ser Gly Leu Asp Ser
<210> 556
<211> 7
<212> PRT
<213> Artificial Sequence
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 556
Asp Thr Ser Gly His Ala Ser
<210> 557
<211> 7
<212> PRT
<213> Artificial Sequence
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
       substitutions
<400> 557
Asp Thr Ser Gly His Ser Ser
<210> 558
 <211> 7
 <212> PRT
 <213> Artificial Sequence
 <220>
 <223> Amino acid sequence derived from Murine monoclonal
       antibody and further modified by amino acid
       substitutions
 <400> 558
 Asp Thr Ser Gly His Lys Ser
 <210> 559
 <211> 7
 <212> PRT
 <213> Artificial Sequence
 <220>
 <223> Amino acid sequence derived from Murine monoclonal
       antibody and further modified by amino acid
       substitutions
 <400> 559
 Asp Thr Ser Gly His Arg Ser
```

```
5
1
<210> 560
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 560
Asp Thr Ser Gly His His Ser
<210> 561
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 561
Asp Thr Ser Gly His Pro Ser
<210> 562
<211> 7
 <212> PRT
 <213> Artificial Sequence
 <220>
 <223> Amino acid sequence derived from Murine monoclonal
       antibody and further modified by amino acid
       substitutions
 <400> 562
 Asp Thr Ser Gly His Thr Ser
 <210> 563
 <211> 7
 <212> PRT
 <213> Artificial Sequence
 <220>
 <223> Amino acid sequence derived from Murine monoclonal
       antibody and further modified by amino acid
       substitutions
 <400> 563
 Asp Thr Ser Gly His Asp Ser
  1
 <210> 564
 <211> 7
 <212> PRT
```

```
<213> Artificial Sequence
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 564
Asp Thr Ser Gly Gln Ala Ser
<210> 565
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 565
Asp Thr Ser Gly Gln Ser Ser
<210> 566
<211> 7
<212> PRT
 <213> Artificial Sequence
 <223> Amino acid sequence derived from Murine monoclonal
       antibody and further modified by amino acid
       substitutions
 <400> 566
 Asp Thr Ser Gly Gln Lys Ser
 <210> 567
 <211> 7
 <212> PRT
 <213> Artificial Sequence
 <220>
 <223> Amino acid sequence derived from Murine monoclonal
       antibody and further modified by amino acid
       substitutions
 <400> 567
 Asp Thr Ser Gly Gln Arg Ser
 <210> 568
 <211> 7
 <212> PRT
 <213> Artificial Sequence
 <220>
 <223> Amino acid sequence derived from Murine monoclonal
        antibody and further modified by amino acid
```

substitutions <400> 568 Asp Thr Ser Gly Gln His Ser 5 <210> 569 <211> 7 <212> PRT <213> Artificial Sequence <223> Amino acid sequence derived from Murine monoclonal antibody and further modified by amino acid substitutions <400> 569 Asp Thr Ser Gly Gln Pro Ser <210> 570 <211> 7 <212> PRT <213> Artificial Sequence <223> Amino acid sequence derived from Murine monoclonal antibody and further modified by amino acid substitutions <400> 570 Asp Thr Ser Gly Gln Thr Ser <210> 571 <211> 7 <212> PRT <213> Artificial Sequence <223> Amino acid sequence derived from Murine monoclonal antibody and further modified by amino acid substitutions <400> 571 Asp Thr Ser Gly Gln Asp Ser <210> 572

<210> 572 <211> 7 <212> PRT <213> Artificial Sequence <220>

<223> Amino acid sequence derived from Murine monoclonal antibody and further modified by amino acid substitutions

<400> 572 Asp Thr Ser Arg Leu Ser Ser

```
5
1
<210> 573
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 573
Asp Thr Ser Arg Leu Lys Ser
<210> 574
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 574
Asp Thr Ser Arg Leu Arg Ser
        5
<210> 575
<211> 7
 <212> PRT
 <213> Artificial Sequence
 <223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
       substitutions
 <400> 575
 Asp Thr Ser Arg Leu His Ser
               5
 <210> 576
 <211> 7
 <212> PRT
 <213> Artificial Sequence
 <220>
 <223> Amino acid sequence derived from Murine monoclonal
       antibody and further modified by amino acid
       substitutions
 <400> 576
 Asp Thr Ser Arg Leu Pro Ser
                 5
 <210> 577
 <211> 7
```

```
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 577
Asp Thr Ser Arg Leu Thr Ser
<210> 578
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 578
Asp Thr Ser Arg Leu Asp Ser
                 5
<210> 579
<211> 7
<212> PRT
 <213> Artificial Sequence
 <220>
 <223> Amino acid sequence derived from Murine monoclonal
       antibody and further modified by amino acid
       substitutions
 <400> 579
 Asp Thr Ser Arg His Ala Ser
                 5
 <210> 580
 <211> 7
 <212> PRT
 <213> Artificial Sequence
 <220>
 <223> Amino acid sequence derived from Murine monoclonal
       antibody and further modified by amino acid
       substitutions
 <400> 580
 Asp Thr Ser Arg His Ser Ser
 <210> 581
 <211> 7
 <212> PRT
 <213> Artificial Sequence
 <223> Amino acid sequence derived from Murine monoclonal
```

antibody and further modified by amino acid substitutions

```
<400> 581
Asp Thr Ser Arg His Lys Ser
<210> 582
<211> 7
<212> PRT
<213> Artificial Sequence
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 582
Asp Thr Ser Arg His Arg Ser
<210> 583
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 583
Asp Thr Ser Arg His His Ser
 <210> 584
 <211> 7
 <212> PRT
 <213> Artificial Sequence
 <223> Amino acid sequence derived from Murine monoclonal
       antibody and further modified by amino acid
       substitutions
 <400> 584
 Asp Thr Ser Arg His Pro Ser
                 5
 <210> 585
 <211> 7
 <212> PRT
 <213> Artificial Sequence
 <220>
 <223> Amino acid sequence derived from Murine monoclonal
       antibody and further modified by amino acid
       substitutions
 <400> 585
 Asp Thr Ser Arg His Thr Ser
```

```
1
                 5
<210> 586
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
     substitutions
<400> 586
Asp Thr Ser Arg His Asp Ser
<210> 587
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
 <400> 587
Asp Thr Ser Arg Gln Ala Ser
 <210> 588
 <211> 7
 <212> PRT
 <213> Artificial Sequence
 <223> Amino acid sequence derived from Murine monoclonal
       antibody and further modified by amino acid
       substitutions
 <400> 588
 Asp Thr Ser Arg Gln Ser Ser
 <210> 589
 <211> 7
 <212> PRT
 <213> Artificial Sequence
 <223> Amino acid sequence derived from Murine monoclonal
       antibody and further modified by amino acid
       substitutions
 <400> 589
 Asp Thr Ser Arg Gln Lys Ser
 <210> 590
 <211> 7
 <212> PRT
```

```
<213> Artificial Sequence
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 590
Asp Thr Ser Arg Gln Arg Ser
<210> 591
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 591
Asp Thr Ser Arg Gln His Ser
<210> 592
<211> 7
<212> PRT
<213> Artificial Sequence
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
       substitutions
 <400> 592
Asp Thr Ser Arg Gln Pro Ser
 <210> 593
 <211> 7
 <212> PRT
 <213> Artificial Sequence
 <220>
 <223> Amino acid sequence derived from Murine monoclonal
       antibody and further modified by amino acid
       substitutions
 <400> 593
 Asp Thr Ser Arg Gln Thr Ser
                 5
 <210> 594
 <211> 7
 <212> PRT
 <213> Artificial Sequence
 <220>
 <223> Amino acid sequence derived from Murine monoclonal
       antibody and further modified by amino acid
```

substitutions <400> 594 Asp Thr Ser Arg Gln Asp Ser <210> 595 <211> 7 <212> PRT <213> Artificial Sequence <220> <223> Amino acid sequence derived from Murine monoclonal antibody and further modified by amino acid substitutions <400> 595 Asp Thr Ser Tyr Leu Lys Ser <210> 596 <211> 7 <212> PRT <213> Artificial Sequence <220> <223> Amino acid sequence derived from Murine monoclonal antibody and further modified by amino acid substitutions <400> 596 Asp Thr Ser Tyr Leu Arg Ser <210> 597 <211> 7 <212> PRT <213> Artificial Sequence <220> <223> Amino acid sequence derived from Murine monoclonal antibody and further modified by amino acid substitutions <400> 597 Asp Thr Ser Tyr Leu His Ser <210> 598 <211> 7 <212> PRT <213> Artificial Sequence <220>

<223> Amino acid sequence derived from Murine monoclonal antibody and further modified by amino acid substitutions

<400> 598
Asp Thr Ser Tyr Leu Pro Ser
1 5

```
<210> 599
<211> 7
<212> PRT
<213> Artificial Sequence
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 599
Asp Thr Ser Tyr Leu Thr Ser
             5
<210> 600
<211> 7
<212> PRT
<213> Artificial Sequence
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 600
Asp Thr Ser Tyr Leu Asp Ser
                5
<210> 601
<211> 7
 <212> PRT
 <213> Artificial Sequence
 <223> Amino acid sequence derived from Murine monoclonal
       antibody and further modified by amino acid
       substitutions
 <400> 601
 Asp Thr Ser Tyr His Ala Ser
 <210> 602
 <211> 7
 <212> PRT
 <213> Artificial Sequence
 <223> Amino acid sequence derived from Murine monoclonal
       antibody and further modified by amino acid
       substitutions
 <400> 602
 Asp Thr Ser Tyr His Ser Ser
 <210> 603
 <211> 7
 <212> PRT
 <213> Artificial Sequence
```

```
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 603
Asp Thr Ser Tyr His Lys Ser
<210> 604
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 604
Asp Thr Ser Tyr His Arg Ser
<210> 605
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 605
Asp Thr Ser Tyr His His Ser
<210> 606
 <211> 7
 <212> PRT
 <213> Artificial Sequence
 <220>
 <223> Amino acid sequence derived from Murine monoclonal
       antibody and further modified by amino acid
       substitutions
 <400> 606
 Asp Thr Ser Tyr His Pro Ser
 <210> 607
 <211> 7
 <212> PRT
 <213> Artificial Sequence
 <223> Amino acid sequence derived from Murine monoclonal
       antibody and further modified by amino acid
       substitutions
```

```
<400> 607
Asp Thr Ser Tyr His Thr Ser
     5
<210> 608
<211> 7
<212> PRT
<213> Artificial Sequence
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 608
Asp Thr Ser Tyr His Asp Ser
<210> 609
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
       antibody and further modified by amino acid
       substitutions
 <400> 609
 Asp Thr Ser Tyr Gln Lys Ser
 1
 <210> 610
 <211> 7
 <212> PRT
 <213> Artificial Sequence
 <223> Amino acid sequence derived from Murine monoclonal
       antibody and further modified by amino acid
       substitutions
 <400> 610
 Asp Thr Ser Tyr Gln Arg Ser
 <210> 611
 <211> 7
 <212> PRT
 <213> Artificial Sequence
 <223> Amino acid sequence derived from Murine monoclonal
       antibody and further modified by amino acid
       substitutions
 <400> 611
 Asp Thr Ser Tyr Gln His Ser
  1
```

```
<210> 612
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 612
Asp Thr Ser Tyr Gln Pro Ser
                 5
<210> 613
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 613
Asp Thr Ser Tyr Gln Thr Ser
<210> 614
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
 <223> Amino acid sequence derived from Murine monoclonal
       antibody and further modified by amino acid
       substitutions
 <400> 614
 Asp Thr Ser Tyr Gln Asp Ser
 <210> 615
 <211> 7
 <212> PRT
 <213> Artificial Sequence
 <220>
 <223> Amino acid sequence derived from Murine monoclonal
       antibody and further modified by amino acid
       substitutions
 <400> 615
 Asp Thr Ser Phe Leu Ala Ser
 <210> 616
 <211> 7
 <212> PRT
 <213> Artificial Sequence
```

```
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 616
Asp Thr Ser Phe Leu Ser Ser
             5
<210> 617
<211> 7
<212> PRT
<213> Artificial Sequence
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 617
Asp Thr Ser Phe Leu Lys Ser
           5
<210> 618
<211> 7
<212> PRT
<213> Artificial Sequence
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
 <400> 618
 Asp Thr Ser Phe Leu Arg Ser
 <210> 619
 <211> 7
 <212> PRT
 <213> Artificial Sequence
 <223> Amino acid sequence derived from Murine monoclonal
       antibody and further modified by amino acid
       substitutions
 <400> 619
 Asp Thr Ser Phe Leu His Ser
 <210> 620
 <211> 7
 <212> PRT
 <213> Artificial Sequence
 <223> Amino acid sequence derived from Murine monoclonal
       antibody and further modified by amino acid
       substitutions
```

```
<400> 620
Asp Thr Ser Phe Leu Pro Ser
           5
<210> 621
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 621
Asp Thr Ser Phe Leu Thr Ser
               5
<210> 622
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 622
Asp Thr Ser Phe His Ala Ser
<210> 623
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
 <223> Amino acid sequence derived from Murine monoclonal
       antibody and further modified by amino acid
       substitutions
 <400> 623
 Asp Thr Ser Phe His Ser Ser
                5
 <210> 624
 <211> 7
 <212> PRT
 <213> Artificial Sequence
 <220>
 <223> Amino acid sequence derived from Murine monoclonal
       antibody and further modified by amino acid
       substitutions
 <400> 624
 Asp Thr Ser Phe His Lys Ser
  1
 <210> 625
```

```
<211> 7
<212> PRT
<213> Artificial Sequence
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 625
Asp Thr Ser Phe His Arg Ser
<210> 626
<211> 7
<212> PRT
<213> Artificial Sequence
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 626
Asp Thr Ser Phe His His Ser
<210> 627
<211> 7
<212> PRT
<213> Artificial Sequence
<223> Amino acid sequence derived from Murine monoclonal
       antibody and further modified by amino acid
       substitutions
 <400> 627
 Asp Thr Ser Phe His Pro Ser
 <210> 628
 <211> 7
 <212> PRT
 <213> Artificial Sequence
 <220>
 <223> Amino acid sequence derived from Murine monoclonal
       antibody and further modified by amino acid
       substitutions
 <400> 628
 Asp Thr Ser Phe His Thr Ser
 <210> 629
 <211> 7
 <212> PRT
 <213> Artificial Sequence
 <220>
```

```
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
     substitutions
<400> 629
Asp Thr Ser Phe His Asp Ser
<210> 630
<211> 7
<212> PRT
<213> Artificial Sequence
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 630
Asp Thr Ser Phe Gln Ala Ser
<210> 631
<211> 7
<212> PRT
<213> Artificial Sequence
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
       substitutions
<400> 631
Asp Thr Ser Phe Gln Ser Ser
 <210> 632
 <211> 7
 <212> PRT
 <213> Artificial Sequence
 <223> Amino acid sequence derived from Murine monoclonal
       antibody and further modified by amino acid
       substitutions
 <400> 632
 Asp Thr Ser Phe Gln Lys Ser
 <210> 633
 <211> 7
 <212> PRT
 <213> Artificial Sequence
 <223> Amino acid sequence derived from Murine monoclonal
       antibody and further modified by amino acid
       substitutions
```

```
<400> 633
Asp Thr Ser Phe Gln Arg Ser
<210> 634
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 634
Asp Thr Ser Phe Gln His Ser
<210> 635
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
 <400> 635
 Asp Thr Ser Phe Gln Pro Ser
                 5
 1
 <210> 636
 <211> 7
 <212> PRT
 <213> Artificial Sequence
 <220>
 <223> Amino acid sequence derived from Murine monoclonal
       antibody and further modified by amino acid
       substitutions
 <400> 636
 Asp Thr Ser Phe Gln Thr Ser
 <210> 637
 <211> 7
 <212> PRT
 <213> Artificial Sequence
 <220>
 <223> Amino acid sequence derived from Murine monoclonal
       antibody and further modified by amino acid
       substitutions
 <400> 637
 Asp Thr Ser Phe Gln Asp Ser
  1
 <210> 638
```

```
<211> 7
<212> PRT
<213> Artificial Sequence
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 638
Asp Thr Ser Leu Leu Ser Ser
<210> 639
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 639
Asp Thr Ser Leu Leu Lys Ser
<210> 640
<211> 7
 <212> PRT
 <213> Artificial Sequence
 <220>
 <223> Amino acid sequence derived from Murine monoclonal
       antibody and further modified by amino acid
       substitutions
 <400> 640
 Asp Thr Ser Leu Leu Arg Ser
 <210> 641
 <211> 7
 <212> PRT
 <213> Artificial Sequence
 <220>
 <223> Amino acid sequence derived from Murine monoclonal
       antibody and further modified by amino acid
       substitutions
 <400> 641
 Asp Thr Ser Leu Leu His Ser
 <210> 642
 <211> 7
 <212> PRT
 <213> Artificial Sequence
```

```
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 642
Asp Thr Ser Leu Leu Pro Ser
      5
<210> 643
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 643
Asp Thr Ser Leu Leu Thr Ser
       5
<210> 644
<211> 7
<212> PRT
<213> Artificial Sequence
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 644
Asp Thr Ser Leu His Ala Ser
<210> 645
<211> 7
<212> PRT
<213> Artificial Sequence
<223> Amino acid sequence derived from Murine monoclonal
       antibody and further modified by amino acid
       substitutions
 <400> 645
 Asp Thr Ser Leu His Ser Ser
 <210> 646
 <211> 7
 <212> PRT
 <213> Artificial Sequence
 <220>
 <223> Amino acid sequence derived from Murine monoclonal
       antibody and further modified by amino acid
       substitutions
```

```
<400> 646
Asp Thr Ser Leu His Lys Ser
<210> 647
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 647
Asp Thr Ser Leu His Arg Ser
<210> 648
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 648
Asp Thr Ser Leu His His Ser
                 5
<210> 649
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 649
Asp Thr Ser Leu His Pro Ser
                 5
<210> 650
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 650
Asp Thr Ser Leu His Thr Ser
 1
<210> 651
```

```
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
     antibody and further modified by amino acid
     substitutions
<400> 651
Asp Thr Ser Leu His Asp Ser
<210> 652
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 652
Asp Thr Ser Leu Gln Ala Ser
<210> 653
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 653
Asp Thr Ser Leu Gln Ser Ser
<210> 654
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 654
Asp Thr Ser Leu Gln Lys Ser
<210> 655
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
```

```
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 655
Asp Thr Ser Leu Gln Arg Ser
<210> 656
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 656
Asp Thr Ser Leu Gln His Ser
<210> 657
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 657
Asp Thr Ser Leu Gln Pro Ser
<210> 658
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
     antibody and further modified by amino acid
      substitutions
<400> 658
Asp Thr Ser Leu Gln Thr Ser
                5
<210> 659
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
     antibody and further modified by amino acid
     substitutions
<400> 659
```

```
Asp Thr Ser Leu Gln Asp Ser
1 5
<210> 660
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
     antibody and further modified by amino acid
   substitutions
<400> 660
Asp Thr Phe Lys Leu Lys Ser
1
               5
<210> 661
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
     antibody and further modified by amino acid
     substitutions
<400> 661
Asp Thr Phe Lys Leu Arg Ser
               5
<210> 662
<211> 7
<212> PRT
<213> Artificial Sequence
<223> Amino acid sequence derived from Murine monoclonal
     antibody and further modified by amino acid
     substitutions
<400> 662
Asp Thr Phe Lys Leu His Ser
               5
<210> 663
<211> 7
<212> PRT
<213> Artificial Sequence
<223> Amino acid sequence derived from Murine monoclonal
     antibody and further modified by amino acid
     substitutions
<400> 663
Asp Thr Phe Lys Leu Pro Ser
1
               5
<210> 664
<211> 7
```

```
<212> PRT
<213> Artificial Sequence
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 664
Asp Thr Phe Lys Leu Thr Ser
<210> 665
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 665
Asp Thr Phe Lys Leu Asp Ser
<210> 666
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 666
Asp Thr Phe Lys His Ala Ser
<210> 667
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 667
Asp Thr Phe Lys His Ser Ser
<210> 668
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
```

substitutions <400> 668 Asp Thr Phe Lys His Lys Ser 1 5 <210> 669 <211> 7 <212> PRT <213> Artificial Sequence <220> <223> Amino acid sequence derived from Murine monoclonal antibody and further modified by amino acid substitutions <400> 669 Asp Thr Phe Lys His Arg Ser 1 5 <210> 670 <211> 7 <212> PRT <213> Artificial Sequence <223> Amino acid sequence derived from Murine monoclonal antibody and further modified by amino acid substitutions <400> 670 Asp Thr Phe Lys His His Ser 1 5 <210> 671 <211> 7 <212> PRT <213> Artificial Sequence <223> Amino acid sequence derived from Murine monoclonal antibody and further modified by amino acid substitutions <400> 671 Asp Thr Phe Lys His Pro Ser 1 5 <210> 672 <211> 7 <212> PRT <213> Artificial Sequence <223> Amino acid sequence derived from Murine monoclonal antibody and further modified by amino acid substitutions <400> 672 Asp Thr Phe Lys His Thr Ser

antibody and further modified by amino acid

```
1
                 5
<210> 673
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 673
Asp Thr Phe Lys His Asp Ser
<210> 674
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 674
Asp Thr Phe Lys Gln Ala Ser
                 5
<210> 675
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 675
Asp Thr Phe Lys Gln Ser Ser
<210> 676
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 676
Asp Thr Phe Lys Gln Lys Ser
<210> 677
<211> 7
<212> PRT
```

```
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
     substitutions
<400> 677
Asp Thr Phe Lys Gln Arg Ser
1
<210> 678
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 678
Asp Thr Phe Lys Gln His Ser
1
                5
<210> 679
<211> 7
<212> PRT
<213> Artificial Sequence
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 679
Asp Thr Phe Lys Gln Pro Ser
                5
 1
<210> 680
<211> 7
<212> PRT
<213> Artificial Sequence
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 680
Asp Thr Phe Lys Gln Thr Ser
 1
                5
<210> 681
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
```

substitutions <400> 681 Asp Thr Phe Lys Gln Asp Ser 1. <210> 682 <211> 7 <212> PRT <213> Artificial Sequence <220> <223> Amino acid sequence derived from Murine monoclonal antibody and further modified by amino acid substitutions <400> 682 Asp Thr Phe Gly Leu Ala Ser 1 <210> 683 <211> 7 <212> PRT <213> Artificial Sequence <220> <223> Amino acid sequence derived from Murine monoclonal antibody and further modified by amino acid substitutions <400> 683 Asp Thr Phe Gly Leu Ser Ser 1 <210> 684 <211> 7 <212> PRT <213> Artificial Sequence <220> <223> Amino acid sequence derived from Murine monoclonal antibody and further modified by amino acid substitutions <400> 684 Asp Thr Phe Gly Leu Lys Ser 1 <210> 685 <211> 7 <212> PRT <213> Artificial Sequence <220> <223> Amino acid sequence derived from Murine monoclonal antibody and further modified by amino acid substitutions <400> 685 Asp Thr Phe Gly Leu Arg Ser 1

```
<210> 686
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 686
Asp Thr Phe Gly Leu His Ser
1
                 5
<210> 687
<211> 7
<212> PRT
<213> Artificial Sequence
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 687
Asp Thr Phe Gly Leu Pro Ser
 1
                 5
<210> 688
<211> 7
<212> PRT
<213> Artificial Sequence
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 688
Asp Thr Phe Gly Leu Thr Ser
                 5
<210> 689
<211> 7
<212> PRT
<213> Artificial Sequence
<223> Amino acid sequence derived from Murine monoclonal
       antibody and further modified by amino acid
      substitutions
<400> 689
Asp Thr Phe Gly Leu Asp Ser
<210> 690
<211> 7
<212> PRT
<213> Artificial Sequence
```

```
<220>
<223> Amino acid sequence derived from Murine monoclonal
     antibody and further modified by amino acid
     substitutions
<400> 690
Asp Thr Phe Gly His Ala Ser
<210> 691
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 691
Asp Thr Phe Gly His Ser Ser
1
<210> 692
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 692
Asp Thr Phe Gly His Lys Ser
1
<210> 693
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 693
Asp Thr Phe Gly His Arg Ser
 1
<210> 694
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
```

```
<400> 694
Asp Thr Phe Gly His His Ser
1
<210> 695
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 695
Asp Thr Phe Gly His Pro Ser
                 5
<210> 696
<211> 7
<212> PRT
<213> Artificial Sequence
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 696
Asp Thr Phe Gly His Thr Ser
 1
                 5
<210> 697
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 697
Asp Thr Phe Gly His Asp Ser
<210> 698
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 698
Asp Thr Phe Gly Gln Ala Ser
                5
```

```
<210> 699
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 699
Asp Thr Phe Gly Gln Ser Ser
                 5
<210> 700
<211> 7
<212> PRT
<213> Artificial Sequence
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 700
Asp Thr Phe Gly Gln Lys Ser
                 5
<210> 701
<211> 7
<212> PRT
<213> Artificial Sequence
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 701
Asp Thr Phe Gly Gln Arg Ser
<210> 702
<211> 7
<212> PRT
<213> Artificial Sequence
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 702
Asp Thr Phe Gly Gln His Ser
<210> 703
<211> 7
<212> PRT
```

```
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 703
Asp Thr Phe Gly Gln Pro Ser
1
                 5
<210> 704
<211> 7
<212> PRT
<213> Artificial Sequence
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 704
Asp Thr Phe Gly Gln Thr Ser
                 5
<210> 705
<211> 7
<212> PRT
<213> Artificial Sequence
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 705
Asp Thr Phe Gly Gln Asp Ser
 1
<210> 706
<211> 7
<212> PRT
<213> Artificial Sequence
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 706
Asp Thr Phe Arg Leu Ala Ser
 1
                 5
<210> 707
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
```

substitutions <400> 707 Asp Thr Phe Arg Leu Ser Ser 1 <210> 708 <211> 7 <212> PRT <213> Artificial Sequence <220> <223> Amino acid sequence derived from Murine monoclonal antibody and further modified by amino acid substitutions <400> 708 Asp Thr Phe Arg Leu Lys Ser 5 <210> 709 <211> 7 <212> PRT <213> Artificial Sequence <220> <223> Amino acid sequence derived from Murine monoclonal antibody and further modified by amino acid substitutions <400> 709 Asp Thr Phe Arg Leu Arg Ser <210> 710 <211> 7 <212> PRT <213> Artificial Sequence <220> <223> Amino acid sequence derived from Murine monoclonal antibody and further modified by amino acid substitutions <400> 710 Asp Thr Phe Arg Leu His Ser <210> 711 <211> 7 <212> PRT <213> Artificial Sequence <220> <223> Amino acid sequence derived from Murine monoclonal antibody and further modified by amino acid substitutions <400> 711 Asp Thr Phe Arg Leu Pro Ser

```
<210> 712
<211> 7
<212> PRT
<213> Artificial Sequence
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 712
Asp Thr Phe Arg Leu Thr Ser
                5
<210> 713
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 713
Asp Thr Phe Arg Leu Asp Ser
<210> 714
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 714
Asp Thr Phe Arg His Ala Ser
       5
<210> 715
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
       antibody and further modified by amino acid
       substitutions
<400> 715
Asp Thr Phe Arg His Ser Ser
         5
 <210> 716
 <211> 7
 <212> PRT
 <213> Artificial Sequence
```

```
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 716
Asp Thr Phe Arg His Lys Ser
<210> 717
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 717
Asp Thr Phe Arg His Arg Ser
<210> 718
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 718
Asp Thr Phe Arg His His Ser
                5
<210> 719
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
       antibody and further modified by amino acid
       substitutions
<400> 719
Asp Thr Phe Arg His Pro Ser
<210> 720
 <211> 7
 <212> PRT
<213> Artificial Sequence
 <220>
 <223> Amino acid sequence derived from Murine monoclonal
       antibody and further modified by amino acid
       substitutions
```

```
<400> 720
Asp Thr Phe Arg His Thr Ser
       5
<210> 721
<211> 7
<212> PRT
<213> Artificial Sequence
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 721
Asp Thr Phe Arg His Asp Ser
       5
<210> 722
<211> 7
<212> PRT
<213> Artificial Sequence
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 722
Asp Thr Phe Arg Gln Ala Ser
<210> 723
<211> 7
 <212> PRT
 <213> Artificial Sequence
 <223> Amino acid sequence derived from Murine monoclonal
       antibody and further modified by amino acid
       substitutions
 <400> 723
 Asp Thr Phe Arg Gln Ser Ser
 <210> 724
 <211> 7
 <212> PRT
 <213> Artificial Sequence
 <223> Amino acid sequence derived from Murine monoclonal
       antibody and further modified by amino acid
       substitutions
 <400> 724
 Asp Thr Phe Arg Gln Lys Ser
```

```
<210> 725
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 725
Asp Thr Phe Arg Gln Arg Ser
<210> 726
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 726
Asp Thr Phe Arg Gln His Ser
<210> 727
<211> 7
<212> PRT
 <213> Artificial Sequence
 <220>
 <223> Amino acid sequence derived from Murine monoclonal
       antibody and further modified by amino acid
       substitutions
 <400> 727
 Asp Thr Phe Arg Gln Pro Ser
  1
 <210> 728
 <211> 7
 <212> PRT
 <213> Artificial Sequence
 <220>
 <223> Amino acid sequence derived from Murine monoclonal
       antibody and further modified by amino acid
       substitutions
 <400> 728
 Asp Thr Phe Arg Gln Thr Ser
 <210> 729
 <211> 7
 <212> PRT
 <213> Artificial Sequence
```

```
<220>
<223> Amino acid sequence derived from Murine monoclonal
     antibody and further modified by amino acid
      substitutions
<400> 729
Asp Thr Phe Arg Gln Asp Ser
<210> 730
<211> 7
<212> PRT
<213> Artificial Sequence
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 730
Asp Thr Phe Tyr Leu Lys Ser
<210> 731
<211> 7
<212> PRT
<213> Artificial Sequence
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 731
Asp Thr Phe Tyr Leu Arg Ser
 <210> 732
 <211> 7
 <212> PRT
 <213> Artificial Sequence
 <223> Amino acid sequence derived from Murine monoclonal
       antibody and further modified by amino acid
       substitutions
 <400> 732
 Asp Thr Phe Tyr Leu His Ser
 <210> 733
 <211> 7
 <212> PRT
 <213> Artificial Sequence
 <223> Amino acid sequence derived from Murine monoclonal
       antibody and further modified by amino acid
       substitutions
```

```
<400> 733
Asp Thr Phe Tyr Leu Pro Ser
<210> 734
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 734
Asp Thr Phe Tyr Leu Thr Ser
<210> 735
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
       substitutions
 <400> 735
 Asp Thr Phe Tyr Leu Asp Ser
 1
 <210> 736
 <211> 7
 <212> PRT
 <213> Artificial Sequence
 <220>
 <223> Amino acid sequence derived from Murine monoclonal
       antibody and further modified by amino acid
       substitutions
 <400> 736
 Asp Thr Phe Tyr His Ala Ser
 <210> 737
 <211> 7
 <212> PRT
 <213> Artificial Sequence
  <223> Amino acid sequence derived from Murine monoclonal
       antibody and further modified by amino acid
        substitutions
  <400> 737
  Asp Thr Phe Tyr His Ser Ser
                  5
  <210> 738
```

```
<211> 7
<212> PRT
<213> Artificial Sequence
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 738
Asp Thr Phe Tyr His Lys Ser
<210> 739
<211> 7
<212> PRT
<213> Artificial Sequence
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 739
Asp Thr Phe Tyr His Arg Ser
<210> 740
<211> 7
 <212> PRT
 <213> Artificial Sequence
 <223> Amino acid sequence derived from Murine monoclonal
       antibody and further modified by amino acid
       substitutions
 <400> 740
 Asp Thr Phe Tyr His His Ser
 <210> 741
 <211> 7
 <212> PRT
 <213> Artificial Sequence
 <223> Amino acid sequence derived from Murine monoclonal
       antibody and further modified by amino acid
       substitutions
 <400> 741
 Asp Thr Phe Tyr His Pro Ser
  <210> 742
  <211> 7
  <212> PRT
  <213> Artificial Sequence
  <220>
```

```
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 742
Asp Thr Phe Tyr His Thr Ser
<210> 743
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 743
Asp Thr Phe Tyr His Asp Ser
                5
<210> 744
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 744
Asp Thr Phe Tyr Gln Ala Ser
1
                5
<210> 745
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 745
Asp Thr Phe Tyr Gln Ser Ser
 1
                 5
<210> 746
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 746
```

```
Asp Thr Phe Tyr Gln Lys Ser
1
<210> 747
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 747
Asp Thr Phe Tyr Gln Arg Ser
<210> 748
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 748
Asp Thr Phe Tyr Gln His Ser
1
<210> 749
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 749
Asp Thr Phe Tyr Gln Pro Ser
1 5
<210> 750
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 750
Asp Thr Phe Tyr Gln Thr Ser
<210> 751
<211> 7
```

```
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
     substitutions
<400> 751
Asp Thr Phe Tyr Gln Asp Ser
<210> 752
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 752
Asp Thr Phe Phe Leu Ala Ser
<210> 753
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 753
Asp Thr Phe Phe Leu Ser Ser
<210> 754
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 754
Asp Thr Phe Phe Leu Lys Ser
<210> 755
<211> 7
<212> PRT
<213> Artificial Sequence
<223> Amino acid sequence derived from Murine monoclonal
```

antibody and further modified by amino acid substitutions

```
<400> 755
Asp Thr Phe Phe Leu Arg Ser
<210> 756
<211> 7
<212> PRT
<213> Artificial Sequence
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 756
Asp Thr Phe Phe Leu His Ser
            5
<210> 757
<211> 7
<212> PRT
<213> Artificial Sequence
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
       substitutions
<400> 757
Asp Thr Phe Phe Leu Pro Ser
<210> 758
 <211> 7
 <212> PRT
 <213> Artificial Sequence
 <220>
 <223> Amino acid sequence derived from Murine monoclonal
       antibody and further modified by amino acid
       substitutions
 <400> 758
 Asp Thr Phe Phe Leu Thr Ser
 <210> 759
 <211> 7
 <212> PRT
 <213> Artificial Sequence
 <220>
 <223> Amino acid sequence derived from Murine monoclonal
       antibody and further modified by amino acid
       substitutions
 <400> 759
```

222

```
Asp Thr Phe Phe His Ala Ser
    5
<210> 760
<211> 7
<212> PRT
<213> Artificial Sequence
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 760
Asp Thr Phe Phe His Ser Ser
<210> 761
<211> 7
<212> PRT
<213> Artificial Sequence
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 761
Asp Thr Phe Phe His Lys Ser
<210> 762
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
 <223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
       substitutions
 <400> 762
 Asp Thr Phe Phe His Arg Ser
         5
 <210> 763
 <211> 7
 <212> PRT
 <213> Artificial Sequence
 <220>
 <223> Amino acid sequence derived from Murine monoclonal
       antibody and further modified by amino acid
       substitutions
 <400> 763
 Asp Thr Phe Phe His His Ser
                 5
 <210> 764
 <211> 7
```

```
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 764
Asp Thr Phe Phe His Pro Ser
<210> 765
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 765
Asp Thr Phe Phe His Thr Ser
                5
<210> 766
<211> 7
<212> PRT
<213> Artificial Sequence
<223> Amino acid sequence derived from Murine monoclonal
       antibody and further modified by amino acid
       substitutions
 <400> 766
Asp Thr Phe Phe His Asp Ser
 <210> 767
 <211> 7
 <212> PRT
 <213> Artificial Sequence
 <223> Amino acid sequence derived from Murine monoclonal
       antibody and further modified by amino acid
       substitutions
 <400> 767
 Asp Thr Phe Phe Gln Ala Ser
 <210> 768
 <211> 7
 <212> PRT
 <213> Artificial Sequence
 <220>
```

```
<223> Amino acid sequence derived from Murine monoclonal
     antibody and further modified by amino acid
      substitutions
<400> 768
Asp Thr Phe Phe Gln Ser Ser
<210> 769
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 769
Asp Thr Phe Phe Gln Lys Ser
<210> 770
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 770
Asp Thr Phe Phe Gln Arg Ser
<210> 771
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
 <223> Amino acid sequence derived from Murine monoclonal
       antibody and further modified by amino acid
       substitutions
 <400> 771
 Asp Thr Phe Phe Gln His Ser
 <210> 772
 <211> 7
 <212> PRT
 <213> Artificial Sequence
 <223> Amino acid sequence derived from Murine monoclonal
       antibody and further modified by amino acid
       substitutions
 <400> 772
```

```
Asp Thr Phe Phe Gln Pro Ser
<210> 773
<211> 7
<212> PRT
<213> Artificial Sequence
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 773
Asp Thr Phe Phe Gln Thr Ser
<210> 774
<211> 7
<212> PRT
<213> Artificial Sequence
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 774
Asp Thr Phe Phe Gln Asp Ser
<210> 775
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
 <223> Amino acid sequence derived from Murine monoclonal
       antibody and further modified by amino acid
       substitutions
 <400> 775
 Asp Thr Phe Leu Leu Ala Ser
 <210> 776
 <211> 7
 <212> PRT
 <213> Artificial Sequence
 <220>
 <223> Amino acid sequence derived from Murine monoclonal
       antibody and further modified by amino acid
       substitutions
 <400> 776
 Asp Thr Phe Leu Leu Ser Ser
 <21.0> 777
 <211> 7
```

```
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 777
Asp Thr Phe Leu Leu Lys Ser
<210> 778
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 778
Asp Thr Phe Leu Leu Arg Ser
<210> 779
<211> 7
 <212> PRT
 <213> Artificial Sequence
 <223> Amino acid sequence derived from Murine monoclonal
       antibody and further modified by amino acid
       substitutions
 <400> 779
 Asp Thr Phe Leu Leu His Ser
 <210> 780
 <211> 7
 <212> PRT
 <213> Artificial Sequence
 <223> Amino acid sequence derived from Murine monoclonal
       antibody and further modified by amino acid
       substitutions
 <400> 780
 Asp Thr Phe Leu Leu Pro Ser
 <210> 781
 <211> 7
 <212> PRT
 <213> Artificial Sequence
 <220>
 <223> Amino acid sequence derived from Murine monoclonal
```

antibody and further modified by amino acid substitutions

```
<400> 781
Asp Thr Phe Leu Leu Thr Ser
<210> 782
<211> 7
<212> PRT
<213> Artificial Sequence
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 782
Asp Thr Phe Leu Leu Asp Ser
<210> 783
<211> 7
<212> PRT
<213> Artificial Sequence
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 783
Asp Thr Phe Leu His Ala Ser
<210> 784
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 784
Asp Thr Phe Leu His Ser Ser
<210> 785
 <211> 7
 <212> PRT
 <213> Artificial Sequence
 <220>
 <223> Amino acid sequence derived from Murine monoclonal
       antibody and further modified by amino acid
       substitutions
 <400> 785
 Asp Thr Phe Leu His Lys Ser
```

228

```
5
<210> 786
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 786
Asp Thr Phe Leu His Arg Ser
<210> 787
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 787
Asp Thr Phe Leu His His Ser
            5
<210> 788
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 788
Asp Thr Phe Leu His Pro Ser
             5
<210> 789
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 789
Asp Thr Phe Leu His Thr Ser
<210> 790
<211> 7
<212> PRT
```

```
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 790
Asp Thr Phe Leu His Asp Ser
<210> 791
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 791
Asp Thr Phe Leu Gln Ala Ser
<210> 792
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 792
Asp Thr Phe Leu Gln Ser Ser
                5
<210> 793
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 793
Asp Thr Phe Leu Gln Lys Ser
<210> 794
<211> 7
<212> PRT
<213> Artificial Sequence
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
```

substitutions <400> 794 Asp Thr Phe Leu Gln Arg Ser <210> 795 <211> 7 <212> PRT <213> Artificial Sequence <220> <223> Amino acid sequence derived from Murine monoclonal antibody and further modified by amino acid substitutions <400> 795 Asp Thr Phe Leu Gln His Ser 5 <210> 796 <211> 7 <212> PRT <213> Artificial Sequence <220> <223> Amino acid sequence derived from Murine monoclonal antibody and further modified by amino acid substitutions <400> 796 Asp Thr Phe Leu Gln Pro Ser 5 <210> 797 <211> 7 <212> PRT <213> Artificial Sequence <220> <223> Amino acid sequence derived from Murine monoclonal antibody and further modified by amino acid substitutions <400> 797 Asp Thr Phe Leu Gln Thr Ser <210> 798 <211> 7 <212> PRT <213> Artificial Sequence <220> <223> Amino acid sequence derived from Murine monoclonal antibody and further modified by amino acid substitutions <400> 798 Asp Thr Phe Leu Gln Asp Ser

1

```
<210> 799
<211> 7
<212> PRT
<213> Artificial Sequence
<223> Amino acid sequence derived from Murine monoclonal
     antibody and further modified by amino acid
      substitutions
<400> 799
Asp Thr Tyr Lys Leu Ala Ser
<210> 800
<211> 7
<212> PRT
<213> Artificial Sequence
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 800
Asp Thr Tyr Lys Leu Ser Ser
<210> 801
<211> 7
<212> PRT
<213> Artificial Sequence
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 801
Asp Thr Tyr Lys Leu Lys Ser
<210> 802
<211> 7
 <212> PRT
 <213> Artificial Sequence
 <223> Amino acid sequence derived from Murine monoclonal
       antibody and further modified by amino acid
       substitutions
 <400> 802
 Asp Thr Tyr Lys Leu Arg Ser
 <210> 803
 <211> 7
 <212> PRT
 <213> Artificial Sequence
```

```
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 803
Asp Thr Tyr Lys Leu His Ser
               5
<210> 804
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 804
Asp Thr Tyr Lys Leu Pro Ser
<210> 805
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 805
Asp Thr Tyr Lys Leu Thr Ser
<210> 806
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
 <400> 806
Asp Thr Tyr Lys Leu Asp Ser
<210> 807
 <211> 7
 <212> PRT
 <213> Artificial Sequence
 <223> Amino acid sequence derived from Murine monoclonal
       antibody and further modified by amino acid
       substitutions
```

```
<400> 807
Asp Thr Tyr Lys His Ala Ser
<210> 808
<211> 7
<212> PRT
<213> Artificial Sequence
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 808
Asp Thr Tyr Lys His Ser Ser
<210> 809
<211> 7
<212> PRT
<213> Artificial Sequence
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 809
Asp Thr Tyr Lys His Lys Ser
<210> 810
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
       antibody and further modified by amino acid
       substitutions
 <400> 810
 Asp Thr Tyr Lys His Arg Ser
                 5
 <210> 811
 <211> 7
 <212> PRT
 <213> Artificial Sequence
 <220>
 <223> Amino acid sequence derived from Murine monoclonal
       antibody and further modified by amino acid
       substitutions
 <400> 811
 Asp Thr Tyr Lys His His Ser
                  5
```

```
<210> 812
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 812
Asp Thr Tyr Lys His Pro Ser
                5
<210> 813
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 813
Asp Thr Tyr Lys His Thr Ser
<210> 814
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 814
Asp Thr Tyr Lys His Asp Ser
<210> 815
 <2.1.1> 7
 <212> PRT
 <213> Artificial Sequence
 <220>
 <223> Amino acid sequence derived from Murine monoclonal
       antibody and further modified by amino acid
       substitutions
 <400> 815
 Asp Thr Tyr Lys Gln Ala Ser
 <210> 816
 <211> 7
 <212> PRT
 <213> Artificial Sequence
```

```
<220>
<223> Amino acid sequence derived from Murine monoclonal
     antibody and further modified by amino acid
     substitutions
<400> 816
Asp Thr Tyr Lys Gln Ser Ser
<210> 817
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 817
Asp Thr Tyr Lys Gln Lys Ser
<210> 818
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 818
Asp Thr Tyr Lys Gln Arg Ser
           5
<210> 819
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 819
Asp Thr Tyr Lys Gln His Ser
<210> 820
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
```

```
<400> 820
Asp Thr Tyr Lys Gln Pro Ser
<210> 821
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 821
Asp Thr Tyr Lys Gln Thr Ser
<210> 822
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 822
Asp Thr Tyr Lys Gln Asp Ser
<210> 823
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 823
Asp Thr Tyr Gly Leu Ala Ser
                5
<210> 824
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
 <223> Amino acid sequence derived from Murine monoclonal
       antibody and further modified by amino acid
       substitutions
 <400> 824
 Asp Thr Tyr Gly Leu Ser Ser
  1
```

```
<210> 825
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 825
Asp Thr Tyr Gly Leu Lys Ser
                 5
<210> 826
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 826
Asp Thr Tyr Gly Leu Arg Ser
1
                 5
<210> 827
<211> 7
<212> PRT
<213> Artificial Sequence
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 827
Asp Thr Tyr Gly Leu His Ser
                 5
<210> 828
<211> 7
<212> PRT
<213> Artificial Sequence
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 828
Asp Thr Tyr Gly Leu Pro Ser
<210> 829
<211> 7
<212> PRT
<213> Artificial Sequence
```

```
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 829
Asp Thr Tyr Gly Leu Thr Ser
<210> 830
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 830
Asp Thr Tyr Gly Leu Asp Ser
                 5
<210> 831
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 831
Asp Thr Tyr Gly His Ala Ser
                 5
<210> 832
<211> 7
<212> PRT
<213> Artificial Sequence
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 832
Asp Thr Tyr Gly His Ser Ser
                 5
<210> 833
<211> 7
<212> PRT
<213> Artificial Sequence
<223> Amino acid sequence derived from Murine monoclonal
     antibody and further modified by amino acid
```

substitutions <400> 833 Asp Thr Tyr Gly His Lys Ser <210> 834 <211> 7 <212> PRT <213> Artificial Sequence <220> <223> Amino acid sequence derived from Murine monoclonal antibody and further modified by amino acid substitutions <400> 834 Asp Thr Tyr Gly His Arg Ser <210> 835 <211> 7 <212> PRT <213> Artificial Sequence <223> Amino acid sequence derived from Murine monoclonal antibody and further modified by amino acid substitutions <400> 835 Asp Thr Tyr Gly His His Ser <210> 836 <211> 7 <212> PRT <213> Artificial Sequence <223> Amino acid sequence derived from Murine monoclonal antibody and further modified by amino acid substitutions <400> 836 Asp Thr Tyr Gly His Pro Ser <210> 837 <211> 7 <212> PRT <213> Artificial Sequence <220>

antibody and further modified by amino acid substitutions <400> 837 Asp Thr Tyr Gly His Thr Ser

1

<223> Amino acid sequence derived from Murine monoclonal

```
<210> 838
<211> 7
<212> PRT
<213> Artificial Sequence
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 838
Asp Thr Tyr Gly His Asp Ser
                 5
<210> 839
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 839
Asp Thr Tyr Gly Gln Ala Ser
                5
1
<210> 840
<211> 7
<212> PRT
<213> Artificial Sequence
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 840
Asp Thr Tyr Gly Gln Ser Ser
                5
1
<210> 841
<211> 7
<212> PRT
<213> Artificial Sequence
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 841
Asp Thr Tyr Gly Gln Lys Ser
                 5
<210> 842
<211> 7
<212> PRT
<213> Artificial Sequence
```

```
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 842
Asp Thr Tyr Gly Gln Arg Ser
<210> 843
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 843
Asp Thr Tyr Gly Gln His Ser
<210> 844
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 844
Asp Thr Tyr Gly Gln Pro Ser
                 5
<210> 845
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 845
Asp Thr Tyr Gly Gln Thr Ser
                 5
<210> 846
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
```

```
<400> 846
Asp Thr Tyr Gly Gln Asp Ser
1
<210> 847
<211> 7
<212> PRT
<213> Artificial Sequence
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 847
Asp Thr Tyr Arg Leu Ala Ser
1
                 5
<210> 848
<211> 7
<212> PRT
<213> Artificial Sequence
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 848
Asp Thr Tyr Arg Leu Ser Ser
                 5
<210> 849
<211> 7
<212> PRT
<213> Artificial Sequence
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 849
Asp Thr Tyr Arg Leu Lys Ser
<210> 850
<211> 7
<212> PRT
<213> Artificial Sequence
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 850
Asp Thr Tyr Arg Leu Arg Ser
                5
```

```
<210> 851
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 851
Asp Thr Tyr Arg Leu His Ser
<210> 852
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 852
Asp Thr Tyr Arg Leu Pro Ser
<210> 853
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
       substitutions
 <400> 853
 Asp Thr Tyr Arg Leu Thr Ser
 <210> 854
 <211> 7
 <212> PRT
 <213> Artificial Sequence
 <220>
 <223> Amino acid sequence derived from Murine monoclonal
       antibody and further modified by amino acid
       substitutions
 <400> 854
 Asp Thr Tyr Arg Leu Asp Ser
 <210> 855
 <211> 7
 <212> PRT
 <213> Artificial Sequence
```

```
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 855
Asp Thr Tyr Arg His Ala Ser
<210> 856
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 856
Asp Thr Tyr Arg His Ser Ser
        5
<210> 857
<211> 7
<212> PRT
<213> Artificial Sequence
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 857
Asp Thr Tyr Arg His Lys Ser
<210> 858
<211> 7
<212> PRT
<213> Artificial Sequence
 <223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
       substitutions
 <400> 858
 Asp Thr Tyr Arg His Arg Ser
                5
 <210> 859
 <211> 7
 <212> PRT
 <213> Artificial Sequence
<223> Amino acid sequence derived from Murine monoclonal
       antibody and further modified by amino acid
       substitutions
```

```
<400> 859
Asp Thr Tyr Arg His His Ser
<210> 860
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 860
Asp Thr Tyr Arg His Pro Ser
                 5
<210> 861
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 861
Asp Thr Tyr Arg His Thr Ser
<210> 862
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
 <223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
       substitutions
 <400> 862
 Asp Thr Tyr Arg His Asp Ser
 <210> 863
 <211> 7
 <212> PRT
 <213> Artificial Sequence
 <220>
 <223> Amino acid sequence derived from Murine monoclonal
       antibody and further modified by amino acid
       substitutions
 <400> 863
 Asp Thr Tyr Arg Gln Ala Ser
  1
 <210> 864
```

```
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 864
Asp Thr Tyr Arg Gln Ser Ser
<210> 865
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 865
Asp Thr Tyr Arg Gln Lys Ser
 1 5
<210> 866
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 866
Asp Thr Tyr Arg Gln Arg Ser
<210> 867
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 867
Asp Thr Tyr Arg Gln His Ser
<210> 868
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
```

```
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
     substitutions
<400> 868
Asp Thr Tyr Arg Gln Pro Ser
<210> 869
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 869
Asp Thr Tyr Arg Gln Thr Ser
<210> 870
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 870
Asp Thr Tyr Arg Gln Asp Ser
<210> 871
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 871
Asp Thr Tyr Tyr Leu Ala Ser
<210> 872
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
       antibody and further modified by amino acid
       substitutions
<400> 872
```

```
Asp Thr Tyr Tyr Leu Ser Ser
1 5
<210> 873
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 873
Asp Thr Tyr Tyr Leu Lys Ser
     5
<210> 874
<211> 7
<212> PRT
<213> Artificial Sequence
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 874
Asp Thr Tyr Tyr Leu Arg Ser
<210> 875
<211> 7
<212> PRT
<213> Artificial Sequence
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
 <400> 875
 Asp Thr Tyr Tyr Leu His Ser
 <210> 876
 <211> 7
 <212> PRT
 <213> Artificial Sequence
 <220>
 <223> Amino acid sequence derived from Murine monoclonal
       antibody and further modified by amino acid
       substitutions
 <400> 876
 Asp Thr Tyr Tyr Leu Pro Ser
 <210> 877
 <211> 7
```

```
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 877
Asp Thr Tyr Tyr Leu Thr Ser
<210> 878
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 878
Asp Thr Tyr Tyr Leu Asp Ser
<210> 879
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 879
Asp Thr Tyr Tyr His Ala Ser
                5
<210> 880
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
       antibody and further modified by amino acid
       substitutions
<400> 880
Asp Thr Tyr Tyr His Ser Ser
<210> 881
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
```

antibody and further modified by amino acid substitutions

```
<400> 881
Asp Thr Tyr Tyr His Lys Ser
<210> 882
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 882
Asp Thr Tyr Tyr His Arg Ser
     5
<210> 883
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 883
Asp Thr Tyr Tyr His His Ser
           5
<210> 884
<211> 7
<212> PRT
<213> Artificial Sequence
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 884
Asp Thr Tyr Tyr His Pro Ser
<210> 885
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
 <400> 885
```

```
Asp Thr Tyr Tyr His Thr Ser
<210> 886
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 886
Asp Thr Tyr Tyr His Asp Ser
<210> 887
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 887
Asp Thr Tyr Tyr Gln Ala Ser
<210> 888
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 888
Asp Thr Tyr Tyr Gln Ser Ser
            5
<210> 889
<211> 7
<212> PRT
<213> Artificial Sequence .
 <223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
       substitutions
 <400> 889
 Asp Thr Tyr Tyr Gln Lys Ser
 <210> 890
 <211> 7
```

```
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 890
Asp Thr Tyr Tyr Gln Arg Ser
                5
<210> 891
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 891
Asp Thr Tyr Tyr Gln His Ser
                 5
<210> 892
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 892
Asp Thr Tyr Tyr Gln Pro Ser
                 5
<210> 893
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 893
Asp Thr Tyr Tyr Gln Thr Ser
 1
<210> 894
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
```

antibody and further modified by amino acid substitutions

<400> 894 Asp Thr Tyr Tyr Gln Asp Ser <210> 895 <211> 7 <212> PRT <213> Artificial Sequence <220> <223> Amino acid sequence derived from Murine monoclonal antibody and further modified by amino acid substitutions <400> 895 Asp Thr Tyr Phe Leu Ala Ser <210> 896 <211> 7 <212> PRT <213> Artificial Sequence <220> <223> Amino acid sequence derived from Murine monoclonal antibody and further modified by amino acid substitutions <400> 896 Asp Thr Tyr Phe Leu Ser Ser <210> 897 <211> 7 <212> PRT <213> Artificial Sequence <220> <223> Amino acid sequence derived from Murine monoclonal antibody and further modified by amino acid substitutions <400> 897 Asp Thr Tyr Phe Leu Lys Ser <210> 898 <211> 7 <212> PRT <213> Artificial Sequence <220> <223> Amino acid sequence derived from Murine monoclonal antibody and further modified by amino acid substitutions

<400> 898

```
Asp Thr Tyr Phe Leu Arg Ser
1
<210> 899
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 899
Asp Thr Tyr Phe Leu His Ser
<210> 900
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 900
Asp Thr Tyr Phe Leu Pro Ser
<210> 901
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 901
Asp Thr Tyr Phe Leu Thr Ser
<210> 902
 <211> 7
 <212> PRT
 <213> Artificial Sequence
 <223> Amino acid sequence derived from Murine monoclonal
       antibody and further modified by amino acid
       substitutions
 <400> 902
 Asp Thr Tyr Phe Leu Asp Ser
 <210> 903
 <211> 7
```

```
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 903
Asp Thr Tyr Phe His Ala Ser
<210> 904
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 904
Asp Thr Tyr Phe His Ser Ser
<210> 905
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 905
Asp Thr Tyr Phe His Lys Ser
<210> 906
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
       antibody and further modified by amino acid
       substitutions
<400> 906
Asp Thr Tyr Phe His Arg Ser
<210> 907
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
```

antibody and further modified by amino acid substitutions

```
<400> 907
Asp Thr Tyr Phe His His Ser
                 5
1
<210> 908
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 908
Asp Thr Tyr Phe His Pro Ser
                 5
<210> 909
<211> 7
<212> PRT
<213> Artificial Sequence
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 909
Asp Thr Tyr Phe His Thr Ser
 1
                 5
<210> 910
<211> 7
<212> PRT
<213> Artificial Sequence
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 910
Asp Thr Tyr Phe His Asp Ser
 1
                 5
<210> 911
<211> 7
<212> PRT
<213> Artificial Sequence
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 911
Asp Thr Tyr Phe Gln Ala Ser
```

257

```
1
                 5
<210> 912
<211> 7
<212> PRT
<213> Artificial Sequence
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 912
Asp Thr Tyr Phe Gln Ser Ser
<210> 913
<211> 7
<212> PRT
<213> Artificial Sequence
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 913
Asp Thr Tyr Phe Gln Lys Ser
<210> 914
<211> 7
<212> PRT
<213> Artificial Sequence
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 914
Asp Thr Tyr Phe Gln Arg Ser
<210> 915
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 915
Asp Thr Tyr Phe Gln His Ser
<210> 916
<211> 7
<212> PRT
```

```
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 916
Asp Thr Tyr Phe Gln Pro Ser
<210> 917
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 917
Asp Thr Tyr Phe Gln Thr Ser
           5
<210> 918
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 918
Asp Thr Tyr Phe Gln Asp Ser
 1
            5
<210> 919
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 919
Asp Thr Tyr Leu Leu Ala Ser
            5
<210> 920
<211> 7
<212> PRT
<213> Artificial Sequence
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
```

substitutions <400> 920 Asp Thr Tyr Leu Leu Ser Ser <210> 921 <211> 7 <212> PRT <213> Artificial Sequence <220> <223> Amino acid sequence derived from Murine monoclonal antibody and further modified by amino acid substitutions <400> 921 Asp Thr Tyr Leu Leu Lys Ser <210> 922 <211> 7 <212> PRT <213> Artificial Sequence <220> <223> Amino acid sequence derived from Murine monoclonal antibody and further modified by amino acid substitutions <400> 922 Asp Thr Tyr Leu Leu Arg Ser <210> 923 <211> 7 <212> PRT <213> Artificial Sequence <220> <223> Amino acid sequence derived from Murine monoclonal antibody and further modified by amino acid substitutions <400> 923 Asp Thr Tyr Leu Leu His Ser <210> 924 <211> 7 <212> PRT <213> Artificial Sequence <220>

<223> Amino acid sequence derived from Murine monoclonal antibody and further modified by amino acid substitutions

<400> 924 Asp Thr Tyr Leu Leu Pro Ser 1 5

```
<210> 925
<211> 7
<212> PRT
<213> Artificial Sequence
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 925
Asp Thr Tyr Leu Leu Thr Ser
<210> 926
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 926
Asp Thr Tyr Leu Leu Asp Ser
<210> 927
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 927
Asp Thr Tyr Leu His Ala Ser
<210> 928
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 928
Asp Thr Tyr Leu His Ser Ser
     5
<210> 929
<211> 7
<212> PRT
<213> Artificial Sequence
```

```
<220>
<223> Amino acid sequence derived from Murine monoclonal
     antibody and further modified by amino acid
     substitutions
<400> 929
Asp Thr Tyr Leu His Lys Ser
                 5
<210> 930
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 930
Asp Thr Tyr Leu His Arg Ser
                 5
<210> 931
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 931
Asp Thr Tyr Leu His His Ser
                 5
<210> 932
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 932
Asp Thr Tyr Leu His Pro Ser
<210> 933
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
```

```
<400> 933
Asp Thr Tyr Leu His Thr Ser
<210> 934
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 934
Asp Thr Tyr Leu His Asp Ser
<210> 935
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 935
Asp Thr Tyr Leu Gln Ala Ser
<210> 936
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 936
Asp Thr Tyr Leu Gln Ser Ser
            5
<210> 937
<211> 7
<212> PRT
<213> Artificial Sequence
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 937
Asp Thr Tyr Leu Gln Lys Ser
```

```
<210> 938
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 938
Asp Thr Tyr Leu Gln Arg Ser
<210> 939
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 939
Asp Thr Tyr Leu Gln His Ser
<210> 940
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 940
Asp Thr Tyr Leu Gln Pro Ser
<210> 941
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 941
Asp Thr Tyr Leu Gln Thr Ser
<210> 942
<211> 7
<212> PRT
<213> Artificial Sequence
```

```
<220>
<223> Amino acid sequence derived from Murine monoclonal
     antibody and further modified by amino acid
     substitutions
<400> 942
Asp Thr Tyr Leu Gln Asp Ser
1
                 5
<210> 943
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
     substitutions
<400> 943
Asp Thr Arg Lys Leu Lys Ser
                5
<210> 944
<211> 7
<212> PRT
<213> Artificial Sequence
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 944
Asp Thr Arg Lys Leu Arg Ser
1
                 5
<210> 945
<211> 7
<212> PRT
<213> Artificial Sequence
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 945
Asp Thr Arg Lys Leu His Ser
                 5
<210> 946
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
```

```
<400> 946
Asp Thr Arg Lys Leu Thr Ser
1
<210> 947
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 947
Asp Thr Arg Lys Leu Asp Ser
<210> 948
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 948
Asp Thr Arg Lys His Ala Ser
1
                5
<210> 949
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 949
Asp Thr Arg Lys His Ser Ser
 1
                 5
<210> 950
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 950
Asp Thr Arg Lys His Lys Ser
<210> 951
```

```
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 951
Asp Thr Arg Lys His Arg Ser
1
                5
<210> 952
<211> 7
<212> PRT
<213> Artificial Sequence
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 952
Asp Thr Arg Lys His His Ser
1
                5
<210> 953
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 953
Asp Thr Arg Lys His Pro Ser
1
                5
<210> 954
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 954
Asp Thr Arg Lys His Thr Ser
<210> 955
<211> 7
<212> PRT
<213> Artificial Sequence
```

```
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 955
Asp Thr Arg Lys His Asp Ser
<210> 956
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 956
Asp Thr Arg Lys Gln Lys Ser
<210> 957
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 957
Asp Thr Arg Lys Gln Arg Ser
<210> 958
<211> 7
<212> PRT
<213> Artificial Sequence
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 958
Asp Thr Arg Lys Gln His Ser
<210> 959
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
```

```
<400> 959
 Asp Thr Arg Lys Gln Pro Ser
                  5
 1
 <210> 960
 <211> 7
 <212> PRT
<213> Artificial Sequence
 <220>
 <223> Amino acid sequence derived from Murine monoclonal
       antibody and further modified by amino acid
       substitutions
 <400> 960
 Asp Thr Arg Lys Gln Thr Ser
  1 5
 <210> 961
 <211> 7
 <212> PRT
 <213> Artificial Sequence
 <223> Amino acid sequence derived from Murine monoclonal
       antibody and further modified by amino acid
       substitutions
 <400> 961
 Asp Thr Arg Lys Gln Asp Ser
                  5
  1
 <210> 962
 <211> 7
 <212> PRT
 <213> Artificial Sequence
 <223> Amino acid sequence derived from Murine monoclonal
       antibody and further modified by amino acid
       substitutions
 <400> 962
 Asp Thr Arg Gly Leu Ser Ser
  1
                  5
 <210> 963
 <211> 7
 <212> PRT
 <213> Artificial Sequence
 <223> Amino acid sequence derived from Murine monoclonal
       antibody and further modified by amino acid
       substitutions
 <400> 963
 Asp Thr Arg Gly Leu Lys Ser
  1
```

```
<210> 964
<211> 7
<212> PRT
<213> Artificial Sequence
<223> Amino acid sequence derived from Murine monoclonal
     antibody and further modified by amino acid
      substitutions
<400> 964
Asp Thr Arg Gly Leu Arg Ser
<210> 965
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 965
Asp Thr Arg Gly Leu His Ser
<210> 966
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 966
Asp Thr Arg Gly Leu Thr Ser
<210> 967
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 967
Asp Thr Arg Gly Leu Asp Ser
<210> 968
<211> 7
<212> PRT
<213> Artificial Sequence
```

```
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 968
Asp Thr Arg Gly His Ala Ser
                5
<210> 969
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 969
Asp Thr Arg Gly His Ser Ser
<210> 970
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 970
Asp Thr Arg Gly His Lys Ser
<210> 971
<211> 7
<212> PRT
<213> Artificial Sequence
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 971
Asp Thr Arg Gly His Arg Ser
                 5
<210> 972
<211> 7
<212> PRT
<213> Artificial Sequence
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
```

```
<400> 972
Asp Thr Arg Gly His His Ser
<210> 973
<211> 7
<212> PRT
<213> Artificial Sequence
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 973
Asp Thr Arg Gly His Pro Ser
<210> 974
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal.
      antibody and further modified by amino acid
      substitutions
<400> 974
Asp Thr Arg Gly His Thr Ser
<210> 975
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 975
Asp Thr Arg Gly His Asp Ser
<210> 976
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 976
Asp Thr Arg Gly Gln Ala Ser
<210> 977
```

272

```
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 977
Asp Thr Arg Gly Gln Ser Ser
<210> 978
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 978
Asp Thr Arg Gly Gln Lys Ser
<210> 979
<211> 7
<212> PRT
<213> Artificial Sequence
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 979
Asp Thr Arg Gly Gln Arg Ser
1
                 5
<210> 980
<211> 7
<212> PRT
<213> Artificial Sequence
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 980
Asp Thr Arg Gly Gln His Ser
1
                5
<210> 981
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
```

273

```
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 981
Asp Thr Arg Gly Gln Pro Ser
<210> 982
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 982
Asp Thr Arg Gly Gln Thr Ser
1
                5
<210> 983
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 983
Asp Thr Arg Gly Gln Asp Ser
 1
                5
<210> 984
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 984
Asp Thr Arg Arg Leu Ala Ser
 1
<210> 985
<211> 7
<212> PRT
<213> Artificial Sequence
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 985
```

```
Asp Thr Arg Arg Leu Ser Ser
<210> 986
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 986
Asp Thr Arg Arg Leu Lys Ser
<210> 987
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 987
Asp Thr Arg Arg Leu Arg Ser
           5
<210> 988
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 988
Asp Thr Arg Arg Leu His Ser
<210> 989
<211> 7
<212> PRT
<213> Artificial Sequence
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 989
Asp Thr Arg Arg Leu Pro Ser
<210> 990
<211> 7
```

```
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
     antibody and further modified by amino acid
     substitutions
<400> 990
Asp Thr Arg Arg Leu Thr Ser
<210> 991
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 991
Asp Thr Arg Arg Leu Asp Ser
<210> 992
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 992
Asp Thr Arg Arg His Ala Ser
                 5
<210> 993
<211> 7
<212> PRT
<213> Artificial Sequence
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 993
Asp Thr Arg Arg His Ser Ser
<210> 994
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
```

antibody and further modified by amino acid substitutions <400> 994 Asp Thr Arg Arg His Lys Ser 1 <210> 995 <211> 7 <212> PRT <213> Artificial Sequence <223> Amino acid sequence derived from Murine monoclonal antibody and further modified by amino acid substitutions <400> 995 Asp Thr Arg Arg His Arg Ser 1 5 <210> 996 <211> 7 <212> PRT <213> Artificial Sequence <223> Amino acid sequence derived from Murine monoclonal antibody and further modified by amino acid substitutions <400> 996 Asp Thr Arg Arg His His Ser 5 <210> 997 <211> 7 <212> PRT <213> Artificial Sequence <220> <223> Amino acid sequence derived from Murine monoclonal antibody and further modified by amino acid substitutions <400> 997 Asp Thr Arg Arg His Pro Ser 5 1 <210> 998 <211> 7 <212> PRT <213> Artificial Sequence <220> <223> Amino acid sequence derived from Murine monoclonal antibody and further modified by amino acid

<400> 998 Asp Thr Arg Arg His Thr Ser

substitutions

```
5
1
<210> 999
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 999
Asp Thr Arg Arg His Asp Ser
<210> 1000
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 1000
Asp Thr Arg Arg Gln Ala Ser
<210> 1001
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 1001
Asp Thr Arg Arg Gln Ser Ser
<210> 1002
<211> 7
 <212> PRT
<213> Artificial Sequence
 <223> Amino acid sequence derived from Murine monoclonal
       antibody and further modified by amino acid
       substitutions
 <400> 1002
 Asp Thr Arg Arg Gln Lys Ser
 <210> 1003
 <211> 7
 <212> PRT
```

```
<213> Artificial Sequence
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 1003
Asp Thr Arg Arg Gln Arg Ser
1
                 5
<210> 1004
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 1004
Asp Thr Arg Arg Gln His Ser
1
           5
<210> 1005
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 1005
Asp Thr Arg Arg Gln Pro Ser
 1
             5
<210> 1006
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 1006
Asp Thr Arg Arg Gln Thr Ser
                5
<210> 1007
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
```

antibody and further modified by amino acid substitutions

```
<400> 1007
Asp Thr Arg Arg Gln Asp Ser
<210> 1008
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 1008
Asp Thr Arg Tyr Leu Lys Ser
                5
<210> 1009
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 1009
Asp Thr Arg Tyr Leu Arg Ser
                5
<210> 1010
<211> 7
<212> PRT
<213> Artificial Sequence
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 1010
Asp Thr Arg Tyr Leu His Ser
                 5
<210> 1011
<211> 7
<212> PRT
<213> Artificial Sequence
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 1011
Asp Thr Arg Tyr Leu Pro Ser
```

```
1
<210> 1012
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 1012
Asp Thr Arg Tyr Leu Thr Ser
                5
<210> 1013
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 1013
Asp Thr Arg Tyr Leu Asp Ser
<210> 1014
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 1014
Asp Thr Arg Tyr His Ala Ser
<210> 1015
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 1015
Asp Thr Arg Tyr His Ser Ser
<210> 1016
<211> 7
<212> PRT
```

```
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 1016
Asp Thr Arg Tyr His Lys Ser
                5
<210> 1017
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 1017
Asp Thr Arg Tyr His Arg Ser
                5
<210> 1018
<211> 7
<212> PRT
<213> Artificial Sequence
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 1018
Asp Thr Arg Tyr His His Ser
                5
<210> 1019
<211> 7
<212> PRT
<213> Artificial Sequence
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 1019
Asp Thr Arg Tyr His Pro Ser
                5
<210> 1020
<211> 7
<212> PRT
<213> Artificial Sequence
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
```

substitutions <400> 1020 Asp Thr Arg Tyr His Thr Ser 5 <210> 1021 <211> 7 <212> PRT <213> Artificial Sequence <220> <223> Amino acid sequence derived from Murine monoclonal antibody and further modified by amino acid substitutions <400> 1021 Asp Thr Arg Tyr His Asp Ser <210> 1022 <211> 7 <212> PRT <213> Artificial Sequence <220> <223> Amino acid sequence derived from Murine monoclonal antibody and further modified by amino acid substitutions <400> 1022 Asp Thr Arg Tyr Gln Lys Ser <210> 1023 <211> 7 <212> PRT <213> Artificial Sequence <220> <223> Amino acid sequence derived from Murine monoclonal antibody and further modified by amino acid substitutions <400> 1023 Asp Thr Arg Tyr Gln Arg Ser <210> 1024 <211> 7 <212> PRT <213> Artificial Sequence <220> <223> Amino acid sequence derived from Murine monoclonal antibody and further modified by amino acid substitutions <400> 1024 Asp Thr Arg Tyr Gln His Ser

5

```
<210> 1025
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 1025
Asp Thr Arg Tyr Gln Pro Ser
<210> 1026
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 1026
Asp Thr Arg Tyr Gln Thr Ser
<210> 1027
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 1027
Asp Thr Arg Tyr Gln Asp Ser
<210> 1028
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 1028
Asp Thr Arg Phe Leu Ala Ser
<210> 1029
<211> 7
<212> PRT
```

```
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 1029
Asp Thr Arg Phe Leu Ser Ser
1
<210> 1030
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 1030
Asp Thr Arg Phe Leu Lys Ser
1
                 5
<210> 1031
<211> 7
<212> PRT
<213> Artificial Sequence
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 1031
Asp Thr Arg Phe Leu Arg Ser
1
                 5
<210> 1032
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 1032
Asp Thr Arg Phe Leu His Ser
                 5
<210> 1033
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
```

antibody and further modified by amino acid substitutions

```
<400> 1033
Asp Thr Arg Phe Leu Pro Ser
<210> 1034
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 1034
Asp Thr Arg Phe Leu Thr Ser
<210> 1035
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 1035
Asp Thr Arg Phe Leu Asp Ser
<210> 1036
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
       antibody and further modified by amino acid
       substitutions
<400> 1036
Asp Thr Arg Phe His Ala Ser
<210> 1037
<211> 7
 <212> PRT
 <213> Artificial Sequence
 <220>
 <223> Amino acid sequence derived from Murine monoclonal
       antibody and further modified by amino acid
       substitutions
 <400> 1037
```

286

```
Asp Thr Arg Phe His Ser Ser
<210> 1038
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 1038
Asp Thr Arg Phe His Lys Ser
           5
<210> 1039
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 1039
Asp Thr Arg Phe His Arg Ser
<210> 1040
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 1040
Asp Thr Arg Phe His His Ser
<210> 1041
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 1041
Asp Thr Arg Phe His Pro Ser
<210> 1042
<211> 7
```

```
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 1042
Asp Thr Arg Phe His Thr Ser
                 5
<210> 1043
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 1043
Asp Thr Arg Phe His Asp Ser
<210> 1044
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 1044
Asp Thr Arg Phe Gln Ala Ser
<210> 1045
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 1045
Asp Thr Arg Phe Gln Ser Ser
                 5
<210> 1046
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
```

antibody and further modified by amino acid substitutions <400> 1046 Asp Thr Arg Phe Gln Lys Ser <210> 1047 <211> 7 <212> PRT <213> Artificial Sequence <220> <223> Amino acid sequence derived from Murine monoclonal antibody and further modified by amino acid substitutions <400> 1047 Asp Thr Arg Phe Gln Arg Ser <210> 1048 <211> 7 <212> PRT <213> Artificial Sequence <220> <223> Amino acid sequence derived from Murine monoclonal antibody and further modified by amino acid substitutions <400> 1048 Asp Thr Arg Phe Gln His Ser 5 <210> 1049 <211> 7 <212> PRT <213> Artificial Sequence <223> Amino acid sequence derived from Murine monoclonal antibody and further modified by amino acid substitutions <400> 1049 Asp Thr Arg Phe Gln Pro Ser 1 <210> 1050 <211> 7 <212> PRT <213> Artificial Sequence <223> Amino acid sequence derived from Murine monoclonal antibody and further modified by amino acid substitutions <400> 1050

```
Asp Thr Arg Phe Gln Thr Ser
<210> 1051
<211> 7
<212> PRT
<213> Artificial Sequence
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 1051
Asp Thr Arg Phe Gln Asp Ser
                 5
<210> 1052
<211> 7
<212> PRT
<213> Artificial Sequence
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 1052
Asp Thr Arg Leu Leu Ala Ser
<210> 1053
<211> 7
<212> PRT
<213> Artificial Sequence
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 1053
Asp Thr Arg Leu Leu Ser Ser
1
                5
<210> 1054
<211> 7
<212> PRT
<213> Artificial Sequence
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 1054
Asp Thr Arg Leu Leu Lys Ser
                 5
<210> 1055
<211> 7
```

```
<212> PRT
<213> Artificial Sequence
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 1055
Asp Thr Arg Leu Leu Arg Ser
<210> 1056
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 1056
Asp Thr Arg Leu Leu His Ser
<210> 1057
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 1057
Asp Thr Arg Leu Leu Pro Ser
<210> 1058
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 1058
Asp Thr Arg Leu Leu Thr Ser
                - 5
<210> 1059
<211> 7
<212> PRT
<213> Artificial Sequence
<223> Amino acid sequence derived from Murine monoclonal
```

substitutions <400> 1059 Asp Thr Arg Leu Leu Asp Ser <210> 1060 <211> 7 <212> PRT <213> Artificial Sequence <220> <223> Amino acid sequence derived from Murine monoclonal antibody and further modified by amino acid substitutions <400> 1060 Asp Thr Arg Leu His Ala Ser <210> 1061 <211> 7 <212> PRT <213> Artificial Sequence <220> <223> Amino acid sequence derived from Murine monoclonal antibody and further modified by amino acid substitutions <400> 1061 Asp Thr Arg Leu His Ser Ser 5 <210> 1062 <211> 7 <212> PRT <213> Artificial Sequence <223> Amino acid sequence derived from Murine monoclonal antibody and further modified by amino acid substitutions <400> 1062 Asp Thr Arg Leu His Lys Ser <210> 1063 <211> 7 <212> PRT <213> Artificial Sequence <223> Amino acid sequence derived from Murine monoclonal antibody and further modified by amino acid substitutions <400> 1063 Asp Thr Arg Leu His Arg Ser

antibody and further modified by amino acid

```
1
<210> 1064
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 1064
Asp Thr Arg Leu His His Ser
<210> 1065
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 1065
Asp Thr Arg Leu His Pro Ser
<210> 1066
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 1066
Asp Thr Arg Leu His Thr Ser
               5
<210> 1067
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 1067
Asp Thr Arg Leu His Asp Ser
<210> 1068
<211> 7
<212> PRT
```

```
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 1068
Asp Thr Arg Leu Gln Ala Ser
                 5
<210> 1069
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 1069
Asp Thr Arg Leu Gln Ser Ser
<210> 1070
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 1070
Asp Thr Arg Leu Gln Lys Ser
<210> 1071
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 1071
Asp Thr Arg Leu Gln Arg Ser
<210> 1072
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
```

antibody and further modified by amino acid substitutions <400> 1072 Asp Thr Arg Leu Gln His Ser 1 <210> 1073 <211> 7 <212> PRT <213> Artificial Sequence <223> Amino acid sequence derived from Murine monoclonal antibody and further modified by amino acid substitutions <400> 1073 Asp Thr Arg Leu Gln Pro Ser <210> 1074 <211> 7 <212> PRT <213> Artificial Sequence <223> Amino acid sequence derived from Murine monoclonal antibody and further modified by amino acid substitutions <400> 1074 Asp Thr Arg Leu Gln Thr Ser <210> 1075 <211> 7 <212> PRT <213> Artificial Sequence <223> Amino acid sequence derived from Murine monoclonal antibody and further modified by amino acid substitutions <400> 1075 Asp Thr Arg Leu Gln Asp Ser <210> 1076 <211> 7 <212> PRT <213> Artificial Sequence <223> Amino acid sequence derived from Murine monoclonal antibody and further modified by amino acid substitutions <400> 1076

Asp Thr Met Lys Leu Lys Ser

```
1
                 5
<210> 1077
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 1077
Asp Thr Met Lys Leu Arg Ser
<210> 1078
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 1078
Asp Thr Met Lys Leu His Ser
<210> 1079
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 1079
Asp Thr Met Lys Leu Pro Ser
 1
<210> 1080
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 1080
Asp Thr Met Lys Leu Thr Ser
<210> 1081
<211> 7
<212> PRT
```

```
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 1081
Asp Thr Met Lys Leu Asp Ser
1
                5
<210> 1082
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 1082
Asp Thr Met Lys His Ala Ser
1
                5
<210> 1083
<211> 7
<212> PRT
<213> Artificial Sequence
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 1083
Asp Thr Met Lys His Ser Ser
 1
                5
<210> 1084
<211> 7
<212> PRT
<213> Artificial Sequence
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 1084
Asp Thr Met Lys His Lys Ser
                5
<210> 1085
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
```

PCT/US2006/024717 WO 2007/002543

substitutions <400> 1085 Asp Thr Met Lys His Arg Ser <210> 1086 <211> 7 <212> PRT <213> Artificial Sequence <220> <223> Amino acid sequence derived from Murine monoclonal antibody and further modified by amino acid substitutions <400> 1086 Asp Thr Met Lys His His Ser <210> 1087 <211> 7 <212> PRT <213> Artificial Sequence <220> <223> Amino acid sequence derived from Murine monoclonal antibody and further modified by amino acid substitutions <400> 1087 Asp Thr Met Lys His Pro Ser <210> 1088 <211> 7 <212> PRT <213> Artificial Sequence <220> <223> Amino acid sequence derived from Murine monoclonal antibody and further modified by amino acid substitutions <400> 1088 Asp Thr Met Lys His Thr Ser <210> 1089 <211> 7 <212> PRT <213> Artificial Sequence <220> <223> Amino acid sequence derived from Murine monoclonal antibody and further modified by amino acid substitutions <400> 1089

Asp Thr Met Lys His Asp Ser

```
1
                 5
<210> 1090
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 1090
Asp Thr Met Lys Gln Lys Ser
<210> 1091
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 1091
Asp Thr Met Lys Gln Arg Ser
<210> 1092
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 1092
Asp Thr Met Lys Gln His Ser
<210> 1093
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 1093
Asp Thr Met Lys Gln Pro Ser
 1
<210> 1094
<211> 7
<212> PRT
```

```
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 1094
Asp Thr Met Lys Gln Thr Ser
 1
<210> 1095
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 1095
Asp Thr Met Lys Gln Asp Ser
1
<210> 1096
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 1096
Asp Thr Met Gly Leu Ala Ser
<210> 1097
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 1097
Asp Thr Met Gly Leu Ser Ser
1
                 5
<210> 1098
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
```

```
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 1098
Asp Thr Met Gly Leu Lys Ser
<210> 1099
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 1099
Asp Thr Met Gly Leu Arg Ser
<210> 1100
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 1100
Asp Thr Met Gly Leu His Ser
<210> 1101
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 1101
Asp Thr Met Gly Leu Pro Ser
                 5
<210> 1102
<211> 7
<212> PRT
<213> Artificial Sequence
<223> Amino acid sequence derived from Murine monoclonal
       antibody and further modified by amino acid
       substitutions
<400> 1102
```

```
Asp Thr Met Gly Leu Thr Ser
1
                 5
<210> 1103
<211> 7
<212> PRT
<213> Artificial Sequence
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 1103
Asp Thr Met Gly Leu Asp Ser
                 5
<210> 1104
<211> 7
<212> PRT
<213> Artificial Sequence
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 1104
Asp Thr Met Gly His Ala Ser
                5
<210> 1105
<211> 7
<212> PRT
<213> Artificial Sequence
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 1105
Asp Thr Met Gly His Ser Ser
               5
<210> 1106
<211> 7
<212> PRT
<213> Artificial Sequence
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 1106
Asp Thr Met Gly His Lys Ser
               5
<210> 1107
<211> 7
```

```
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
     antibody and further modified by amino acid
     substitutions
<400> 1107
Asp Thr Met Gly His Arg Ser
<210> 1108
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 1108
Asp Thr Met Gly His His Ser
<210> 1109
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 1109
Asp Thr Met Gly His Pro Ser
                5
 1
<210> 1110
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 1110
Asp Thr Met Gly His Thr Ser
                 5
<210> 1111
<211> 7
<212> PRT
<213> Artificial Sequence
<223> Amino acid sequence derived from Murine monoclonal
```

antibody and further modified by amino acid substitutions <400> 1111 Asp Thr Met Gly His Asp Ser 5 <210> 1112 <211> 7 <212> PRT <213> Artificial Sequence <220> <223> Amino acid sequence derived from Murine monoclonal antibody and further modified by amino acid substitutions <400> 1112 Asp Thr Met Gly Gln Ala Ser 1 <210> 1113 <211> 7 <212> PRT <213> Artificial Sequence <220> <223> Amino acid sequence derived from Murine monoclonal antibody and further modified by amino acid substitutions <400> 1113 Asp Thr Met Gly Gln Ser Ser 1 5 <210> 1114 <211> 7 <212> PRT <213> Artificial Sequence <223> Amino acid sequence derived from Murine monoclonal antibody and further modified by amino acid substitutions <400> 1114 Asp Thr Met Gly Gln Lys Ser <210> 1115 <211> 7 <212> PRT <213> Artificial Sequence <220> <223> Amino acid sequence derived from Murine monoclonal antibody and further modified by amino acid substitutions <400> 1115

304

```
Asp Thr Met Gly Gln Arg Ser
 <210> 1116
 <211> 7
 <212> PRT
 <213> Artificial Sequence
 <220>
<223> Amino acid sequence derived from Murine monoclonal
       antibody and further modified by amino acid
       substitutions
<400> 1116
Asp Thr Met Gly Gln His Ser
<210> 1117
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 1117
Asp Thr Met Gly Gln Pro Ser
                 5
<210> 1118
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 1118
Asp Thr Met Gly Gln Thr Ser
                 5
<210> 1119
<211> 7
<212> PRT
<213> Artificial Sequence
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 1119
Asp Thr Met Gly Gln Asp Ser
1
<210> 1120
<211> 7
```

```
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 1120
Asp Thr Met Arg Leu Ser Ser
                  5
<210> 1121
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 1121
Asp Thr Met Arg Leu Lys Ser
<210> 1122
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 1122
Asp Thr Met Arg Leu Arg Ser
<210> 1123
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 1123
Asp Thr Met Arg Leu His Ser
<210> 1124
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
```

antibody and further modified by amino acid substitutions <400> 1124 Asp Thr Met Arg Leu Pro Ser 5 <210> 1125 <211> 7 <212> PRT <213> Artificial Sequence <223> Amino acid sequence derived from Murine monoclonal antibody and further modified by amino acid substitutions <400> 1125 Asp Thr Met Arg Leu Thr Ser 1 5 <210> 1126 <211> 7 <212> PRT <213> Artificial Sequence <223> Amino acid sequence derived from Murine monoclonal antibody and further modified by amino acid substitutions <400> 1126 Asp Thr Met Arg Leu Asp Ser 1 5 <210> 1127 <211> 7 <212> PRT <213> Artificial Sequence <220> <223> Amino acid sequence derived from Murine monoclonal antibody and further modified by amino acid substitutions <400> 1127 Asp Thr Met Arg His Ala Ser 1 5 <210> 1128 <211> 7 <212> PRT <213> Artificial Sequence <223> Amino acid sequence derived from Murine monoclonal antibody and further modified by amino acid substitutions <400> 1128 Asp Thr Met Arg His Ser Ser

```
1
                 5
<210> 1129
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 1129
Asp Thr Met Arg His Lys Ser
<210> 1130
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 1130
Asp Thr Met Arg His Arg Ser
                5
<210> 1131
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 1131
Asp Thr Met Arg His His Ser
<210> 1132
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 1132
Asp Thr Met Arg His Pro Ser
                 5
 1
<210> 1133
<211> 7
<212> PRT
```

```
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 1133
Asp Thr Met Arg His Thr Ser
<210> 1134
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 1134
Asp Thr Met Arg His Asp Ser
                 5
<210> 1135
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 1135
Asp Thr Met Arg Gln Ala Ser
                 5
<210> 1136
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 1136
Asp Thr Met Arg Gln Ser Ser
1
                 5
<210> 1137
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
```

antibody and further modified by amino acid substitutions <400> 1137 Asp Thr Met Arg Gln Lys Ser <210> 1138 <211> 7 <212> PRT <213> Artificial Sequence <220> <223> Amino acid sequence derived from Murine monoclonal antibody and further modified by amino acid substitutions <400> 1138 Asp Thr Met Arg Gln Arg Ser 1 <210> 1139 <211> 7 <212> PRT <213> Artificial Sequence <223> Amino acid sequence derived from Murine monoclonal antibody and further modified by amino acid substitutions <400> 1139 Asp Thr Met Arg Gln His Ser 1 <210> 1140 <211> 7 <212> PRT <213> Artificial Sequence <223> Amino acid sequence derived from Murine monoclonal antibody and further modified by amino acid substitutions <400> 1140 Asp Thr Met Arg Gln Pro Ser 1 5 <210> 1141 <211> 7 <212> PRT <213> Artificial Sequence <223> Amino acid sequence derived from Murine monoclonal antibody and further modified by amino acid substitutions <400> 1141

Asp Thr Met Arg Gln Thr Ser

```
<210> 1142
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 1142
Asp Thr Met Arg Gln Asp Ser
<210> 1143
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 1143
Asp Thr Met Tyr Leu Lys Ser
<210> 1144
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 1144
Asp Thr Met Tyr Leu Arg Ser
<210> 1145
<211> 7
<212> PRT
<213> Artificial Sequence
<223> Amino acid sequence derived from Murine monoclonal
       antibody and further modified by amino acid
       substitutions
 <400> 1145
 Asp Thr Met Tyr Leu His Ser
 <210> 1146
 <211> 7
 <212> PRT
```

```
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 1146
Asp Thr Met Tyr Leu Pro Ser
                 5
<210> 1147
<211> 7
<212> PRT
<213> Artificial Sequence
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 1147
Asp Thr Met Tyr Leu Thr Ser
                 5
<210> 1148
<211> 7
<212> PRT
<213> Artificial Sequence
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 1148
Asp Thr Met Tyr Leu Asp Ser
                 5
1
<210> 1149
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 1149
Asp Thr Met Tyr His Ala Ser
              5
<210> 1150
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
```

substitutions <400> 1150 Asp Thr Met Tyr His Ser Ser <210> 1151 <211> 7 <212> PRT <213> Artificial Sequence <220> <223> Amino acid sequence derived from Murine monoclonal antibody and further modified by amino acid substitutions <400> 1151 Asp Thr Met Tyr His Lys Ser <210> 1152 <211> 7 <212> PRT <213> Artificial Sequence <220> <223> Amino acid sequence derived from Murine monoclonal antibody and further modified by amino acid substitutions <400> 1152 Asp Thr Met Tyr His Arg Ser <210> 1153 <211> 7 <212> PRT <213> Artificial Sequence <220> <223> Amino acid sequence derived from Murine monoclonal antibody and further modified by amino acid substitutions <400> 1153 Asp Thr Met Tyr His His Ser <210> 1154 <211> 7 <212> PRT <213> Artificial Sequence <223> Amino acid sequence derived from Murine monoclonal

antibody and further modified by amino acid

<400> 1154 Asp Thr Met Tyr His Pro Ser

substitutions

```
5
1.
<210> 1155
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
     antibody and further modified by amino acid
     substitutions
<400> 1155
Asp Thr Met Tyr His Thr Ser
<210> 1156
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 1156
Asp Thr Met Tyr His Asp Ser
<210> 1157
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 1157
Asp Thr Met Tyr Gln Lys Ser
 1
<210> 1158
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 1158
Asp Thr Met Tyr Gln Arg Ser
                 5
<210> 1159
<211> 7
<212> PRT
```

```
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 1159
Asp Thr Met Tyr Gln His Ser
<210> 1160
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 1160
Asp Thr Met Tyr Gln Pro Ser
<210> 1161
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 1161
Asp Thr Met Tyr Gln Thr Ser
<210> 1162
<211> 7
<212> PRT
<213> Artificial Sequence
<223> Amino acid sequence derived from Murine monoclonal
       antibody and further modified by amino acid
       substitutions
<400> 1162
Asp Thr Met Tyr Gln Asp Ser
<210> 1163
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
```

```
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 1163
Asp Thr Met Phe Leu Ala Ser
<210> 1164
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 1164
Asp Thr Met Phe Leu Ser Ser
<210> 1165
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 1165
Asp Thr Met Phe Leu Lys Ser
<210> 1166
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 1166
Asp Thr Met Phe Leu Arg Ser
<210> 1167
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 1167
```

```
Asp Thr Met Phe Leu His Ser
<210> 1168
<211> 7
<212> PRT
<213> Artificial Sequence
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 1168
Asp Thr Met Phe Leu Pro Ser
<210> 1169
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 1169
Asp Thr Met Phe Leu Thr Ser
<210> 1170
<211> 7
<212> PRT
<213> Artificial Sequence
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 1170
Asp Thr Met Phe Leu Asp Ser
<210> 1171
<211> 7
<212> PRT
<213> Artificial Sequence
 <223> Amino acid sequence derived from Murine monoclonal
       antibody and further modified by amino acid
       substitutions
 <400> 1171
 Asp Thr Met Phe His Ala Ser
                 5
 <210> 1172
 <211> 7
```

```
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 1172
Asp Thr Met Phe His Ser Ser
<210> 1173
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 1173
Asp Thr Met Phe His Lys Ser
                5
<210> 1174
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 1174
Asp Thr Met Phe His Arg Ser
<210> 1175
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 1175
Asp Thr Met Phe His His Ser
                 5
<210> 1176
<211> 7
<212> PRT
<213> Artificial Sequence
<223> Amino acid sequence derived from Murine monoclonal
```

antibody and further modified by amino acid substitutions <400> 1176 Asp Thr Met Phe His Pro Ser <210> 1177 <211> 7 <212> PRT <213> Artificial Sequence <220> <223> Amino acid sequence derived from Murine monoclonal antibody and further modified by amino acid substitutions <400> 1177 Asp Thr Met Phe His Thr Ser <210> 1178 <211> 7 <212> PRT <213> Artificial Sequence <220> <223> Amino acid sequence derived from Murine monoclonal antibody and further modified by amino acid substitutions <400> 1178 Asp Thr Met Phe His Asp Ser <210> 1179 <211> 7 <212> PRT <213> Artificial Sequence <220> <223> Amino acid sequence derived from Murine monoclonal antibody and further modified by amino acid substitutions <400> 1179 Asp Thr Met Phe Gln Ala Ser 1 <210> 1180 <211> 7 <212> PRT <213> Artificial Sequence <220> <223> Amino acid sequence derived from Murine monoclonal antibody and further modified by amino acid substitutions <400> 1180

```
Asp Thr Met Phe Gln Ser Ser
<210> 1181
<211> 7
<212> PRT
<213> Artificial Sequence
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 1181
Asp Thr Met Phe Gln Lys Ser
<210> 1182
<211> 7
<212> PRT
<213> Artificial Sequence
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 1182
Asp Thr Met Phe Gln Arg Ser
<210> 1183
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 1183
Asp Thr Met Phe Gln His Ser
<210> 1184
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
       antibody and further modified by amino acid
       substitutions
<400> 1184
Asp Thr Met Phe Gln Pro Ser
                 5
<210> 1185
<211> 7
```

```
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 1185
Asp Thr Met Phe Gln Thr Ser
<210> 1186
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 1186
Asp Thr Met Phe Gln Asp Ser
                5
<210> 1187
<211> 7
<212> PRT
<213> Artificial Sequence
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
       substitutions
 <400> 1187
 Asp Thr Met Leu Leu Ala Ser
                 5
 <210> 1188
 <211> 7
 <212> PRT
 <213> Artificial Sequence
 <223> Amino acid sequence derived from Murine monoclonal
       antibody and further modified by amino acid
       substitutions
 <400> 1188
 Asp Thr Met Leu Leu Ser Ser
 <210> 1189
 <211> 7
 <212> PRT
 <213> Artificial Sequence
 <223> Amino acid sequence derived from Murine monoclonal
```

substitutions <400> 1189 Asp Thr Met Leu Leu Lys Ser 5 <210> 1190 <211> 7 <212> PRT <213> Artificial Sequence <223> Amino acid sequence derived from Murine monoclonal antibody and further modified by amino acid substitutions <400> 1190 Asp Thr Met Leu Leu Arg Ser <210> 1191 <211> 7 <212> PRT <213> Artificial Sequence <223> Amino acid sequence derived from Murine monoclonal antibody and further modified by amino acid substitutions <400> 1191 Asp Thr Met Leu Leu His Ser <210> 1192 <211> 7 <212> PRT <213> Artificial Sequence <223> Amino acid sequence derived from Murine monoclonal antibody and further modified by amino acid substitutions <400> 1192 Asp Thr Met Leu Leu Pro Ser <210> 1193 <211> 7 <212> PRT <213> Artificial Sequence <220> <223> Amino acid sequence derived from Murine monoclonal antibody and further modified by amino acid substitutions <400> 1193 Asp Thr Met Leu Leu Thr Ser

antibody and further modified by amino acid

```
1
<210> 1194
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 1194
Asp Thr Met Leu Leu Asp Ser
<210> 1195
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 1195
Asp Thr Met Leu His Ala Ser
                5
<210> 1196
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 1196
Asp Thr Met Leu His Ser Ser
                5
<210> 1197
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 1197
Asp Thr Met Leu His Lys Ser
                 5
 1
<210> 1198
<211> 7
<212> PRT
```

```
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
     antibody and further modified by amino acid
     substitutions
<400> 1198
Asp Thr Met Leu His Arg Ser
                5
<210> 1199
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 1199
Asp Thr Met Leu His His Ser
<210> 1200
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 1200
Asp Thr Met Leu His Pro Ser
<210> 1201
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 1201
Asp Thr Met Leu His Thr Ser
                  5
<210> 1202
<211> 7
<212> PRT
<213> Artificial Sequence
<223> Amino acid sequence derived from Murine monoclonal
```

antibody and further modified by amino acid substitutions
1202

<400> 1202 Asp Thr Met Leu His Asp Ser 5 <210> 1203 <211> 7 <212> PRT <213> Artificial Sequence <223> Amino acid sequence derived from Murine monoclonal antibody and further modified by amino acid substitutions <400> 1203 Asp Thr Met Leu Gln Ala Ser <210> 1204 <211> 7 <212> PRT <213> Artificial Sequence <223> Amino acid sequence derived from Murine monoclonal antibody and further modified by amino acid substitutions <400> 1204 Asp Thr Met Leu Gln Ser Ser <210> 1205 <211> 7 <212> PRT <213> Artificial Sequence <220> <223> Amino acid sequence derived from Murine monoclonal antibody and further modified by amino acid substitutions <400> 1205 Asp Thr Met Leu Gln Lys Ser <210> 1206 <211> 7 <212> PRT <213> Artificial Sequence <220> <223> Amino acid sequence derived from Murine monoclonal antibody and further modified by amino acid

<400> 1206 Asp Thr Met Leu Gln Arg Ser

substitutions

```
5
1
<210> 1207
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 1207
Asp Thr Met Leu Gln His Ser
<210> 1208
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 1208
Asp Thr Met Leu Gln Pro Ser
<210> 1209
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 1209
Asp Thr Met Leu Gln Thr Ser
<210> 1210
<211> 7
<212> PRT
<213> Artificial Sequence
 <223> Amino acid sequence derived from Murine monoclonal
       antibody and further modified by amino acid
       substitutions
 <400> 1210
 Asp Thr Met Leu Gln Asp Ser
 <210> 1211
 <211> 7
 <212> PRT
```

```
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 1211
Asp Thr Lys Lys Leu Ala Ser
               5
<210> 1212
<211> 7
<212> PRT
<213> Artificial Sequence
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 1212
Asp Thr Lys Lys Leu Ser Ser
<210> 1213
<211> 7
<212> PRT
<213> Artificial Sequence
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 1213
Asp Thr Lys Lys Leu Lys Ser
<210> 1214
<211> 7
<212> PRT
<213> Artificial Sequence
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 1214
Asp Thr Lys Lys Leu Arg Ser
<210> 1215
<211> 7
<212> PRT
<213> Artificial Sequence
<223> Amino acid sequence derived from Murine monoclonal
       antibody and further modified by amino acid
```

substitutions

```
<400> 1215
Asp Thr Lys Lys Leu His Ser
<210> 1216
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 1216
Asp Thr Lys Lys Leu Pro Ser
<210> 1217
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 1217
Asp Thr Lys Lys Leu Thr Ser
<210> 1218
<211> 7
<212> PRT
<213> Artificial Sequence
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 1218
Asp Thr Lys Lys Leu Asp Ser
<210> 1219
<211> 7
<212> PRT
<213> Artificial Sequence
 <223> Amino acid sequence derived from Murine monoclonal
       antibody and further modified by amino acid
      substitutions
 <400> 1219
 Asp Thr Lys Lys His Ala Ser
```

```
<210> 1220
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
     antibody and further modified by amino acid
     substitutions
<400> 1220
Asp Thr Lys Lys His Ser Ser
<210> 1221
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 1221
Asp Thr Lys Lys His Lys Ser
<210> 1222
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 1222
Asp Thr Lys Lys His Arg Ser
<210> 1223
<211> 7
<212> PRT
<213> Artificial Sequence
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 1223
Asp Thr Lys Lys His His Ser
                 5
<210> 1224
<211> 7
<212> PRT
```

```
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 1224
Asp Thr Lys Lys His Pro Ser
<210> 1225
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 1225
Asp Thr Lys Lys His Thr Ser
<210> 1226
<211> 7
<212> PRT
<213> Artificial Sequence
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 1226
Asp Thr Lys Lys His Asp Ser
                5
<210> 1227
<211> 7
<212> PRT
<213> Artificial Sequence
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
       substitutions
<400> 1227
 Asp Thr Lys Lys Gln Ala Ser
 <210> 1228
 <211> 7
 <212> PRT
 <213> Artificial Sequence
 <220>
```

```
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 1228
Asp Thr Lys Lys Gln Ser Ser
<210> 1229
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 1229
Asp Thr Lys Lys Gln Lys Ser
<210> 1230
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 1230
Asp Thr Lys Lys Gln Arg Ser
<210> 1231
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 1231
Asp Thr Lys Lys Gln His Ser
<210> 1232
<211> 7
<212> PRT
<213> Artificial Sequence
<22.0>
<223> Amino acid sequence derived from Murine monoclonal
       antibody and further modified by amino acid
       substitutions
<400> 1232
```

```
Asp Thr Lys Lys Gln Pro Ser
1 5
<210> 1233
<211> 7
<212> PRT
<213> Artificial Sequence
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 1233
Asp Thr Lys Lys Gln Thr Ser
<210> 1234
<211> 7
<212> PRT
<213> Artificial Sequence
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 1234
Asp Thr Lys Lys Gln Asp Ser
<210> 1235
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 1235
Asp Thr Lys Gly Leu Ala Ser
<210> 1236
<211> 7
<212> PRT
<213> Artificial Sequence
 <220>
 <223> Amino acid sequence derived from Murine monoclonal
       antibody and further modified by amino acid
       substitutions
 <400> 1236
 Asp Thr Lys Gly Leu Ser Ser
 <210> 1237
 <211> 7
```

```
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 1237
Asp Thr Lys Gly Leu Lys Ser
<210> 1238
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 1238
Asp Thr Lys Gly Leu Arg Ser
<210> 1239
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 1239
Asp Thr Lys Gly Leu His Ser
<210> 1240
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 1240
Asp Thr Lys Gly Leu Pro Ser
 1
<210> 1241
 <211> 7
 <212> PRT
 <213> Artificial Sequence
 <223> Amino acid sequence derived from Murine monoclonal
```

antibody and further modified by amino acid substitutions
<400> 1241
Asp Thr Lys Gly Leu Thr Ser
1 5

<210> 1242

<211> 7

<212> PRT

<213> Artificial Sequence

<220>

<223> Amino acid sequence derived from Murine monoclonal antibody and further modified by amino acid substitutions

<400> 1242

Asp Thr Lys Gly Leu Asp Ser

<210> 1243

<211> 7

<212> PRT

<213> Artificial Sequence

<220>

<223> Amino acid sequence derived from Murine monoclonal antibody and further modified by amino acid substitutions

<400> 1243

Asp Thr Lys Gly His Ala Ser

Τ.

<210> 1244 <211> 7

<211> / <212> PRT

<213> Artificial Sequence

<220>

<223> Amino acid sequence derived from Murine monoclonal antibody and further modified by amino acid substitutions

<400> 1244

Asp Thr Lys Gly His Ser Ser

.

<210> 1245

<211> 7

<212> PRT

<213> Artificial Sequence

<220>

<223> Amino acid sequence derived from Murine monoclonal antibody and further modified by amino acid substitutions

<400> 1245

```
Asp Thr Lys Gly His Lys Ser
<210> 1246
<211> 7
<212> PRT
<213> Artificial Sequence
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 1246
Asp Thr Lys Gly His Arg Ser
<210> 1247
<211> 7
<212> PRT
<213> Artificial Sequence
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 1247
Asp Thr Lys Gly His His Ser
<210> 1248
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 1248
Asp Thr Lys Gly His Pro Ser
                 5
<210> 1249
 <211> 7
 <212> PRT
 <213> Artificial Sequence
 <220>
 <223> Amino acid sequence derived from Murine monoclonal
       antibody and further modified by amino acid
       substitutions
 <400> 1249
 Asp Thr Lys Gly His Thr Ser
                 5
 <210> 1250
 <211> 7
```

```
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
     antibody and further modified by amino acid
     substitutions
<400> 1250
Asp Thr Lys Gly His Asp Ser
<210> 1251
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 1251
Asp Thr Lys Gly Gln Ala Ser
<210> 1252
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 1252
Asp Thr Lys Gly Gln Ser Ser
                5
<210> 1253
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 1253
Asp Thr Lys Gly Gln Lys Ser
<210> 1254
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
```

antibody and further modified by amino acid substitutions

```
<400> 1254
Asp Thr Lys Gly Gln Arg Ser
           5
<210> 1255
<211> 7
<212> PRT
<213> Artificial Sequence
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 1255
Asp Thr Lys Gly Gln His Ser
<210> 1256
<211> 7
<212> PRT
<213> Artificial Sequence
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 1256
Asp Thr Lys Gly Gln Pro Ser
<210> 1257
<211> 7
<212> PRT
<213> Artificial Sequence
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 1257
Asp Thr Lys Gly Gln Thr Ser
<210> 1258
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
       antibody and further modified by amino acid
       substitutions
 <400> 1258
Asp Thr Lys Gly Gln Asp Ser
```

```
5
1
<210> 1259
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 1259
Asp Thr Lys Arg Leu Ala Ser
<210> 1260
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 1260
Asp Thr Lys Arg Leu Ser Ser
<210> 1261
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 1261
Asp Thr Lys Arg Leu Lys Ser
<210> 1262
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
       antibody and further modified by amino acid
       substitutions
<400> 1262
Asp Thr Lys Arg Leu Arg Ser
<210> 1263
<211> 7
```

```
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 1263
Asp Thr Lys Arg Leu His Ser
<210> 1264
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 1264
Asp Thr Lys Arg Leu Pro Ser
<210> 1265
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 1265
Asp Thr Lys Arg Leu Thr Ser
<210> 1266
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
       antibody and further modified by amino acid
       substitutions
 <400> 1266
Asp Thr Lys Arg Leu Asp Ser
<210> 1267
 <211> 7
 <212> PRT
 <213> Artificial Sequence
 <220>
 <223> Amino acid sequence derived from Murine monoclonal
```

antibody and further modified by amino acid substitutions <400> 1267 Asp Thr Lys Arg His Ala Ser

<210> 1268

<211> 7 <212> PRT

<213> Artificial Sequence

<220>

<223> Amino acid sequence derived from Murine monoclonal antibody and further modified by amino acid substitutions

<400> 1268 Asp Thr Lys Arg His Ser Ser 1 5

<210> 1269 <211> 7

<212> PRT <213> Artificial Sequence

<220>

<223> Amino acid sequence derived from Murine monoclonal antibody and further modified by amino acid substitutions

<400> 1269 Asp Thr Lys Arg His Lys Ser 1 5

<210> 1270 <211> 7

<212> PRT

<213> Artificial Sequence

(220>

<223> Amino acid sequence derived from Murine monoclonal antibody and further modified by amino acid substitutions

<400> 1270 Asp Thr Lys Arg His Arg Ser 1 5

<210> 1271 <211> 7

<211> /

<213> Artificial Sequence

<220N

<223> Amino acid sequence derived from Murine monoclonal antibody and further modified by amino acid substitutions

<400> 1271

Asp Thr Lys Arg His His Ser

```
5
1
<210> 1272
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 1272
Asp Thr Lys Arg His Pro Ser
<210> 1273
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 1273
Asp Thr Lys Arg His Thr Ser
<210> 1274
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 1274
Asp Thr Lys Arg His Asp Ser
<210> 1275
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
 <223> Amino acid sequence derived from Murine monoclonal
       antibody and further modified by amino acid
       substitutions
 <400> 1275
 Asp Thr Lys Arg Gln Ala Ser
  1
 <210> 1276
 <211> 7
 <212> PRT
```

```
<213> Artificial Sequence
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
     substitutions
<400> 1276
Asp Thr Lys Arg Gln Ser Ser
     5
<210> 1277
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
     antibody and further modified by amino acid
     substitutions
<400> 1277
Asp Thr Lys Arg Gln Lys Ser
1 5
<210> 1278
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 1278
Asp Thr Lys Arg Gln Arg Ser
1
         5
<210> 1279
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 1279
Asp Thr Lys Arg Gln His Ser
<210> 1280
<211> 7
<212> PRT
<213> Artificial Sequence
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
```

substitutions <400> 1280 Asp Thr Lys Arg Gln Pro Ser 1 <210> 1281 <211> 7 <212> PRT <213> Artificial Sequence <220> <223> Amino acid sequence derived from Murine monoclonal antibody and further modified by amino acid substitutions <400> 1281 Asp Thr Lys Arg Gln Thr Ser <210> 1282 <211> 7 <212> PRT <213> Artificial Sequence <220> <223> Amino acid sequence derived from Murine monoclonal antibody and further modified by amino acid substitutions <400> 1282 Asp Thr Lys Arg Gln Asp Ser 5 <210> 1283 <211> 7 <212> PRT <213> Artificial Sequence <220> <223> Amino acid sequence derived from Murine monoclonal antibody and further modified by amino acid substitutions <400> 1283 Asp Thr Lys Tyr Leu Ala Ser <210> 1284 <211> 7 <212> PRT <213> Artificial Sequence <223> Amino acid sequence derived from Murine monoclonal antibody and further modified by amino acid substitutions <400> 1284 Asp Thr Lys Tyr Leu Ser Ser

```
1
<210> 1285
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 1285
Asp Thr Lys Tyr Leu Lys Ser
<210> 1286
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 1286
Asp Thr Lys Tyr Leu Arg Ser
<210> 1287
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 1287
Asp Thr Lys Tyr Leu His Ser
<210> 1288
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 1288
Asp Thr Lys Tyr Leu Pro Ser
<210> 1289
 <211> 7
```

```
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 1289
Asp Thr Lys Tyr Leu Thr Ser
<210> 1290
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 1290
Asp Thr Lys Tyr Leu Asp Ser
         5
<210> 1291
<211> 7
<212> PRT
<213> Artificial Sequence
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 1291
Asp Thr Lys Tyr His Ala Ser
               5
<210> 1292
<211> 7
<212> PRT
<213> Artificial Sequence
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 1292
Asp Thr Lys Tyr His Ser Ser
<210> 1293
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
```

```
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 1293
Asp Thr Lys Tyr His Lys Ser
1
                5
<210> 1294
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 1294
Asp Thr Lys Tyr His Arg Ser
<210> 1295
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 1295
Asp Thr Lys Tyr His His Ser
<210> 1296
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 1296
Asp Thr Lys Tyr His Pro Ser
<210> 1297
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 1297
```

```
Asp Thr Lys Tyr His Thr Ser
1
                5
<210> 1298
<211> 7
<212> PRT
<213> Artificial Sequence
<223> Amino acid sequence derived from Murine monoclonal
     antibody and further modified by amino acid
     substitutions
<400> 1298
Asp Thr Lys Tyr His Asp Ser
                5
<210> 1299
<211> 7
<212> PRT
<213> Artificial Sequence
<223> Amino acid sequence derived from Murine monoclonal
     antibody and further modified by amino acid
     substitutions
<400> 1299
Asp Thr Lys Tyr Gln Ala Ser
1
<210> 1300
<211> 7
<212> PRT
<213> Artificial Sequence
<223> Amino acid sequence derived from Murine monoclonal
     antibody and further modified by amino acid
     substitutions
<400> 1300
Asp Thr Lys Tyr Gln Ser Ser
1 5
<210> 1301
<211> 7
<212> PRT
<213> Artificial Sequence
<223> Amino acid sequence derived from Murine monoclonal
     antibody and further modified by amino acid
     substitutions
<400> 1301
Asp Thr Lys Tyr Gln Lys Ser
             5 ,
<210> 1302
<211> 7
```

```
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 1302
Asp Thr Lys Tyr Gln Arg Ser
                5
1
<210> 1303
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 1303
Asp Thr Lys Tyr Gln His Ser
1
                5
<210> 1304
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 1304
Asp Thr Lys Tyr Gln Pro Ser
1
                 5
<210> 1305
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 1305
Asp Thr Lys Tyr Gln Thr Ser
<210> 1306
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
```

antibody and further modified by amino acid substitutions <400> 1306 Asp Thr Lys Tyr Gln Asp Ser <210> 1307 <211> 7 <212> PRT <213> Artificial Sequence <220> <223> Amino acid sequence derived from Murine monoclonal antibody and further modified by amino acid substitutions <400> 1307 Asp Thr Lys Phe Leu Ala Ser 5 <210> 1308 <211> 7 <212> PRT <213> Artificial Sequence <220> <223> Amino acid sequence derived from Murine monoclonal antibody and further modified by amino acid substitutions <400> 1308 Asp Thr Lys Phe Leu Ser Ser 5 <210> 1309 <211> 7 <212> PRT <213> Artificial Sequence <220> <223> Amino acid sequence derived from Murine monoclonal antibody and further modified by amino acid substitutions <400> 1309 Asp Thr Lys Phe Leu Lys Ser <210> 1310 <211> 7 <212> PRT <213> Artificial Sequence <220> <223> Amino acid sequence derived from Murine monoclonal antibody and further modified by amino acid substitutions <400> 1310

```
Asp Thr Lys Phe Leu Arg Ser
<210> 1311
<211> 7
<212> PRT
<213> Artificial Sequence
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 1311
Asp Thr Lys Phe Leu His Ser
<210> 1312
<211> 7
<212> PRT
<213> Artificial Sequence
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 1312
Asp Thr Lys Phe Leu Pro Ser
<210> 1313
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
       antibody and further modified by amino acid
       substitutions
<400> 1313
Asp Thr Lys Phe Leu Thr Ser
<210> 1314
 <211> 7
 <212> PRT
 <213> Artificial Sequence
 <220>
 <223> Amino acid sequence derived from Murine monoclonal
       antibody and further modified by amino acid
       substitutions
 <400> 1314
 Asp Thr Lys Phe Leu Asp Ser
 <210> 1315
 <211> 7
```

```
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 1315
Asp Thr Lys Phe His Ala Ser
<210> 1316
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 1316
Asp Thr Lys Phe His Ser Ser
<210> 1317
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 1317
Asp Thr Lys Phe His Lys Ser
<210> 1318
<211> 7
<212> PRT
<213> Artificial Sequence
 <220>
 <223> Amino acid sequence derived from Murine monoclonal
       antibody and further modified by amino acid
       substitutions
 <400> 1318
 Asp Thr Lys Phe His Arg Ser
 <210> 1319
 <211> 7
 <212> PRT
 <213> Artificial Sequence
 <223> Amino acid sequence derived from Murine monoclonal
```

antibody and further modified by amino acid substitutions

```
<400> 1319
Asp Thr Lys Phe His His Ser
<210> 1320
<211> 7
<212> PRT
<213> Artificial Sequence
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 1320
Asp Thr Lys Phe His Pro Ser
<210> 1321
<211> 7
<212> PRT
<213> Artificial Sequence
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 1321
Asp Thr Lys Phe His Thr Ser
<210> 1322
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
       substitutions
 <400> 1322
 Asp Thr Lys Phe His Asp Ser
                 5
 <210> 1323
 <211> 7
 <212> PRT
 <213> Artificial Sequence
 <223> Amino acid sequence derived from Murine monoclonal
       antibody and further modified by amino acid
       substitutions
 <400> 1323
 Asp Thr Lys Phe Gln Ala Ser
```

```
1
                 5
<210> 1324
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 1324
Asp Thr Lys Phe Gln Ser Ser
<210> 1325
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 1325
Asp Thr Lys Phe Gln Lys Ser
<210> 1326
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
       antibody and further modified by amino acid
       substitutions
 <400> 1326
 Asp Thr Lys Phe Gln Arg Ser
 <210> 1327
 <211> 7
 <212> PRT
 <213> Artificial Sequence
 <223> Amino acid sequence derived from Murine monoclonal
       antibody and further modified by amino acid
       substitutions
 <400> 1327
 Asp Thr Lys Phe Gln His Ser
 <210> 1328
 <211> 7
```

```
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 1328
Asp Thr Lys Phe Gln Pro Ser
                5
<210> 1329
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 1329
Asp Thr Lys Phe Gln Thr Ser
<210> 1330
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 1330
Asp Thr Lys Phe Gln Asp Ser
<210> 1331
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
       antibody and further modified by amino acid
       substitutions
<400> 1331
Asp Thr Lys Leu Leu Ala Ser
<210> 1332
 <211> 7
 <212> PRT
 <213> Artificial Sequence
 <223> Amino acid sequence derived from Murine monoclonal
```

antibody and further modified by amino acid substitutions

```
<400> 1332
Asp Thr Lys Leu Leu Ser Ser
<210> 1333
<211> 7
<212> PRT
<213> Artificial Sequence
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 1333
Asp Thr Lys Leu Leu Lys Ser
<210> 1334
<211> 7
<212> PRT
<213> Artificial Sequence
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 1334
Asp Thr Lys Leu Leu Arg Ser
<210> 1335
<211> 7
<212> PRT
<213> Artificial Sequence
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 1335
Asp Thr Lys Leu Leu His Ser
<210> 1336
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
 <223> Amino acid sequence derived from Murine monoclonal
       antibody and further modified by amino acid
       substitutions
 <400> 1336
 Asp Thr Lys Leu Leu Pro Ser
```

355

```
5
<210> 1337
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 1337
Asp Thr Lys Leu Leu Thr Ser
<210> 1338
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 1338
Asp Thr Lys Leu Leu Asp Ser
<210> 1339
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 1339
Asp Thr Lys Leu His Ala Ser
                5
<210> 1340
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
 <223> Amino acid sequence derived from Murine monoclonal
       antibody and further modified by amino acid
       substitutions
 <400> 1340
 Asp Thr Lys Leu His Ser Ser
 <210> 1341
 <211> 7
 <212> PRT
```

```
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 1341
Asp Thr Lys Leu His Lys Ser
<210> 1342
<211> 7
<212> PRT
<213> Artificial Sequence
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 1342
Asp Thr Lys Leu His Arg Ser
<210> 1343
<211> 7
<212> PRT
<213> Artificial Sequence
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
       substitutions
<400> 1343
Asp Thr Lys Leu His His Ser
<210> 1344
<211> 7
<212> PRT
<213> Artificial Sequence
 <223> Amino acid sequence derived from Murine monoclonal
       antibody and further modified by amino acid
       substitutions
 <400> 1344
 Asp Thr Lys Leu His Pro Ser
                 5
 <210> 1345
 <211> 7
 <212> PRT
 <213> Artificial Sequence
 <223> Amino acid sequence derived from Murine monoclonal
       antibody and further modified by amino acid
```

substitutions

```
<400> 1345
Asp Thr Lys Leu His Thr Ser
        5
<210> 1346
<211> 7
<212> PRT
<213> Artificial Sequence
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 1346
Asp Thr Lys Leu His Asp Ser
<210> 1347
<211> 7
<212> PRT
<213> Artificial Sequence
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 1347
Asp Thr Lys Leu Gln Ala Ser
<210> 1348
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 1348
Asp Thr Lys Leu Gln Ser Ser
<210> 1349
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
       antibody and further modified by amino acid
       substitutions
<400> 1349
Asp Thr Lys Leu Gln Lys Ser
```

```
5
1
<210> 1350
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 1350
Asp Thr Lys Leu Gln Arg Ser
<210> 1351
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 1351
Asp Thr Lys Leu Gln His Ser
<210> 1352
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
       substitutions
<400> 1352
Asp Thr Lys Leu Gln Pro Ser
 1
 <210> 1353
 <211> 7
 <212> PRT
 <213> Artificial Sequence
 <223> Amino acid sequence derived from Murine monoclonal
       antibody and further modified by amino acid
       substitutions
 <400> 1353
 Asp Thr Lys Leu Gln Thr Ser
 <210> 1354
```

```
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 1354
Asp Thr Lys Leu Gln Asp Ser
<210> 1355
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 1355
Asp Thr Leu Lys Leu Ser Ser
<210> 1356
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 1356
Asp Thr Leu Lys Leu Lys Ser
<210> 1357
<211> 7
<212> PRT
<213> Artificial Sequence
 <223> Amino acid sequence derived from Murine monoclonal
       antibody and further modified by amino acid
       substitutions
 <400> 1357
 Asp Thr Leu Lys Leu Arg Ser
 <210> 1358
 <211> 7
 <212> PRT
 <213> Artificial Sequence
```

```
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 1358
Asp Thr Leu Lys Leu His Ser
       5
<210> 1359
<211> 7
<212> PRT
<213> Artificial Sequence
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 1359
Asp Thr Leu Lys Leu Pro Ser
       5
<210> 1360
<211> 7
<212> PRT
<213> Artificial Sequence
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 1360
Asp Thr Leu Lys Leu Thr Ser
<210> 1361
<211> 7
<212> PRT
<213> Artificial Sequence
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
 <400> 1361
Asp Thr Leu Lys His Ala Ser
 <210> 1362
 <211> 7
 <212> PRT
 <213> Artificial Sequence
 <220>
 <223> Amino acid sequence derived from Murine monoclonal
       antibody and further modified by amino acid
       substitutions
```

```
<400> 1362
Asp Thr Leu Lys His Ser Ser
           5
<210> 1363
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
     antibody and further modified by amino acid
      substitutions
<400> 1363
Asp Thr Leu Lys His Lys Ser
<210> 1364
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 1364
Asp Thr Leu Lys His Arg Ser
<210> 1365
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 1365
Asp Thr Leu Lys His His Ser
<210> 1366
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
 <223> Amino acid sequence derived from Murine monoclonal
       antibody and further modified by amino acid
      substitutions
 <400> 1366
 Asp Thr Leu Lys His Pro Ser
                 5
  1
 <210> 1367
```

```
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 1367
Asp Thr Leu Lys His Thr Ser
<210> 1368
<211> 7
<212> PRT
<213> Artificial Sequence
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 1368
Asp Thr Leu Lys His Asp Ser
<210> 1369
<211> 7
<212> PRT
<213> Artificial Sequence
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
       substitutions
<400> 1369
Asp Thr Leu Lys Gln Ala Ser
<210> 1370
 <211> 7
 <212> PRT
 <213> Artificial Sequence
 <223> Amino acid sequence derived from Murine monoclonal
       antibody and further modified by amino acid
       substitutions
 <400> 1370
 Asp Thr Leu Lys Gln Ser Ser
 <210> 1371
 <211> 7
 <212> PRT
 <213> Artificial Sequence
```

```
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 1371
Asp Thr Leu Lys Gln Lys Ser
<210> 1372
<211> 7
<212> PRT
<213> Artificial Sequence
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 1372
Asp Thr Leu Lys Gln Arg Ser
<210> 1373
<211> 7
<212> PRT
<213> Artificial Sequence
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
       substitutions
<400> 1373
Asp Thr Leu Lys Gln His Ser
<210> 1374
 <211> 7
 <212> PRT
 <213> Artificial Sequence
 <223> Amino acid sequence derived from Murine monoclonal
       antibody and further modified by amino acid
       substitutions
 <400> 1374
 Asp Thr Leu Lys Gln Pro Ser
 <210> 1375
 <211> 7
 <212> PRT
 <213> Artificial Sequence
 <223> Amino acid sequence derived from Murine monoclonal
       antibody and further modified by amino acid
       substitutions
```

```
<400> 1375
Asp Thr Leu Lys Gln Thr Ser
<210> 1376
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 1376
Asp Thr Leu Lys Gln Asp Ser
                5
<210> 1377
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 1377
Asp Thr Leu Gly Leu Ala Ser
<210> 1378
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 1378
Asp Thr Leu Gly Leu Ser Ser
<210> 1379
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
 <223> Amino acid sequence derived from Murine monoclonal
       antibody and further modified by amino acid
       substitutions
 <400> 1379
 Asp Thr Leu Gly Leu Lys Ser
  1
 <210> 1380
```

```
<211> 7
<212> PRT
<213> Artificial Sequence
<223> Amino acid sequence derived from Murine monoclonal
     antibody and further modified by amino acid
     substitutions
<400> 1380
Asp Thr Leu Gly Leu Arg Ser
<210> 1381
<211> 7
<212> PRT
<213> Artificial Sequence
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 1381
Asp Thr Leu Gly Leu His Ser
<210> 1382
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 1382
Asp Thr Leu Gly Leu Pro Ser
 <210> 1383
 <211> 7
 <212> PRT
 <213> Artificial Sequence
 <223> Amino acid sequence derived from Murine monoclonal
       antibody and further modified by amino acid
       substitutions
 <400> 1383
 Asp Thr Leu Gly Leu Thr Ser
 <210> 1384
 <211> 7
 <212> PRT
 <213> Artificial Sequence
 <220>
```

```
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 1384
Asp Thr Leu Gly Leu Asp Ser
<210> 1385
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 1385
Asp Thr Leu Gly His Ala Ser
                 5
<210> 1386
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 1386
Asp Thr Leu Gly His Ser Ser
<210> 1387
<211> 7
<212> PRT
<213> Artificial Sequence
 <223> Amino acid sequence derived from Murine monoclonal
       antibody and further modified by amino acid
       substitutions
 <400> 1387
 Asp Thr Leu Gly His Lys Ser
 <210> 1388
 <211> 7
 <212> PRT
 <213> Artificial Sequence
 <220>
 <223> Amino acid sequence derived from Murine monoclonal
       antibody and further modified by amino acid
       substitutions
 <400> 1388
```

```
Asp Thr Leu Gly His Arg Ser
<210> 1389
<211> 7
<212> PRT
<213> Artificial Sequence
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 1389
Asp Thr Leu Gly His His Ser
            5
<210> 1390
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 1390
Asp Thr Leu Gly His Pro Ser
<210> 1391
<211> 7
 <212> PRT
 <213> Artificial Sequence
 <220>
 <223> Amino acid sequence derived from Murine monoclonal
       antibody and further modified by amino acid
       substitutions
 <400> 1391
 Asp Thr Leu Gly His Thr Ser
 <210> 1392
 <211> 7
 <212> PRT
 <213> Artificial Sequence
 <220>
 <223> Amino acid sequence derived from Murine monoclonal
       antibody and further modified by amino acid
       substitutions
 <400> 1392
 Asp Thr Leu Gly His Asp Ser
                  5
  1
 <210> 1393
```

```
<211> 7
<212> PRT
<213> Artificial Sequence
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 1393
Asp Thr Leu Gly Gln Ala Ser
        5
<210> 1394
<211> 7
<212> PRT
<213> Artificial Sequence
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 1394
Asp Thr Leu Gly Gln Ser Ser
<210> 1395
<211> 7
<212> PRT
<213> Artificial Sequence
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
       substitutions
 <400> 1395
 Asp Thr Leu Gly Gln Lys Ser
 <210> 1396
 <211> 7
 <212> PRT
 <213> Artificial Sequence
 <220>
 <223> Amino acid sequence derived from Murine monoclonal
       antibody and further modified by amino acid
       substitutions
 <400> 1396
 Asp Thr Leu Gly Gln Arg Ser
 <210> 1397
 <211> 7
 <212> PRT
 <213> Artificial Sequence
 <220>
```

```
<223> Amino acid sequence derived from Murine monoclonal
     antibody and further modified by amino acid
     substitutions
<400> 1397
Asp Thr Leu Gly Gln His Ser
               5
<210> 1398
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
     antibody and further modified by amino acid
   substitutions
<400> 1398
Asp Thr Leu Gly Gln Pro Ser
<210> 1399
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 1399
Asp Thr Leu Gly Gln Thr Ser
                5
<210> 1400
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
 <400> 1400
Asp Thr Leu Gly Gln Asp Ser
 <210> 1401
 <211> 7
 <212> PRT
 <213> Artificial Sequence
 <223> Amino acid sequence derived from Murine monoclonal
       antibody and further modified by amino acid
       substitutions
 <400> 1401
```

```
Asp Thr Leu Arg Leu Ala Ser
<210> 1402
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 1402
Asp Thr Leu Arg Leu Ser Ser
<210> 1403
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 1403
Asp Thr Leu Arg Leu Lys Ser
 1 5
<210> 1404
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 1404
Asp Thr Leu Arg Leu Arg Ser
         5
<210> 1405
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 1405
Asp Thr Leu Arg Leu His Ser
<210> 1406
<211> 7
```

```
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 1406
Asp Thr Leu Arg Leu Pro Ser
                5
<210> 1407
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 1407
Asp Thr Leu Arg Leu Thr Ser
<210> 1408
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 1408
Asp Thr Leu Arg Leu Asp Ser
<210> 1409
<211> 7
 <212> PRT
 <213> Artificial Sequence
 <220>
 <223> Amino acid sequence derived from Murine monoclonal
       antibody and further modified by amino acid
       substitutions
 <400> 1409
 Asp Thr Leu Arg His Ala Ser
 <210> 1410
 <211> 7
 <212> PRT
 <213> Artificial Sequence
 <220>
```

```
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 1410
Asp Thr Leu Arg His Ser Ser
<210> 1411
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 1411
Asp Thr Leu Arg His Lys Ser
<210> 1412
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 1412
Asp Thr Leu Arg His Arg Ser
<210> 1413
 <211> 7
 <212> PRT
 <213> Artificial Sequence
 <220>
 <223> Amino acid sequence derived from Murine monoclonal
       antibody and further modified by amino acid
       substitutions
 <400> 1413
 Asp Thr Leu Arg His His Ser
 <210> 1414
 <211> 7
 <212> PRT
 <213> Artificial Sequence
 <220>
 <223> Amino acid sequence derived from Murine monoclonal
       antibody and further modified by amino acid
       substitutions
 <400> 1414
```

```
Asp Thr Leu Arg His Pro Ser
<210> 1415
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 1415
Asp Thr Leu Arg His Thr Ser
<210> 1416
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 1416
Asp Thr Leu Arg His Asp Ser
                5
<210> 1417
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
       antibody and further modified by amino acid
       substitutions
<400> 1417
Asp Thr Leu Arg Gln Ala Ser
                 5
<210> 1418
<211> 7
 <212> PRT
 <213> Artificial Sequence
 <223> Amino acid sequence derived from Murine monoclonal
       antibody and further modified by amino acid
       substitutions
 <400> 1418
 Asp Thr Leu Arg Gln Ser Ser
  1
```

```
<210> 1419
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 1419
Asp Thr Leu Arg Gln Lys Ser
<210> 1420
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 1420
Asp Thr Leu Arg Gln Arg Ser
<210> 1421
<211> 7
<212> PRT
<213> Artificial Sequence
 <220>
 <223> Amino acid sequence derived from Murine monoclonal
       antibody and further modified by amino acid
       substitutions
 <400> 1421
 Asp Thr Leu Arg Gln His Ser
 <210> 1422
 <211> 7
 <212> PRT
 <213> Artificial Sequence
 <220>
 <223> Amino acid sequence derived from Murine monoclonal
       antibody and further modified by amino acid
       substitutions
 <400> 1422
 Asp Thr Leu Arg Gln Pro Ser
                   5
 <210> 1423
 <211> 7
 <212> PRT
 <213> Artificial Sequence
```

```
<220>
<223> Amino acid sequence derived from Murine monoclonal
     antibody and further modified by amino acid
     substitutions
<400> 1423
Asp Thr Leu Arg Gln Thr Ser
1
<210> 1424
<211> 7
<212> PRT
<213> Artificial Sequence
<223> Amino acid sequence derived from Murine monoclonal
     antibody and further modified by amino acid
     substitutions
<400> 1424
Asp Thr Leu Arg Gln Asp Ser
1
   5
<210> 1425
<211> 7
<212> PRT
<213> Artificial Sequence
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
     substitutions
<400> 1425
Asp Thr Leu Tyr Leu Ala Ser
1
                5
<210> 1426
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 1426
Asp Thr Leu Tyr Leu Ser Ser
     5
<210> 1427
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
```

```
<400> 1427
Asp Thr Leu Tyr Leu Lys Ser
<210> 1428
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 1428
Asp Thr Leu Tyr Leu Arg Ser
<210> 1429
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 1429
Asp Thr Leu Tyr Leu His Ser
<210> 1430
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 1430
Asp Thr Leu Tyr Leu Pro Ser
<210> 1431
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
     antibody and further modified by amino acid
     substitutions
<400> 1431
Asp Thr Leu Tyr Leu Thr Ser
1
<210> 1432
```

```
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 1432
Asp Thr Leu Tyr Leu Asp Ser
<210> 1433
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 1433
Asp Thr Leu Tyr His Ala Ser
<210> 1434
<211> 7
<212> PRT
<213> Artificial Sequence
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 1434
Asp Thr Leu Tyr His Ser Ser
<210> 1435
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
     antibody and further modified by amino acid
     substitutions
<400> 1435
Asp Thr Leu Tyr His Lys Ser
<210> 1436
<211> 7
<212> PRT
<213> Artificial Sequence
```

```
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 1436
Asp Thr Leu Tyr His Arg Ser
<210> 1437
<211> 7
<212> PRT
<213> Artificial Sequence
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 1437
Asp Thr Leu Tyr His His Ser
<210> 1438
<211> 7
<212> PRT
<213> Artificial Sequence
<223> Amino acid sequence derived from Murine monoclonal
     antibody and further modified by amino acid
     substitutions
<400> 1438
Asp Thr Leu Tyr His Pro Ser
<210> 1439
<211> 7
<212> PRT
<213> Artificial Sequence
<223> Amino acid sequence derived from Murine monoclonal
     antibody and further modified by amino acid
     substitutions
<400> 1439
Asp Thr Leu Tyr His Thr Ser
<210> 1440
<211> 7
<212> PRT
<213> Artificial Sequence
<223> Amino acid sequence derived from Murine monoclonal
     antibody and further modified by amino acid
     substitutions
```

```
<400> 1440
Asp Thr Leu Tyr His Asp Ser
<210> 1441
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
     antibody and further modified by amino acid
     substitutions
<400> 1441
Asp Thr Leu Tyr Gln Ala Ser
<210> 1442
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
     antibody and further modified by amino acid
     substitutions
<400> 1442
Asp Thr Leu Tyr Gln Ser Ser
<210> 1443
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
     antibody and further modified by amino acid
     substitutions
<400> 1443
Asp Thr Leu Tyr Gln Lys Ser
<210> 1444
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
     antibody and further modified by amino acid
     substitutions
<400> 1444
Asp Thr Leu Tyr Gln Arg Ser
<210> 1445
```

```
<211> 7
<212> PRT
<213> Artificial Sequence
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 1445
Asp Thr Leu Tyr Gln His Ser
<210> 1446
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 1446
Asp Thr Leu Tyr Gln Pro Ser
<210> 1447
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 1447
Asp Thr Leu Tyr Gln Thr Ser
<210> 1448
<211> 7
<212> PRT
<213> Artificial Sequence
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 1448
Asp Thr Leu Tyr Gln Asp Ser
<210> 1449
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
```

```
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 1449
Asp Thr Leu Phe Leu Ala Ser
<210> 1450
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 1450
Asp Thr Leu Phe Leu Ser Ser
<210> 1451
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 1451
Asp Thr Leu Phe Leu Lys Ser
<210> 1452
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 1452
Asp Thr Leu Phe Leu Arg Ser
<210> 1453
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 1453
```

```
Asp Thr Leu Phe Leu His Ser
<210> 1454
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 1454
Asp Thr Leu Phe Leu Pro Ser
          5
<210> 1455
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 1455 .
Asp Thr Leu Phe Leu Thr Ser
<210> 1456
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 1456
Asp Thr Leu Phe Leu Asp Ser
<210> 1457
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 1457
Asp Thr Leu Phe His Ala Ser
1
<210> 1458
```

```
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 1458
Asp Thr Leu Phe His Ser Ser
<210> 1459
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 1459
Asp Thr Leu Phe His Lys Ser
<210> 1460
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 1460
Asp Thr Leu Phe His Arg Ser
                5
<210> 1461
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 1461
Asp Thr Leu Phe His His Ser
                 5
<210> 1462
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
```

```
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 1462
Asp Thr Leu Phe His Pro Ser
<210> 1463
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 1463
Asp Thr Leu Phe His Thr Ser
<210> 1464
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 1464
Asp Thr Leu Phe His Asp Ser
<210> 1465
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 1465
Asp Thr Leu Phe Gln Ala Ser
<210> 1466
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 1466
```

385

```
Asp Thr Leu Phe Gln Ser Ser
               5
<210> 1467
<211> 7
<212> PRT
<213> Artificial Sequence
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 1467
Asp Thr Leu Phe Gln Lys Ser
<210> 1468
<211> 7
<212> PRT
<213> Artificial Sequence
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 1468
Asp Thr Leu Phe Gln Arg Ser
1
                5
<210> 1469
<211> 7
<212> PRT
<213> Artificial Sequence
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 1469
Asp Thr Leu Phe Gln His Ser
1
<210> 1470
<211> 7
<212> PRT
<213> Artificial Sequence
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 1470
Asp Thr Leu Phe Gln Pro Ser
1
<210> 1471
<211> 7
```

```
<212> PRT
<213> Artificial Sequence
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 1471
Asp Thr Leu Phe Gln Thr Ser
<210> 1472
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 1472
Asp Thr Leu Phe Gln Asp Ser
<210> 1473
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 1473
Asp Thr Leu Leu Leu Ser Ser
                5
<210> 1474
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 1474
Asp Thr Leu Leu Leu Lys Ser
                 5
<210> 1475
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
```

```
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 1475
Asp Thr Leu Leu Leu Arg Ser
<210> 1476
<211> 7
<212> PRT
<213> Artificial Sequence
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 1476
Asp Thr Leu Leu Leu His Ser
<210> 1477
<211> 7
<212> PRT
<213> Artificial Sequence
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 1477
Asp Thr Leu Leu Pro Ser
<210> 1478
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 1478
Asp Thr Leu Leu Leu Thr Ser
<210> 1479
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 1479
```

388

```
Asp Thr Leu Leu His Ala Ser
<210> 1480
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 1480
Asp Thr Leu Leu His Ser Ser
<210> 1481
<211> 7
<212> PRT
<213> Artificial Sequence
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 1481
Asp Thr Leu Leu His Lys Ser
<210> 1482
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 1482
Asp Thr Leu Leu His Arg Ser
<210> 1483
<211> 7
<212> PRT
<213> Artificial Sequence
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 1483
Asp Thr Leu Leu His His Ser
         5
```

```
<210> 1484
<211> 7
<212> PRT
<213> Artificial Sequence
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 1484
Asp Thr Leu Leu His Pro Ser
<210> 1485
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 1485
Asp Thr Leu Leu His Thr Ser
               5
<210> 1486
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 1486
Asp Thr Leu Leu His Asp Ser
<210> 1487
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 1487
Asp Thr Leu Leu Gln Ala Ser
<210> 1488
<211> 7
<212> PRT
<213> Artificial Sequence
```

```
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 1488
Asp Thr Leu Leu Gln Ser Ser
                5
<210> 1489
<211> 7
<212> PRT
<213> Artificial Sequence
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 1489
Asp Thr Leu Leu Gln Lys Ser
<210> 1490
<211> 7
<212> PRT
<213> Artificial Sequence
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 1490
Asp Thr Leu Leu Gln Arg Ser
1
                5
<210> 1491
<211> 7
<212> PRT
<213> Artificial Sequence
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 1491
Asp Thr Leu Leu Gln His Ser
                5
1
<210> 1492
<211> 7
<212> PRT
<213> Artificial Sequence
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
```

```
<400> 1492
Asp Thr Leu Leu Gln Pro Ser
<210> 1493
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 1493
Asp Thr Leu Leu Gln Thr Ser
<210> 1494
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 1494
Asp Thr Leu Leu Gln Asp Ser
<210> 1495
<211> 9
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 1495
Phe Gln Gly Ser Tyr Tyr Pro Phe Thr
<210> 1496
<211> 9
<212> PRT
<213> Artificial Sequence
<220>
<223> Amino acid sequence derived from Murine monoclonal
      antibody and further modified by amino acid
      substitutions
<400> 1496
Phe Gln Gly Ser Trp Tyr Pro Phe Thr
```